

**TREATMENT OF STUTTERING
AND OTHER COMMUNICATION DISORDERS
WITH NOREPINEPHRINE REUPTAKE INHIBITORS**

5

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention relates to the fields of pharmaceutical chemistry and central nervous system medicine. More specifically, the present invention relates to methods of
10 treating communication disorders, such as stuttering, in children, adolescents, and adults by administering selective norepinephrine reuptake inhibitors to patients in need of such treatment.

Description of Related Art

15 The *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition (DSM-IV) (1994), American Psychiatric Association, Washington, D.C., pp. 55-65, describes a number of communication disorders usually first diagnosed in infancy, childhood, or adolescence. These include stuttering, expressive language disorder, mixed receptive-expressive language disorder, phonological disorder, and communication
20 disorder not otherwise specified. Stuttering is perhaps the most well known of these disorders.

Stuttering is a speech disorder in which the normal flow of speech is disrupted by frequent repetitions or prolongations of speech sounds, syllables, or words, or by an individual's inability to start a word. The speech disruptions may be accompanied by
25 rapid eye blinks, tremors of the lips and/or jaw, or other struggle behaviors of the face or upper body that a person who stutters may use in an attempt to speak. Certain situations, such as speaking before a group of people or talking on the telephone, tend to make stuttering more severe, whereas other situations, such as singing or speaking alone, often improve fluency. Stuttering is also referred to as stammering, especially in England, and
30 by a broader term, dysfluent speech.

Characteristics of stuttering are described in Section 307.0 of the DSM-IV at pp. 63-65. While all individuals are dysfluent at times, the person who stutters is differentiated from someone with normal speech disfluencies by the kind and amount of dysfluencies.

- 5 Characteristics of stuttering include:
- Repetition of sounds, syllables, parts of words, whole words, and phrases
 - Prolongation, or stretching, of sounds or syllables
 - Tense pauses, hesitations, and/or no sound between words
 - Speech that occurs in spurts, as the client tries to initiate or maintain voice
 - 10 - Related behaviors, for example reactions that accompany stuttering such as tense muscles in the lips, jaw, and/or neck; tremor of the lips, jaw, and/or tongue during attempts to speak; foot tapping, eye blinks, head turns, etc. (to try to escape from the stuttering); etc. There are many related behaviors that can occur and vary from person to person.
 - 15 - Variability in stuttering behavior, depending on the speaking situation, the communication partner(s), and the speaking task. A person who stutters may experience more fluency in the speech-language pathologist's office than in a classroom or workplace. There may be no difficulty making a special dinner request at home, but extreme difficulty ordering a meal in a restaurant.
 - 20 Conversation with a spouse may be easier, and more fluent, than that with a boss. A person may be completely fluent when singing, but experience significant stuttering when talking on the telephone.
 - A feeling of loss of control. The person who stutters may experience sound and word fears, situational fears, anticipation of stuttering, embarrassment,
 - 25 and a sense of shame. Certain sounds or words may be avoided. One word may be substituted for another that is thought to be harder to say. Or, certain speaking situations may be avoided altogether. For example, a person who stutters may always wait for someone else to answer the phone. Or, he or she may walk around a store for an hour rather than ask sales staff where an item
 - 30 can be found. These reactions to stuttering occur in more advanced stages.

Repetitions and prolongations are essential features of stuttering. The presence of the other listed behaviors varies from person to person.

Developmental stuttering (DS), with or without associated psychiatric illness, is the most common form, and includes all cases with gradual onset in childhood that are not the result of acquired brain damage. Persistent developmental stuttering (PDS) is DS that has not undergone spontaneous or speech therapy-induced remission. Acquired stuttering, which is much rarer than DS, may occur in previously fluent individuals. This form may be neurogenic, resulting from brain damage associated with, for example, stroke, traumatic brain injury, Alzheimer's disease, renal dialysis, Parkinson's disease, and progressive supranuclear palsy (Heuer et al. (1996) *Ear Nose Throat J.* 75:161-168; Brazis et al. (1996) *Localization in Clinical Neurology*, Third Ed., Little, Brown and Company, Boston, MA, p.515).

Based on neuroimaging research data and the effectiveness of dopamine receptor antagonists in DS, this form of stuttering appears to have a hyperdopaminergic origin.

It is estimated that over three million Americans stutter. Stuttering affects individuals of all ages, but occurs most frequently in young children between the ages of 2 and 6 who are developing language. The prevalence of stuttering in prepubertal children is 1%, and drops to 0.8% in adolescence. The male-to-female ratio is approximately 3:1. Most children outgrow their stuttering, and it is estimated that less than 1 percent of adults stutter.

Family and twin studies strongly suggest a genetic factor in the etiology of stuttering. The risk of stuttering among first-degree biological relatives is more than three times that in the general population. About 10% of daughters, and 20% of sons, of men who stutter will also stutter.

There is at present no cure for stuttering. However, a variety of treatments are available that may improve stuttering to some degree. These include speech therapy to improve fluency and success in communication; parent education to restructure the child's speaking environment to reduce episodes of stuttering; and the use of interventions such as electronic devices or medications. Electronic devices which help an individual control fluency may be more of a bother than a help in most speaking situations, and are often

abandoned by individuals who stutter. Medications that affect brain function often have side effects that make them difficult to use for long-term treatment.

Many medications have been studied for use in treating stuttering. Evidence suggests that persons who stutter exhibit hypometabolism of the striatum and increased dopamine activity (Wu et al. (1995) *Neuroreport* 6:501-5; Wu et al. (1997) *Neuroreport* 8:767-70; Wu et al. (1997) In: Hulstijn W, Peters HRM, van Lieshout PHHM, eds. *Speech production: motor control, brain research and fluency disorders*. International Congress Series 1146. Amsterdam: Excerpta Medica 339-41). Drugs that boost dopamine levels exacerbate stuttering. Ritalin has a similar effect. Tricyclic antidepressants have proved ineffective, and in fact stuttering has been reported as an adverse event with the use of these compounds.

In contrast, the dopamine antagonist haloperidol has been shown in replicated, double-blind trials to reduce the symptoms of stuttering, leading to the hypothesis that D₂ receptor antagonists may be important in the treatment of developmental stuttering (J.P. Brady (1991) *Am. J. Psychiatry* 148:1309-16). Unfortunately, as this drug is not well tolerated by this patient population and carries substantial risk of extrapyramidal symptoms and tardive dyskinesia, it is not recommended for the treatment of stuttering.

In a recent small study, Maguire et al. ((2000) *J. Clin. Psychopharmacology* 20:479-482) demonstrated that the serotonin-dopamine antagonist risperidone may be effective in the treatment of developmental stuttering, and recommended further investigations of risperidone for this purpose. While extrapyramidal symptoms and akathisia were not found with the use of risperidone, sedation was common, and some participants developed transient sexual and menstrual cycle side effects that resolved with discontinuance of the medication, or with a reduction in dose. These side effects are thought to be due to the elevation of the hormone prolactin by risperidone (haloperidol also raises prolactin levels in some patients).

Paroxetine and Sertraline, selective serotonin reuptake inhibitors, are also used for the treatment of stuttering, but cause a number of undesirable side effects.

Newer medications more narrowly target dopamine receptors. Olanzapine (Zyprexa), has been used successfully to treat developmental and acquired stuttering in children, adolescents, and adults (Lavid et al. (1999) *Annals of Clinical Psychiatry* 11(4):

233-236; Lavid et al. (2000) Presented at the annual meeting of the American Psychiatric Association, Chicago IL, 2000). Side effects were mostly limited to slight weight gain and drowsiness.

The present invention addresses the need in the art for improved treatments for
5 stuttering that are both safe and effective.

SUMMARY OF THE INVENTION

Accordingly, in a first aspect, the present invention provides a method of treating
stuttering or another communication disorder, comprising administering to a patient in
10 need of such treatment an effective amount of a selective norepinephrine reuptake
inhibitor. The selective norepinephrine reuptake inhibitor can be, but is not limited to,
any of the compounds disclosed herein.

In another aspect, the present invention provides the use of a selective
norepinephrine reuptake inhibitor, such as any of the compounds disclosed herein, or
15 other selective norepinephrine reuptake inhibitors, for the manufacture of a medicament
for the treatment of stuttering or another communication disorder.

Further scope of the applicability of the present invention will become apparent
from the detailed description provided below. However, it should be understood that the
detailed description and specific examples, while indicating preferred embodiments of the
20 present invention, are given by way of illustration only since various changes and
modifications within the spirit and scope of the invention will become apparent to those
skilled in the art from this detailed description.

DETAILED DESCRIPTION OF THE INVENTION

25 The following detailed description of the invention is provided to aid those skilled
in the in practicing the present invention. Even so, the following detailed description
should not be construed to unduly limit the present invention as modifications and
variations in the embodiments discussed herein can be made by those of ordinary skill in
the art without departing from the spirit or scope of the present inventive discovery.

30 The contents of each of the references cited herein are herein incorporated by
reference in their entirety.

Costa and Kroll ((2000) *Canadian Medical Association Journal* 162(13):1849-1855) disclose definitions and features of the various types of stuttering:

Developmental Stuttering – Stuttering with gradual onset in childhood as a disturbance in the normal fluency and time patterning of speech.

5 **Persistent Developmental Stuttering** – Developmental stuttering that has not undergone spontaneous or speech therapy-induced remission.

10 **Acquired Stuttering** – Stuttering that occurs more or less abruptly in previously fluent individuals. Neurogenic acquired stuttering results from brain damage, involves repetitions, prolongations, and blocks, and is not associated with grimaces, eye-blinking, or social anxiety. Psychogenic acquired stuttering begins suddenly after emotional trauma, and involves repetition of initial or stressed syllables, indifference toward dysfluency, dysfluency that never fluctuates, and persistence of normal eye contact.

Diagnostic criteria for stuttering set forth in the DSM-IV (p. 65) are:

15 **Diagnostic criteria for 307.0 Stuttering**

A. Disturbance in the normal fluency and time patterning of speech (inappropriate for the individual's age), characterized by frequent occurrences of one or more of the following:

- 20 (1) sound and syllable repetitions
 (2) sound prolongations
 (3) interjections
 (4) broken words (e.g., pauses within a word)
 (5) audible or silent blocking (filled or unfilled pauses in speech)
25 (6) circumlocutions (word substitutions to avoid problematic words)
 (7) words produced with an excess of physical tension
 (8) monosyllabic whole-word repetitions (e.g., "I-I-I-I see him")

30 B. The disturbance in fluency interferes with academic or occupational achievement or with social communication.

C. If a speech-motor or sensory deficit is present, the speech difficulties are in excess of those usually associated with these problems.

Any of the types of stuttering described above, whether presenting alone, or
5 comorbidly with cluttering, phonological disorder, expressive or mixed receptive-
expressive language disorder, attention-deficit hyperactivity disorder (ADHD), or mental
retardation, can be treated or prevented by the methods of the present invention. Patients
will receive benefit from the use of norepinephrine reuptake inhibitors in the amelioration
of the symptoms of stuttering regardless of whether a comorbid condition is present.
10 Patients suffering from both stuttering and a comorbid condition, for example attention-
deficit hyperactivity disorder, will receive benefit in the amelioration of symptoms of
both conditions via the methods of the present invention. Therefore, in addition to
methods for treating stuttering presenting alone, the present invention encompasses
methods of treating stuttering comorbid with any of the conditions listed immediately
15 above, comprising administering to a patient in need of treatment of both stuttering and
such comorbid condition an effective amount of a selective norepinephrine reuptake
inhibitor.

For clarification, cluttering is characterized by unpredictable, fast, and jerky
outpourings of words and phrases, including slurred or omitted syllables and improper
20 phrasing and pauses (Kaplan and Sadock (1998) *Synopsis of Psychiatry*, Eighth Ed.,
Williams and Wilkins, Baltimore, MD, pp.1175-7). Phonological disorder (DSM-IV
section 315.39) involves, in part, failure to produce developmentally expected speech
sounds appropriate for the individual's age and dialect, and can involve errors in sound
production, use, representation, or organization. Expressive language disorder (DSM-IV
25 section 315.31) involves, in part, an impairment in expressive language development as
demonstrated by scores on standardized individually administered measures of expressive
language development substantially below those obtained from standardized measures of
both nonverbal intellectual capacity and receptive language development. Mixed
receptive-expressive language disorder (DSM-IV section 315.32) involves, in part, an
30 impairment in both receptive and expressive language development as demonstrated by
scores on standardized individually administered measures of both receptive and

expressive language development that are substantially below those obtained from standardized measures of nonverbal intellectual capacity. Attention-deficit hyperactivity disorder (DSM-IV sections 314.00 and 314.01) involves, in part, a persistent pattern of inattention and/or hyperactivity-impulsivity that is more frequent and severe than is typically observed in individuals at a comparable level of development.

Specific diagnostic criteria in the DSM-IV for expressive language disorder, mixed receptive-expressive language disorder, phonological disorder, communication disorder not otherwise specified, as well as for attention-deficit hyperactivity disorder, each of which can exist comorbidly with developmental stuttering, are as follows:

Diagnostic criteria for 315.31 Expressive Language Disorder

A. The scores obtained from standardized individually administered measures of expressive language development are substantially below those obtained from standardized measures of both nonverbal intellectual capacity and receptive language development. The disturbance may be manifest clinically by symptoms that include having a markedly limited vocabulary, making errors in tense, or having difficulty recalling words or producing sentences with developmentally appropriate length or complexity.

B. The difficulties with expressive language interfere with academic or occupational achievement or with social communication.

C. Criteria are not met for Mixed Receptive-Expressive Language Disorder or a Pervasive Developmental Disorder.

D. If Mental Retardation, a speech-motor or sensory deficit, or environmental deprivation is present, the language difficulties are in excess of those usually associated with these problems.

Diagnostic criteria for 315.32 Mixed Receptive-Expressive Language Disorder

- 5 A. The scores obtained from a battery of standardized individually administered measures of both receptive and expressive language development are substantially below those obtained from standardized measures of nonverbal intellectual capacity. Symptoms include those for Expressive Language Disorder as well as difficulty understanding words, sentences, or specific types of words, such as spatial terms.
- 10 B. The difficulties with receptive and expressive language significantly interfere with academic or occupational achievement or with social communication.
- C. Criteria are not met for a Pervasive Developmental Disorder.
- 15 D. If Mental Retardation, a speech-motor or sensory deficit, or environmental deprivation is present, the language difficulties are in excess of those usually associated with these problems.

Diagnostic criteria for 315.39 Phonological Disorder

- 20 A. Failure to use developmentally expected speech sounds that are appropriate for age and dialect (e.g., errors in sound production, use, representation, or organization such as, but not limited to, substitutions of one sound for another [use of /t/ for target /k/ sound] or omissions of sounds such
- 25 as final consonants).
- B. The difficulties in speech sound production interfere with academic or occupational achievement or with social communication.

C. If Mental Retardation, a speech-motor or sensory deficit, or environmental deprivation is present, the speech difficulties are in excess of those usually associated with these problems.

5 **307.9 Communication Disorder Not Otherwise Specified**

This category is for disorders in communication that do not meet criteria for any specific Communication Disorder; for example, a voice disorder (i.e., an abnormality of vocal pitch, loudness, quality, tone, or resonance).

10 **Diagnostic criteria for Attention-Deficit/Hyperactivity Disorder**

A. Either (1) or (2):

(1) six (or more) of the following symptoms of **inattention** have persisted for at least 6 months to a degree that is maladaptive and inconsistent with developmental level:

15

Inattention

(a) often fails to give close attention to details or makes careless mistakes in schoolwork, work, or other activities

(b) often has difficulty sustaining attention in tasks or play activities

20

(c) often does not seem to listen when spoken to directly

(d) often does not follow through on instructions and fails to finish schoolwork, chores, or duties in the workplace (not due to oppositional behavior or failure to understand instructions)

(e) often has difficulty organizing tasks and activities

25

(f) often avoids, dislikes, or is reluctant to engage in tasks that require sustained mental effort (such as schoolwork or homework)

(g) often loses things necessary for tasks or activities (e.g., toys, school assignments, pencils, books, or tools)

(h) is often easily distracted by extraneous stimuli

30

(i) is often forgetful in daily activities

(2) six (or more) of the following symptoms of **hyperactivity-impulsivity** have persisted for at least 6 months to a degree that is maladaptive and inconsistent with developmental level:

5

Hyperactivity

(a) often fidgets with hands or feet or squirms in seat

(b) often leaves seat in classroom or in other situations in which remaining seated is expected

10

(c) often runs about or climbs excessively in situations in which it is inappropriate (in adolescents or adults, may be limited to subjective feelings of restlessness)

(d) often has difficulty playing or engaging in leisure activities quietly

(e) is often "on the go" or often acts as if "driven by a motor"

(f) often talks excessively

15

Impulsivity

(g) often blurts out answers before questions have been completed

(h) often has difficulty awaiting turn

20

(i) often interrupts or intrudes on others (e.g., butts into conversations or games)

B. Some hyperactive-impulsive or inattentive symptoms that caused impairment were present before age 7 years.

25

C. Some impairment from the symptoms is present in two or more settings (e.g., at school [or work] and at home).

D. There must be clear evidence of clinically significant impairment in social, academic, or occupational functioning.

30

E. The symptoms do not occur exclusively during the course of a Pervasive Developmental Disorder, Schizophrenia, or other Psychotic Disorder and are not better accounted for by another mental disorder (e.g., Mood Disorder, Anxiety Disorder, Dissociative Disorder, or a Personality Disorder).

5

Code based on type:

314.01 Attention-Deficit/Hyperactivity Disorder, Combined Type: if both Criteria A1 and A2 are met for the past 6 months

10

314.00 Attention-Deficit/Hyperactivity Disorder, Predominantly Inattentive Type: if Criterion A1 is met but Criterion A2 is not met for the past 6 months

15

314.01 Attention-Deficit/Hyperactivity Disorder, Predominantly Hyperactive-Impulsive Type: if Criterion A2 is met but Criterion A1 is not met for the past 6 months

The methods of the present invention are effective in the treatment of patients who are children, adolescents, or adults, and there is no significant difference in the symptoms or the details of the manner of treatment among patients of different ages. In general terms, for purposes of the present invention, a child is considered to be a patient below the age of puberty, an adolescent is considered to be a patient from the age of puberty up to about 18 years of age, and an adult is considered to be a patient of 18 years or older.

25

Norepinephrine Reuptake Inhibitors Useful in the Present Invention

To the best of the inventor's knowledge, norepinephrine reuptake inhibitors have not been employed to treat stuttering.

Many compounds, including those discussed at length below, are selective norepinephrine reuptake inhibitors, and no doubt many more will be identified in the future. Practice of the present invention encompasses the use of norepinephrine reuptake

30

inhibitors that exhibit 50% effective concentrations of about 1000 nM or less in the protocol described by Wong et al. (1985) *Drug Development Research*, 6:397. Preferred norepinephrine reuptake inhibitors useful in the methods of the present invention are those that are selective for the inhibition of norepinephrine reuptake relative to their ability to act as direct agonists or antagonists at other receptors. Preferably, the compounds useful in the methods of the present invention are selective for the inhibition of norepinephrine reuptake relative to direct agonist or antagonist activity at other receptors by a factor of at least ten, and even more preferably by a factor of at least one hundred.


10 Norepinephrine reuptake inhibitors useful in the methods of the present invention include, but are not limited to:

1. Atomoxetine (formerly known as tomoxetine), (R)-(-)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine, is usually administered as the hydrochloride salt. Atomoxetine was first disclosed in U.S. Patent No. 4,314,081. The term "atomoxetine" will be used here to refer to any acid addition salt or the free base of the molecule. See, for example, Gehlert et al. (1993) *Neuroscience Letters* 157:203-206, for a discussion of atomoxetine's activity as a norepinephrine reuptake inhibitor;

2. Reboxetine (EdronaxTM; ProliftTM; VestraTM; NoreboxTM), 2-[α -(2-ethoxy)phenoxy-benzyl]morpholine, first disclosed in U.S. Patent 4,229,449 for the treatment of depression, is usually administered as the racemate. Reboxetine is a selective norepinephrine reuptake inhibitor. The term "reboxetine" as used herein refers to any acid addition salt or the free base of the molecule existing as the racemate or either enantiomer, i.e., (S,S)-reboxetine or (R,R)-reboxetine. The use of (S,S)-reboxetine as a preferred selective norepinephrine reuptake inhibitor is disclosed in PCT International Publication No. WO 01/01973.

3. Compounds of formula I:

14




(I)

wherein X is C₁-C₄ alkylthio, and Y is C₁-C₂ alkyl or a pharmaceutically acceptable salt thereof. The compounds of formula I have been described in U.S. Patent No. 5,281,624, and in Gehlert et al. (1995) *Life Sciences*, 55(22):1915-1920. These compounds are disclosed as being inhibitors of norepinephrine reuptake in the brain. It should be noted that these compounds exist as stereoisomers, and accordingly include not only the racemates, but also the isolated individual isomers as well as mixtures of the individual isomers. For example, the compounds of formula I include the following exemplary species:

N-ethyl-3-phenyl-3-(2-methylthiophenoxy)propyl-amine benzoate;
 (R)-N-methyl-3-phenyl-3-(2-propylthiophenoxy)-propylamine hydrochloride;
 (S)-N-ethyl-3-phenyl-3-(2-butylthiophenoxy)propyl-amine;
 N-methyl-3-phenyl-3-(2-ethylthiophenoxy)propyl-amine malonate;
 (S)-N-methyl-3-phenyl-3-(2-tert-butylthiophenoxy)-propylamine naphthalene-2-sulfonate; and
 (R)-N-methyl-3-(2-methylthiophenoxy)-3-phenyl-propylamine.

4. A compound of formula (IA)



(IA)

wherein n is 1, 2 or 3; R₁ is C₂-C₁₀alkyl, C₂-C₁₀alkenyl, C₃-C₈cycloalkyl or C₄-C₁₀cycloalkylalkyl, wherein one C-C bond within any cycloalkyl moiety is optionally substituted by an O-C or C=C bond and wherein each group is optionally substituted with from 1 to 7 halogen substituents and/or with from 1 to 3 substituents each independently selected from hydroxy, cyano, C₁-C₄alkyl and C₁-C₄alkoxy; R₂ is H, C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0, 1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl), or together with R₃ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy); R₃ is H, C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0, 1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl), or together with R₂ or R₄ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy); R₄ is H, C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0, 1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or

-CO₂(C₁-C₄alkyl), or together with R₃ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy); R₅ is H, C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms) or halogen; R₆ is H, C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms) or halogen; R₇ is H or C₁-C₄alkyl; R₈ is H or C₁-C₄alkyl; R₉ is H, halogen, hydroxy, cyano, C₁-C₄alkyl or C₁-C₄alkoxy; and R₁₀ is H, halogen, hydroxy, cyano, C₁-C₄alkyl or C₁-C₄alkoxy; or a pharmaceutically acceptable salt thereof, with the proviso that the compound N-ethyl-N-benzyl-4-piperidinamine is excluded.

With respect to compounds of formula (IA), the term "C₂-C₁₀alkyl" means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 2 to 10 carbon atoms.


With respect to compounds of formula (IA), the term "C₂-C₁₀alkenyl" means a monovalent unsubstituted unsaturated straight-chain or branched-chain hydrocarbon radical having from 2 to 10 carbon atoms and containing at least one carbon-carbon double bond.

With respect to compounds of formula (IA), the term "C₃-C₈cycloalkyl" means a monovalent unsubstituted saturated cyclic hydrocarbon radical having from 3 to 8 carbon atoms.

With respect to compounds of formula (IA), the term "C₄-C₁₀cycloalkylalkyl" means a monovalent unsubstituted saturated cyclic hydrocarbon radical having from 3 to 9 carbon atoms linked to the point of substitution by a divalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having at least 1 carbon atom.

With respect to compounds of formula (IA), the phrase "wherein one C-C bond within any cycloalkyl moiety is optionally substituted by an O-C or C=C bond" means that either (i) any two adjacent carbon atoms within a cycloalkyl ring may be linked by a double bond rather than a single bond (with the number of substituents on each carbon atom being reduced accordingly), or that (ii) one of any two adjacent C atoms within a

cycloalkyl ring (and any substituents thereon) may be replaced by an oxygen atom. Examples of R1 groups encompassed by this phrase include but are not limited to:



With respect to compounds of formula (IA), the term “halo” or “halogen” means
 5 F, Cl, Br or I.

With respect to compounds of formula (IA), the term “C₁-C₄alkoxy” means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 4 carbon atoms linked to the point of substitution by an O atom.

With respect to compounds of formula (IA), the term “phenoxy” means a
 10 monovalent unsubstituted phenyl radical linked to the point of substitution by an O atom.

With respect to compounds of formula (IA), in the above definitions, similar terms specifying different numbers of C atoms take an analogous meaning.

Preferred compounds of formula (IA) are those wherein n is 1 or 2. More preferably, n is 1.

15 Preferred compounds of formula (IA) are those wherein R7 is H or methyl. More preferably R7 is H.

Preferred compounds of formula (IA) are those wherein R8 is H.

Preferred compounds of formula (IA) are those wherein R9 is H or fluoro. More preferably, R9 is H.

20 Preferred compounds of formula (IA) are those wherein R10 is H or fluoro. More preferably, R10 is H.

Preferred compounds of formula (IA) are those wherein R1 is C₂-C₆alkyl, C₂-C₆alkenyl, C₃-C₆cycloalkyl or C₄-C₇cycloalkylalkyl, each of which is optionally substituted with from 1 to 3 halogen atoms or a methoxy radical. More preferably, R1 is
 25 C₂-C₆alkyl (optionally substituted with from 1 to 3 halogen atoms or a methoxy radical), C₂-C₆alkenyl, C₃-C₆cycloalkyl or C₄-C₇cycloalkylalkyl. Suitable C₂-C₆alkyl groups (optionally substituted with from 1 to 3 halogen atoms or a methoxy radical) include, for example, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, n-pentyl, 3-methylbutyl, 1,2-dimethylpropyl, 1-ethylpropyl, 3,3-dimethylbutyl, 2-ethylbutyl, 3,3,3-trifluoropropyl,

4,4,4-trifluorobutyl and 2-methoxyethyl. Suitable C₂-C₆alkenyl groups include, for example, 2-methyl-2-propenyl. Suitable C₃-C₆cycloalkyl groups include, for example, cyclopentyl. Suitable C₄-C₇cycloalkylalkyl groups include, for example, cyclohexylmethyl or cyclopropylmethyl.

- 5 Preferred compounds of formula (IA) are those wherein R₁ is a C₂-C₁₀alkyl group optionally substituted with from 1 to 7 halogen substituents and/or with from 1 to 3 substituents each independently selected from hydroxy, cyano and C₁-C₄alkoxy. More preferably, R₁ is a C₂-C₁₀alkyl group optionally substituted with from 1 to 3 substituents each independently selected from halogen, hydroxy and C₁-C₄alkoxy. More preferably
- 10 R₁ is C₂-C₆alkyl optionally substituted with from 1 to 3 halogen atoms or a methoxy radical. Still more preferably R₁ is C₂-C₆alkyl. Still more preferably, R₁ is selected from ethyl, n-propyl, isopropyl, n-butyl, isobutyl, n-pentyl, 3-methylbutyl, 1,2-dimethylpropyl, 1-ethylpropyl, 3,3-dimethylbutyl and 2-ethylbutyl. Most preferably R₁ is selected from n-propyl, n-butyl and isobutyl.
- 15 Preferred compounds of formula (IA) are those wherein R₂ is H, C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-
- 20 C₄alkyl and C₁-C₄alkoxy) or phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), or together with R₃ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy). More preferably, R₂ is H, C₁-C₂alkyl (optionally substituted with from 1 to 5 halogen
- 25 atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0 or 2 (optionally substituted with from 1 to 5 halogen atoms), C₁-C₂alkoxy (optionally substituted with from 1 to 5 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy) or phenoxy

(optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy), or together with R₃ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy). Still more preferably, R₂ is H, methyl, trifluoromethyl, methylthio, tert-butylthio, trifluoromethylthio, methylsulfonyl, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy, cyano, fluoro, chloro, bromo, phenyl or phenoxy, or together with R₃ forms a further benzene ring.

Preferred compounds of formula (IA) are those wherein R₂ is not H. More preferably, R₂ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), or together with R₃ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy). More preferably, R₂ is C₁-C₂alkyl (optionally substituted with from 1 to 5 halogen atoms), C₁-C₂alkyl-S(O)_x- wherein x is 0 or 2 (optionally substituted with from 1 to 5 halogen atoms), C₁-C₂alkoxy (optionally substituted with from 1 to 5 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy) or phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy), or together with R₃ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy). Still more preferably, R₂ is methyl, trifluoromethyl, methylthio, tert-butylthio, trifluoromethylthio, methylsulfonyl, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy, cyano, fluoro, chloro, bromo, phenyl or phenoxy, or together with R₃ forms a further benzene ring.

Preferred compounds of formula (IA) are those wherein R₃ is H, C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S- (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl), or together with R₂ or R₄ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy). More preferably, R₃ is H, C₁-C₂alkyl (optionally substituted with from 1 to 5 halogen atoms), C₁-C₂alkyl-S- (optionally substituted with from 1 to 5 halogen atoms), C₁-C₂alkoxy (optionally substituted with from 1 to 5 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy) or -CO₂(C₁-C₂alkyl), or together with R₂ or R₄ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy). Still more preferably, R₃ is H, methyl, trifluoromethyl, trifluoromethylthio, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy, cyano, fluoro, chloro, bromo, phenyl, phenoxy or CO₂CH₃, or together with R₂ or R₄ forms a further benzene ring.


Preferred compounds of formula (IA) are those wherein R₄ is H, C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S- (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), or -CO₂(C₁-C₄alkyl), or together with R₃ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-

C₄alkyl and C₁-C₄alkoxy). More preferably, R₄ is H, C₁-C₂alkyl (optionally substituted with from 1 to 5 halogen atoms), C₁-C₂alkyl-S- (optionally substituted with from 1 to 5 halogen atoms), C₁-C₂alkoxy (optionally substituted with from 1 to 5 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy), or -CO₂(C₁-C₂alkyl), or together with R₃ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₂alkyl and C₁-C₂alkoxy). Still more preferably, R₄ is H, methyl, trifluoromethyl, methylthio, methoxy, trifluoromethoxy, cyano, fluoro, chloro, phenyl or CO₂CH₃, or together with R₃ forms a further benzene ring.

Preferred compounds of formula (IA) are those wherein R₅ is H, C₁-C₄alkyl (optionally substituted with from 1 to 5 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 5 halogen atoms) or halogen. More preferably, R₅ is H, C₁-C₄alkyl, C₁-C₄alkoxy or halogen. Still more preferably, R₅ is H, methyl, methoxy, fluoro or chloro.

Preferred compounds of formula (IA) are those wherein R₆ is H, C₁-C₄alkyl (optionally substituted with from 1 to 5 halogen atoms) or halogen. More preferably, R₆ is H, C₁-C₄alkyl or halogen. Still more preferably, R₆ is H, methyl, fluoro or chloro.

Preferred compounds of formula (IA) are those wherein the group



is phenyl, 2-methylphenyl, 2-(trifluoromethyl)phenyl, 2-(methylthio)phenyl, 2-(tertbutylthio)phenyl, 2-(trifluoromethylthio)phenyl, 2-(methysulfonyl)phenyl, 2-methoxyphenyl, 2-ethoxyphenyl, 2-(difluoromethoxy)phenyl, 2-(trifluoromethoxy)phenyl, 2-cyanophenyl, 2-fluorophenyl, 2-chlorophenyl, 2-bromophenyl, 2-biphenyl, 2-phenoxyphenyl, 3-methylphenyl, 3-(trifluoromethyl)phenyl, 3-(trifluoromethylthio)phenyl, 3-methoxyphenyl, 3-ethoxyphenyl, 3-

(difluoromethoxy)phenyl, 3-(trifluoromethoxy)phenyl, 3-cyanophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-biphenyl, 3-phenoxyphenyl, 3-(methoxycarbonyl)phenyl, 4-methylphenyl, 4-(trifluoromethyl)phenyl, 4-(methylthio)phenyl, 4-methoxyphenyl, 4-(trifluoromethoxy)phenyl, 4-cyanophenyl, 4-fluorophenyl, 4-chlorophenyl, 4-biphenyl, 4-(methoxycarbonyl)phenyl, 2,3-dichlorophenyl, 2,4-dimethylphenyl, 2,4-bis(trifluoromethyl)phenyl, 2,4-dimethoxyphenyl, 2,4-difluorophenyl, 2,4-dichlorophenyl, 2,5-dimethylphenyl, 2,6-dimethylphenyl, 2,6-dichlorophenyl, 2-chloro-6-fluorophenyl, 2-fluoro-6-(trifluoromethyl)phenyl, 3,4-dichlorophenyl, 3,5-dimethylphenyl, 3,5-dimethoxyphenyl, 3,5-difluorophenyl, 3,5-dichlorophenyl, 3-fluoro-5-(trifluoromethyl)phenyl, 5-fluoro-2-(trifluoromethyl)phenyl, 5-fluoro-2-methoxyphenyl, 4-fluoro-2-(trifluoromethyl)phenyl, 1-naphthyl or 2-naphthyl.

A further embodiment provides a group (Group A) of compounds of formula (IA) above, wherein R₂, R₃, R₄, R₅ and R₆ are all H.

A further embodiment provides a group (Group B) of compounds of formula (IA) above, wherein one of R₂, R₃, R₄, R₅ and R₆ is not H and the others are H.

Compounds of Group B include those (Group B2) wherein R₃, R₄, R₅ and R₆ are all H and R₂ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl).

Compounds of Group B also include those (Group B3) wherein R₂, R₄, R₅ and R₆ are all H and R₃ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy

(optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl).

Compounds of Group B also include those (Group B4) wherein R₂, R₃, R₅ and R₆ are all H and R₄ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl).

A further embodiment provides a group (Group C) of compounds of formula (IA) above, wherein two of R₂, R₃, R₄, R₅ and R₆ are not H and the others are H.

Compounds of Group C include those (Group C2,3) wherein R₄, R₅ and R₆ are all H; R₂ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl), or together with R₃ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy); and R₃ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl), or together with R₂

forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy).

Compounds of Group C also include those (Group C2,4) wherein R₃, R₅ and R₆ are all H; R₂ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl); and R₄ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl).

Compounds of Group C also include those (Group C2,5) wherein R₃, R₄ and R₆ are all H; R₂ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl); and R₅ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms) or halogen.

Compounds of Group C also include those (Group C2,6) wherein R₃, R₄ and R₅ are all H; R₂ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-

C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted
5 with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl); and R₆ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms) or halogen.

Compounds of Group C also include those (Group C3,4) wherein R₂, R₅ and R₆
10 are all H; R₃ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted
15 with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl), or together with R₄ forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy); and R₄ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally
20 substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl), or together with R₃
25 forms a further benzene ring (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy).

Compounds of Group C also include those (Group C3,5) wherein R₂, R₄ and R₆ are all H; R₃ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-

C₄alkyl-S(O)_x- wherein x is 0,1 or 2 (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms), cyano, halogen, phenyl (optionally substituted with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy), phenoxy (optionally substituted
5 with from 1 to 3 substituents each independently selected from halogen, C₁-C₄alkyl and C₁-C₄alkoxy) or -CO₂(C₁-C₄alkyl); and R₅ is C₁-C₄alkyl (optionally substituted with from 1 to 7 halogen atoms), C₁-C₄alkoxy (optionally substituted with from 1 to 7 halogen atoms) or halogen.

For compounds of Formula (IA) falling within any one of groups A, B, B₂, B₃,
10 B₄, C, C_{2,3}, C_{2,4}, C_{2,5}, C_{2,6}, C_{3,4} and C_{3,5} described above, n is preferably 1 or 2, more preferably 1.

For compounds of Formula (IA) falling within any one of groups A, B, B₂, B₃, B₄, C, C_{2,3}, C_{2,4}, C_{2,5}, C_{2,6}, C_{3,4} and C_{3,5} described above, R₇ is preferably H or methyl, more preferably H.

15 For compounds of Formula (IA) falling within any one of groups A, B, B₂, B₃, B₄, C, C_{2,3}, C_{2,4}, C_{2,5}, C_{2,6}, C_{3,4} and C_{3,5} described above, R₈ is preferably H.

For compounds of Formula (IA) falling within any one of groups A, B, B₂, B₃, B₄, C, C_{2,3}, C_{2,4}, C_{2,5}, C_{2,6}, C_{3,4} and C_{3,5} described above, R₉ is preferably H or fluoro, more preferably H.


20 For compounds of Formula (IA) falling within any one of groups A, B, B₂, B₃, B₄, C, C_{2,3}, C_{2,4}, C_{2,5}, C_{2,6}, C_{3,4} and C_{3,5} described above, R₁₀ is preferably H or fluoro, more preferably H.

For compounds of Formula (IA) falling within any one of groups A, B, B₂, B₃, B₄, C, C_{2,3}, C_{2,4}, C_{2,5}, C_{2,6}, C_{3,4} and C_{3,5} described above, R₁ is preferably a C₂-
25 C₁₀alkyl group optionally substituted with from 1 to 7 halogen substituents and/or with from 1 to 3 substituents each independently selected from hydroxy, cyano and C₁-C₄alkoxy.

For compounds of Formula (IA) falling within any one of groups A, B, B₂, B₃, B₄, C, C_{2,3}, C_{2,4}, C_{2,5}, C_{2,6}, C_{3,4} and C_{3,5} described above, n is preferably 1, R₇, R₈,

R9 and R10 are preferably H and R1 is preferably a C₂-C₁₀alkyl group optionally substituted with from 1 to 7 halogen substituents and/or with from 1 to 3 substituents each independently selected from hydroxy, cyano and C₁-C₄alkoxy.

5. A compound of formula (IB)



(IB)

wherein Rx is H; Ry is H or C₁-C₄ alkyl; each Rz is independently H or C₁-C₄ alkyl; X represents O; Y represents OH or OR; R is C₁-C₄ alkyl; Ar₁ is a phenyl ring or a 5- or 6-membered heteroaryl ring each of which may be substituted with 1, 2, 3, 4 or 5 substituents (depending upon the number of available substitution positions) each independently selected from C₁-C₄ alkyl, O(C₁-C₄ alkyl), S(C₁-C₄ alkyl), halo, hydroxy, pyridyl, thiophenyl and phenyl optionally substituted with 1, 2, 3, 4 or 5 substituents each independently selected from halo, C₁-C₄ alkyl, or O(C₁-C₄ alkyl); and Ar₂ is a phenyl ring or a 5- or 6-membered heteroaryl ring each of which may be substituted with 1, 2, 3, 4 or 5 substituents (depending upon the number of available substitution positions) each independently selected from C₁-C₄ alkyl, O(C₁-C₄ alkyl) and halo; wherein each above-mentioned C₁-C₄ alkyl group is optionally substituted with one or more halo atoms; or a pharmaceutically acceptable salt thereof.

Preferred compounds of formula (IB) above are those wherein Ar₁ is phenyl, pyridyl, pyrimidyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, thiophenyl, furanyl, imidazolyl, triazolyl, oxadiazolyl or thiadiazolyl, each of which may be substituted with 1, 2, 3, 4 or 5 substituents (depending upon the number of available substitution positions) each independently selected from C₁-C₄ alkyl, O(C₁-C₄ alkyl), S(C₁-C₄ alkyl), halo, hydroxy, pyridyl, thiophenyl and phenyl optionally substituted with 1, 2, 3, 4 or 5 substituents each independently selected from halo, C₁-C₄ alkyl, or O(C₁-C₄ alkyl);

and Ar₂ is phenyl, pyridyl, pyrimidyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, thiophenyl, furanyl, imidazolyl or triazolyl each of which may be substituted with 1, 2, 3, 4 or 5 substituents (depending upon the number of available substitution positions) each independently selected from C₁-C₄ alkyl, O(C₁-C₄ alkyl) and halo; wherein each above-
5 mentioned C₁-C₄ alkyl group is optionally substituted with one or more halo atoms.

For the compounds of formula (IB) above, it is preferred that Ar₁ is a phenyl ring or a 5- or 6-membered heteroaryl ring substituted with 1, 2, 3, 4 or 5 substituents, more preferably with 1 or 2 substituents.

For the compounds of formula (IB) above, when Ar₁ is a substituted phenyl ring or
10 a substituted 5- or 6-membered heteroaryl ring, it is preferred that not more than one of those substituents is a pyridyl, thiophenyl or optionally substituted phenyl group.

Preferred compounds of formula (IB) above are those wherein Ar₁ includes a substituent attached at the 2-position. That is, the substituent is attached to the atom adjacent to that which forms the point of attachment of Ar₁ to the methylene group
15 connecting Ar₁ to the rest of the molecule. For example, when Ar₁ is phenyl, it is preferably ortho-substituted.

Further preferred compounds of formula (IB) above are those wherein R_x is H; R_y is H or C₁-C₄ alkyl; each R_z is independently H or C₁-C₄ alkyl; X represents O; Y represents OH or OR; R is C₁-C₄ alkyl; and Ar₁ and Ar₂ are each independently
20 selected from the group consisting of phenyl, and substituted phenyl; and pharmaceutically acceptable salts thereof. In this further preferred embodiment, the group Ar₁ may be substituted or unsubstituted phenyl. For example, Ar₁ may be unsubstituted phenyl or, preferably phenyl substituted with 1, 2, 3, 4 or 5 substituents, preferably with 1 or 2, for example 1, substituent. When disubstituted, the substituted phenyl group is
25 preferably substituted at the 2- and 5- positions. When monosubstituted, the substituted phenyl group is preferably substituted in the 2- position. Suitable substituents include C₁-C₄ alkyl, O(C₁-C₄ alkyl), S(C₁-C₄ alkyl), halo, and phenyl, optionally substituted with, for example, halo, C₁-C₄ alkyl, or O(C₁-C₄ alkyl). In this further preferred embodiment, the group Ar₂ may be substituted or unsubstituted phenyl. For example, Ar₂

may be phenyl substituted with 1, 2, 3, 4 or 5 substituents, preferably with 1 substituent. Suitable substituents include C₁-C₄ alkyl, O(C₁-C₄ alkyl), and especially, halo.

“C₁-C₄ alkyl” as used in respect of compounds of formula (IB) includes straight and branched chain alkyl groups of 1, 2, 3 or 4 carbon atoms, and may be unsubstituted or substituted. C₁-C₂ alkyl groups are preferred. Suitable substituents include halo, especially Cl and/or F. Thus the term “C₁-C₄ alkyl” includes haloalkyl. A particularly preferred substituted C₁-C₄ alkyl group is trifluoromethyl. Similar terms defining different numbers of C atoms (e.g. “C₁-C₃ alkyl”) take an analogous meaning. When R_y is C₁-C₄ alkyl it is preferably unsubstituted. When R_z is C₁-C₄ alkyl it is preferably unsubstituted. When R is C₁-C₄ alkyl it is preferably unsubstituted.

“5-membered heteroaryl ring” as used in respect of compounds of formula (IB) means a 5-membered aromatic ring including at least one heteroatom independently selected from N, O and S. Preferably there are not more than three heteroatoms in total in the ring. More preferably there are not more than two heteroatoms in total in the ring. More preferably there is not more than one heteroatom in total in the ring. The term includes, for example, the groups thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, thiophenyl, furanyl, pyrrolyl, imidazolyl, triazolyl, oxadiazolyl and thiadiazolyl.

“6-membered heteroaryl ring” as used in respect of compounds of formula (IB) means a 6-membered aromatic ring including at least one heteroatom independently selected from N, O and S. Preferably there are not more than three heteroatoms in total in the ring. More preferably there are not more than two heteroatoms in total in the ring. More preferably there is not more than one heteroatom in total in the ring. The term includes, for example, the groups pyridyl, pyrimidyl, pyrazinyl, pyridazinyl and triazinyl.

“Halo” as used in respect of compounds of formula (IB) includes F, Cl, Br and I, and is preferably F or Cl.

“Pyridyl” as used in respect of compounds of formula (IB) includes 2-pyridyl, 3-pyridyl and 4-pyridyl.

“Pyrimidyl” as used in respect of compounds of formula (IB) includes 2-pyrimidyl, 4-pyrimidyl and 5-pyrimidyl.

“Pyridazinyl” as used in respect of compounds of formula (IB) includes 3-pyridazinyl and 4-pyridazinyl.

“Pyrazinyl” as used in respect of compounds of formula (IB) includes 2-pyrazinyl and 3-pyrazinyl.

5 “Triazinyl” as used in respect of compounds of formula (IB) includes 2-(1,3,5-triazinyl), 3-, 5- and 6-(1,2,4-triazinyl) and 4- and 5-(1,2,3-triazinyl).

“Thiazolyl” as used in respect of compounds of formula (IB) includes 2-thiazolyl, 4-thiazolyl and 5-thiazolyl.

10 “Isothiazolyl” as used in respect of compounds of formula (IB) includes 3-isothiazolyl, 4-isothiazolyl, and 5-isothiazolyl.

“Oxazolyl” as used in respect of compounds of formula (IB) includes 2-oxazolyl, 4-oxazolyl and 5-oxazolyl.

“Isoxazolyl” as used in respect of compounds of formula (IB) includes 3-isoxazolyl, 4-isoxazolyl, and 5-isoxazolyl.

15 “Thiophenyl” as used in respect of compounds of formula (IB) includes 2-thiophenyl and 3-thiophenyl.

“Furanyl” as used in respect of compounds of formula (IB) includes 2-furanyl and 3-furanyl.

20 “Pyrrolyl” as used in respect of compounds of formula (IB) includes 2-pyrrolyl and 3-pyrrolyl.

“Imidazolyl” as used in respect of compounds of formula (IB) includes 2-imidazolyl and 4-imidazolyl.

“Triazolyl” as used in respect of compounds of formula (IB) includes 1-triazolyl, 4-triazolyl and 5-triazolyl.

25 “Oxadiazolyl” as used in respect of compounds of formula (IB) includes 4- and 5-(1,2,3-oxadiazolyl), 3- and 5-(1,2,4-oxadiazolyl), 3-(1,2,5-oxadiazolyl), 2-(1,3,4-oxadiazolyl).

30 “Thiadiazolyl” as used in respect of compounds of formula (IB) includes 4- and 5-(1,2,3-thiadiazolyl), 3- and 5-(1,2,4-thiadiazolyl), 3-(1,2,5-thiadiazolyl), 2-(1,3,4-thiadiazolyl).


For the compounds of formula (IB) above, Ry is preferably H or Me. More preferably Ry is H.

For the compounds of formula (IB) above, each Rz is preferably H or Me with 0, 1, 2 or 3 of Rz being Me. More preferably only 1 Rz is Me. Most preferably all Rz are H.


5 For the compounds of formula (IB) above, Y is preferably OH or OMe. More preferably, Y is OH.

For the compounds of formula (IB) above, it is preferred that Ry and all Rz are H and Y is OH.

10 For the compounds of formula (IB) above, the preferred stereochemistry is shown below:



A preferred group of compounds of formula (IB) is represented by the formula (IIB)



15


(IIB)

wherein R₁ and R₂ are each independently selected from H, C₁-C₄ alkyl, O(C₁-C₄ alkyl), S(C₁-C₄ alkyl), halo and phenyl; and R₃ is selected from H, C₁-C₄ alkyl and halo; and pharmaceutically acceptable salts thereof.

For the compounds of formula (IB) or (IIB) above, R_1 is preferably C_1 - C_3 alkyl (especially trifluoromethyl), $O(C_1$ - C_3 alkyl) (especially methoxy or trifluoromethoxy), F or phenyl (Ph). R_2 is preferably H. R_2 is also preferably F. R_3 is preferably H.

Especially preferred compounds of formula (IB) are 1-morpholin-2-yl-1-phenyl-2-(2-trifluoromethoxy-phenyl)-ethanol and 2-(5-fluoro-2-methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol. For both of these compounds the (S,R) stereoisomer is preferred. For both of these compounds the preferred salt form is the hydrochloride salt.

6. A compound of formula (IC)



(IC)

wherein: A is S or O; R is H; Ar is a phenyl group optionally substituted with 1, 2, 3, 4 or 5 substituents each independently selected from C_1 - C_4 alkyl, $O(C_1$ - C_4 alkyl), $S(C_1$ - C_4 alkyl), halo, hydroxy, $CO_2(C_1$ - C_4 alkyl), pyridyl, thiophenyl and phenyl optionally substituted with 1, 2, 3, 4 or 5 substituents each independently selected from halo, C_1 - C_4 alkyl, or $O(C_1$ - C_4 alkyl); X is a phenyl group optionally substituted with 1, 2, 3, 4 or 5 substituents each independently selected from halo, C_1 - C_4 alkyl, or $O(C_1$ - C_4 alkyl); a C_1 - C_4 alkyl group; a C_3 - C_6 cycloalkyl group or a $CH_2(C_3$ - C_6 cycloalkyl) group; R' is H or C_1 - C_4 alkyl; each R¹ is independently H or C_1 - C_4 alkyl; wherein each above-mentioned C_1 - C_4 alkyl group is optionally substituted with one or more halo atoms; or a pharmaceutically acceptable salt thereof; with the proviso that, when A is O, X is a C_1 - C_4 alkyl group, a C_3 - C_6 cycloalkyl group or a $CH_2(C_3$ - C_6 cycloalkyl) group.


For the compounds of formula (IC) above, it is preferred that A is S.

For the compounds of formula (IC) above, it is preferred that Ar is phenyl substituted with 1, 2, 3, 4 or 5 substituents, more preferably with 1 or 2 substituents.

When Ar is a substituted phenyl, it is preferred that not more than one of those substituents is a pyridyl, thiophenyl or optionally substituted phenyl group.

Preferred compounds of formula (IC) above are those wherein Ar is ortho-substituted.

5 Further preferred compounds of formula (IC) above are those of formula (ICa)



(ICa)

wherein: R is H; Ar is a phenyl group; X is a phenyl group; R' is H or C₁-C₄ alkyl; each R¹ is independently H or C₁-C₄ alkyl; and pharmaceutically acceptable salts thereof. For these further preferred compounds, the group Ar may be substituted or unsubstituted phenyl. For example, Ar may be unsubstituted phenyl or, preferably phenyl substituted with 1, 2, 3, 4 or 5 substituents, preferably with 1 or 2, for example 1, substituent. When disubstituted, the substituted phenyl group is preferably substituted at the 2- and 5-positions. When monosubstituted, the substituted phenyl group is preferably substituted in the 2- position. Suitable substituents include C₁-C₄ alkyl, O(C₁-C₄ alkyl), S(C₁-C₄ alkyl), halo, and phenyl optionally substituted with, for example, halo, C₁-C₄ alkyl, or O(C₁-C₄ alkyl). For these further preferred compounds, the group X may be substituted or unsubstituted phenyl. For example, X may be phenyl substituted with 1, 2, 3, 4 or 5 substituents, preferably with 1 substituent. Suitable substituents include C₁-C₄ alkyl, O(C₁-C₄ alkyl), and halo.

“C₁-C₄ alkyl” as used in respect of compounds of formula (IC) includes straight and branched chain alkyl groups of 1, 2, 3 or 4 carbon atoms, and may be unsubstituted or substituted. C₁-C₂ alkyl groups are preferred. Suitable substituents include halo. Thus the term “C₁-C₄ alkyl” includes haloalkyl. Similar terms defining different numbers of C

atoms (e.g. "C₁-C₃ alkyl") take an analogous meaning. When R' is C₁-C₄ alkyl it is preferably unsubstituted. When R¹ is C₁-C₄ alkyl it is preferably unsubstituted.

"C₃-C₆ cycloalkyl" as used in respect of compounds of formula (IC) includes cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

5 "Halo" as used in respect of compounds of formula (IC) includes F, Cl, Br and I, and is preferably F or Cl.

"Pyridyl" as used in respect of compounds of formula (IC) includes 2-pyridyl, 3-pyridyl and 4-pyridyl.


10 "Thiophenyl" as used in respect of compounds of formula (IC) includes 2-thiophenyl and 3-thiophenyl.

For the compounds of formula (IC) above, R' is preferably H or Me. More preferably R' is H.

For the compounds of formula (IC) above, each R¹ is preferably H or Me with 0, 1, 2 or 3 of R¹ being Me. More preferably only 1 R¹ is Me. Most preferably all R¹ are H.

15 For the compounds of formula (IC) above, it is preferred that R' and all R¹ are H. A particularly preferred substituted C₁-C₄ alkyl group for the group Ar is trifluoromethyl.

A preferred group of compounds of formula (IC) is represented by the formula (IIC);




(IIC)

wherein R₂ and R₃ are each independently selected from H, C₁-C₄ alkyl, O(C₁-C₄ alkyl), S(C₁-C₄ alkyl), halo and phenyl; and R₄ is selected from H and C₁-C₄ alkyl; and

pharmaceutically acceptable salts thereof. R_2 is preferably C_1 - C_3 alkyl (especially trifluoromethyl), $O(C_1$ - C_3 alkyl) (especially methoxy or trifluoromethoxy), F or Ph. R_3 is preferably H. R_3 is also preferably F. R_4 is preferably H.


7. A compound of formula (ID)



(ID)

wherein -X- is $-C(R^4R^5)-$, -O- or -S-; n is 2 or 3; R^1 is H or C_1 - C_4 alkyl; R^3 is H, halo, C_1 - C_4 alkyl, $O(C_1$ - C_4 alkyl), nitrile, phenyl or substituted phenyl; R^4 and R^5 are each independently selected from H or C_1 - C_4 alkyl; Ar- is selected from the group consisting

of



in which R^{2a} is H, halo, methyl or ethyl; R^{2b} is H, halo or methyl; R^{2c} is H, halo, methyl, trifluoromethyl, nitrile, or methoxy; R^{2d} is H, halo, methyl or ethyl; R^{2e} is H, halo, methyl, trifluoromethyl, nitrile, or methoxy; R^{2f} is H, or fluoro; -Y- is -O-, -S- or $-N(R^6)-$; and R^6 is H or methyl and pharmaceutically acceptable salts thereof.

The term " C_1 - C_4 alkyl" as used in respect of compounds of formula (ID)


includes straight and branched chain alkyl groups of 1, 2, 3 or 4 carbon atoms. Thus the term " C_1 - C_4 alkyl" includes methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl. C_1 - C_2 alkyl groups are preferred. A particularly preferred C_1 - C_4 alkyl group is methyl or ethyl.

The term "halo" as used in respect of compounds of formula (ID) includes F, Cl, Br and I, and is preferably F or Cl.

The term "substituted phenyl" as used in respect of compounds of formula (ID) means phenyl substituted with 1, 2, 3, 4 or 5 substituents, preferably with 1 or 2, for example 1, substituent. Suitable substituents include C₁-C₄ alkyl, O(C₁-C₄ alkyl), S(C₁-C₄ alkyl), halo, and phenyl optionally substituted with, for example, C₁-C₄ alkyl, O(C₁-C₄ alkyl), S(C₁-C₄ alkyl), or halo.

The terms "O(C₁-C₄ alkyl)" or "S(C₁-C₄ alkyl)" as used in respect of compounds of formula (ID) mean a C₁-C₄ alkyl group as defined above linked to the point of substitution via an oxygen or a sulphur atom. An O(C₁-C₄ alkyl) or S(C₁-C₄ alkyl) group includes for example methoxy, ethoxy, thiomethyl or thioethyl.

Preferred compounds of formula (ID) are represented by the formula (IDa)



wherein -X-, n, R¹, R³ and Ar have the values as defined for formula (ID) above.


Compounds of formula (ID) or (IDa) wherein -X- is -C(R⁴R⁵)- are preferred. Even more preferred are compounds of formula (ID) or (IDa) wherein -X- is -C(R⁴R⁵)- and R⁴ and R⁵ are both H or R⁴ and R⁵ are both the same C₁-C₄ alkyl.

Compounds of formula (ID) or (IDa) wherein Ar is (i) are also preferred. Preferably Ar is (i) and R^{2c} is H. Even more preferred are compounds of formula (ID) or (IDa) wherein Ar is (i), R^{2c} is H, and (a) R^{2a} is H or methyl, R^{2b} is H and R^{2f} is H or (b) R^{2a} is H, R^{2b} is halo, preferably fluoro or chloro and R^{2f} is H or fluoro.

Another group of preferred compounds of formula (ID) or (IDa) are compounds wherein Ar is (ii) and -Y- is -S-. More preferably Ar is 2-thiophenyl or 3-thiophenyl.

A further preferred group of compounds of formula (ID) is represented by the formula (IID)

37




(IID)

wherein n is 2 or 3; R^1 is H or C_1 - C_4 alkyl; R^3 is H, halo, phenyl or substituted phenyl; R^{2a} is H, halo, methyl or ethyl; R^{2b} is H, halo or methyl; and pharmaceutically acceptable salts thereof.


- 5 Preferred compounds of formulae (ID), (IDa) and (IID) are those wherein n is 3, or wherein R^1 is H, methyl, ethyl or n -propyl, or wherein R^3 is H or halo.

8. A compound of formula (IE)



(IE)

- 10 wherein R^1 is C_1 - C_6 alkyl (optionally substituted with 1, 2 or 3 halo substituents and/or with 1 substituent selected from -S-(C_1 - C_3 alkyl), -O-(C_1 - C_3 alkyl) (optionally substituted with 1, 2 or 3 F atoms), -O-(C_3 - C_6 cycloalkyl), -SO₂-(C_1 - C_3 alkyl), -CN, -COO-(C_1 - C_2 alkyl) and -OH); C_2 - C_6 alkenyl; -(CH₂)_q-Ar₂; or a group of formula (i) or (ii)



15

R^2 , R^3 and R^4 are each independently selected from hydrogen or C_1 - C_2 alkyl; R^5 , R^6 , R^7 and R^8 are at each occurrence independently selected from hydrogen or C_1 - C_2 alkyl; -X- is a bond, -CH₂-, -CH=CH-, -O-, -S-, or -SO₂-; -Y- is a bond, -CH₂- or -O-; -Z is

hydrogen, -OH or -O-(C₁-C₃ alkyl); p is 0, 1 or 2; q is 0, 1 or 2; r is 0 or 1; s is 0, 1, 2 or 3; t is 0, 1, 2 or 3; Ar₁ is phenyl, pyridyl, thiazolyl, benzothiophenyl or naphthyl; wherein said phenyl, pyridyl or thiazolyl group may be substituted with 1, 2 or 3 substituents each independently selected from halo, cyano, C₁-C₄ alkyl (optionally substituted with 1, 2 or 3 F atoms), -O-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms) and -S-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms) and/or with 1 substituent selected from pyridyl, pyrazole, phenyl (optionally substituted with 1, 2 or 3 halo substituents) and phenoxy (optionally substituted with 1, 2 or 3 halo substituents); and wherein said benzothiophenyl or naphthyl group may be optionally substituted with 1, 2 or 3 substituents each independently selected from halo, cyano, C₁-C₄ alkyl (optionally substituted with 1, 2 or 3 F atoms), -O-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms), and -S-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms); Ar₂ is naphthyl, pyridyl, thiazolyl, furyl, thiophenyl, benzothiophenyl, or phenyl, wherein said naphthyl, pyridyl, thiazolyl, furyl, thiophenyl, benzothiophenyl, or phenyl may be substituted with 1, 2 or 3 substituents each independently selected from halo, C₁-C₄ alkyl (optionally substituted with 1, 2 or 3 F atoms) and -O-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms); and pharmaceutically acceptable salts thereof; provided that (a) the cyclic portion of the group of formula (i) must contain at least three carbon atoms and not more than seven ring atoms; (b) when -X- is -CH=CH-, then the cyclic portion of the group of formula (i) must contain at least five carbon atoms; and (c) when -Z is -OH or -O-(C₁-C₃ alkyl), then -X- is -CH₂-; (d) when -Y- is -O- then p cannot be 0; and (e) the compound 3-[(phenylmethyl)-(3S)-3-pyrrolidinylamino]-propanenitrile is excluded.

With respect to formula (IE) the term "C₁-C₆ alkyl" means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 6 carbon atoms.

With respect to formula (IE) the term "C₂-C₆ alkenyl" means a monovalent unsubstituted unsaturated straight-chain or branched-chain hydrocarbon radical having from 2 to 6 carbon atoms and containing at least one carbon-carbon double bond.

With respect to formula (IE) the term "C₃-C₆ cycloalkyl" means a monovalent unsubstituted saturated cyclic hydrocarbon radical having from 3 to 6 carbon atoms.

With respect to formula (IE) the term "C₁-C₆ alkylene" means a divalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 6 carbon atoms.

With respect to formula (IE) the term "halo" or "halogen" means F, Cl, Br or I.

5 With respect to formula (IE) the term "C₁-C₄ difluoroalkyl" means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 4 carbon atoms wherein two hydrogen atoms are substituted with two fluoro atoms. Preferably the two fluoro atoms are attached to the same carbon atom.

10 With respect to formula (IE) the term "C₁-C₄ trifluoroalkyl" means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 4 carbon atoms wherein three hydrogen atoms are substituted with three fluoro atoms. Preferably the three fluoro atoms are attached to the same carbon atom.

With respect to formula (IE) the term "phenoxy" means a monovalent unsubstituted phenyl radical linked to the point of substitution by an O atom.

15 With respect to formula (IE) the term "pyridyl" includes 2-pyridyl, 3-pyridyl and 4-pyridyl.

With respect to formula (IE) the term "furyl" includes 2-furyl and 3-furyl. 2-furyl is preferred.

20 With respect to formula (IE) the term "thiophenyl" includes 2-thiophenyl and 3-thiophenyl.

With respect to formula (IE) the term "thiazolyl" includes 2-thiazolyl, 4-thiazolyl and 5-thiazolyl.

With respect to formula (IE) the term "pyrazole" includes 1-pyrazole, 3-pyrazole and 4-pyrazole. 1-pyrazole is preferred.

25 With respect to formula (IE) the term "benzothiophenyl" includes 2-benzo[b]thiophenyl, 3-benzo[b]thiophenyl, 4-benzo[b]thiophenyl, 5-benzo[b]thiophenyl, 6-benzo[b]thiophenyl and 7-benzo[b]thiophenyl.


With respect to formula (IE) the term "naphthyl" includes 1-naphthyl, and 2-naphthyl. 1-naphthyl is preferred.

With respect to formula (IE), similar terms specifying different numbers of C atoms take an analogous meaning. For example the terms "C₁-C₄ alkyl" and "C₁-C₃ alkyl" mean a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 4 and 1 to 3 carbon atoms respectively. The term
 5 "C₁-C₄ alkyl" includes methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and tert-butyl. The term "C₁-C₃ alkyl" includes methyl, ethyl, n-propyl and iso-propyl.

With respect to formula (IE) it will be appreciated that when s is 2 or 3, then each R⁵ and/or each R⁶ can be different. In the same way when t is 2 or 3, then each R⁷ and/or each R⁸ can be different.

10 Preferred compounds of formula (IE) are those wherein R¹ is C₁-C₆ alkyl, C₂-C₆ alkenyl, -(CH₂)_m-CF₃, -(CH₂)_n-S-(C₁-C₃ alkyl), -CH₂-COO-(C₁-C₂ alkyl), -(C₁-C₅ alkylene)-O-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-O-(C₃-C₆ cycloalkyl), -(C₁-C₅ alkylene)-SO₂-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-OCF₃, -(C₁-C₆ alkylene)-OH, -(C₁-C₅ alkylene)-CN, -(CH₂)_q-Ar₂ or a group of formula (ia), (ib) or (ii)

15



R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, -X-, -Y-, p, q, r and s have the values defined above; m is 1, 2 or 3; n is 1, 2 or 3; t is 2, 3 or 4; -Ar₁ is phenyl, pyridyl, thiazolyl or naphthyl; wherein
 20 said phenyl, pyridyl or thiazolyl group may be substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl, cyano, C₁-C₄ alkyl, -O-(C₁-C₄ alkyl), -O-(C₁-C₄ difluoroalkyl), -O-(C₁-C₄ trifluoroalkyl), -S-(C₁-C₄ alkyl), -S-(C₁-C₂ trifluoroalkyl) and/or with 1 substituent selected from pyridyl, pyrazole, phenyl (optionally substituted with 1, 2 or 3 halo substituents) and phenoxy (optionally
 25 substituted with 1, 2 or 3 halo substituents); and wherein said naphthyl group may be optionally substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl, cyano, C₁-C₄ alkyl, -O-(C₁-C₄ alkyl), -O-(C₁-C₄ difluoroalkyl), -O-(C₁-C₄ trifluoroalkyl), -S-(C₁-C₄ alkyl), -S-(C₁-C₂ trifluoroalkyl); Ar₂ is naphthyl, pyridyl,

thiazolyl, furyl, thiophenyl, benzothiophenyl, or phenyl, wherein said naphthyl, pyridyl, thiazolyl, furyl, thiophenyl, benzothiophenyl, or phenyl may be substituted with 1, 2 or 3 substituents each independently selected from halo, C₁-C₄ alkyl, trifluoromethyl and -O-(C₁-C₄ alkyl); and pharmaceutically acceptable salts thereof.

5 Preferred compounds of formula (IE) are those wherein R² is hydrogen. In another preferred embodiment R³ and R⁴ are hydrogen. More preferably R², R³ and R⁴ are hydrogen.

Preferred compounds of formula (IE) are those wherein each R⁵ and R⁶ is hydrogen. In another preferred embodiment each R⁷ and R⁸ is hydrogen. More preferably
10 R⁵, R⁶, R⁷ and R⁸ are hydrogen.

Preferred compounds of formula (IE) are those wherein R¹ is C₁-C₆ alkyl. More preferably R¹ is n-propyl, 1-methylethyl, 2-methylpropyl, 3,3-dimethylpropyl.

Preferred compounds of formula (IE) are those wherein R¹ is -(C₄-C₅ alkylene)-OH. More preferably R¹ is 2,2-dimethyl-2-hydroxyethyl or 3,3-dimethyl-3-
15 hydroxypropyl.

Preferred compounds of formula (IE) are those wherein R¹ is a group of formula (i) and each R⁵ and R⁶ is hydrogen. More preferably each R⁵, R⁶, R⁷ and R⁸ is hydrogen.

Preferred compounds of formula (IE) are those wherein R¹ is a group of formula (ii) and each R⁵ and R⁶ is hydrogen. More preferably each R⁵, R⁶, R⁷ and R⁸ is hydrogen.

20 Preferred compounds of formula (IE) are those wherein R¹ is a group of formula (i), r is 0, s is 2, t is 2, -Z is hydrogen and -X- is -O-, -S- or -SO₂-. More preferably R¹ is a group of formula (i), r is 0, s is 2, t is 1 or 2, -Z is hydrogen and -X- is -O-.

Preferred compounds of formula (IE) are those wherein R¹ is a group of formula (i), r is 0, s is 1, 2 or 3, t is 1, -Z is hydrogen and -X- is -CH₂-.

25 Preferred compounds of formula (IE) are those wherein R¹ is a group of formula (i), r is 1, s is 0, 1, 2 or 3, t is 1, -Z is hydrogen and -X- is -CH₂-.

Preferred compounds of formula (IE) are those wherein R¹ is a group of the formula (ia). More preferably R¹ is a group of the formula (ia) and each R⁵, R⁶, R⁷ and R⁸ is hydrogen.

Preferred compounds of formula (IE) are those wherein R^1 is a group of the formula (ib). More preferably R^1 is a group of the formula (ib), r is 1, t is 3, and each R^7 and R^8 is hydrogen.

Preferred compounds of formula (IE) are those wherein R^1 is $-(CH_2)_m-CF_3$. More preferably R^1 is $-(CH_2)_m-CF_3$ and m is 1, 2, or 3.

Preferred compounds of formula (IE) are those wherein R^1 is $-(CH_2)_n-S-(C_1-C_3 \text{ alkyl})$. More preferably R^1 is $-(CH_2)_3-S-CH_3$.

Preferred compounds of formula (IE) are those wherein R^1 is $-CH_2-COO-(C_1-C_2 \text{ alkyl})$. More preferably R^1 is $-CH_2-COOCH_3$.

Preferred compounds of formula (IE) are those wherein R^1 is $-(C_1-C_5 \text{ alkylene})-O-(C_1-C_3 \text{ alkyl})$. More preferably R^1 is $-(C_3-C_4 \text{ alkylene})-OCH_3$.

Preferred compounds of formula (IE) are those wherein R^1 is $-(C_1-C_5 \text{ alkylene})-(C_3-C_6 \text{ cycloalkyl})$. More preferably R^1 is $-CH_2-CH_2-O\text{-cyclobutyl}$.

Preferred compounds of formula (IE) are those wherein R^1 is $-(C_1-C_5 \text{ alkylene})-SO_2-(C_1-C_3 \text{ alkyl})$.

Preferred compounds of formula (IE) are those wherein R^1 is $-(C_1-C_5 \text{ alkylene})-OCF_3$. More preferably R^1 is $-CH_2-CH_2-OCF_3$.

Preferred compounds of formula (IE) are those wherein R^1 is $-(C_1-C_5 \text{ alkylene})-CN$. More preferably R^1 is $-(C_2-C_4 \text{ alkylene})-CN$. Most preferably $-CH_2-CH_2-CN$ or $-CH_2-C(CH_3)_2-CN$.


Preferred compounds of formula (IE) are those wherein R^1 is $-(CH_2)_q-Ar_2$, and q is 1. More preferably R^1 is $-(CH_2)_q-Ar_2$, q is 1 and $-Ar_2$ is pyridyl, phenyl or phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl or C_1-C_4 alkyl.

Preferred compounds of formula (IE) are those wherein $-Ar_1$ is phenyl; phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl and C_1-C_4 alkyl and/or with 1 substituent selected from phenyl, phenyl substituted with 1, 2 or 3 halo substituents, pyridyl, pyrazole, phenoxy and phenoxy substituted with 1, 2 or 3 halo substituents; pyridyl; or pyridyl substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl and C_1-C_4 alkyl and/or with 1 substituent selected from phenyl and phenyl substituted with 1, 2 or 3 halo



substituents. More preferably $-Ar_1$ is phenyl or phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl and C_1 - C_4 alkyl and/or with 1 substituent selected from phenyl, phenyl substituted with 1, 2 or 3 halo substituents, pyridyl, pyrazole, phenoxy and phenoxy substituted with 1, 2 or 3 halo substituents. Most preferably $-Ar_1$ is phenyl substituted with 1 or 2 substituents each independently selected from halo, trifluoromethyl and C_1 - C_4 alkyl and/or with 1 substituent selected from phenyl, phenyl substituted with 1, 2 or 3 halo substituents, pyridyl, pyrazole, phenoxy and phenoxy substituted with 1, 2 or 3 halo substituents. Suitable $-Ar_1$ groups include, for example, 2-methylthiophenyl, 2-methylphenyl, 2-fluorophenyl, 2-chlorophenyl, 2-isopropoxyphenyl, 2-trifluoromethylphenyl, 2-difluoromethoxyphenyl, 2-methoxyphenyl, 2-ethoxyphenyl, 2-(1,1'-biphenyl), 2-phenoxyphenyl, 2-benzylphenyl, 3-trifluoromethoxyphenyl, 3-chlorophenyl, 3-trifluoromethylphenyl, 3-methylphenyl, 3-trifluoromethoxyphenyl, 3-methoxyphenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 4-fluorophenyl, 3,5-dichlorophenyl, 3,5-dimethylphenyl, 3-trifluoromethyl-5-fluorophenyl, 3,5-difluorophenyl, 2,3-dichlorophenyl, 2,3-dimethylphenyl, 2-chloro-3-trifluoromethylphenyl, 2-chloro-3-methylphenyl, 2-methyl-3-chlorophenyl, 2,4-dichlorophenyl, 2,4-dimethyl, 2,4-difluorophenyl, 2-chloro-4-fluorophenyl, 2-trifluoromethyl-4-fluorophenyl, 2-fluoro-4-trifluoromethylphenyl, 2-methyl-4-chlorophenyl, 2-methoxy-4-fluorophenyl, 2-trifluoromethyl-5-fluorophenyl, 2,5-dimethylphenyl, 4-fluoro-[1,1'-biphenyl]-2-yl, 2-chloro-5-fluorophenyl, 2-(trifluoromethyl)-6-fluorophenyl, 2-chloro-6-fluorophenyl, 3,4-dichlorophenyl, and 3-chloro-4-fluorophenyl. In general when $-Ar_1$ is phenyl substituted with pyridyl, 3-pyridyl is preferred.

Preferred compounds of formula (IE) are those wherein $-Ar_1$ is pyridyl or pyridyl substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl and C_1 - C_4 alkyl and/or with 1 substituent selected from phenyl and phenyl substituted with 1, 2 or 3 halo substituents. More preferably $-Ar_1$ is pyridyl substituted with 1 or 2 substituents each independently selected from halo, trifluoromethyl and C_1 - C_4 alkyl and/or with 1 substituent selected from phenyl and phenyl substituted with 1, 2 or 3 halo substituents. Suitable $-Ar_1$ groups include, for example, 3-phenyl-2-pyridyl. In general when $-Ar_1$ is a substituted pyridyl, substituted 2-pyridyl is preferred.


9. A compound of formula (IF)





5 wherein



or



R^1 is C_1 - C_6 alkyl (optionally substituted with 1, 2 or 3 halo substituents and/or with 1 substituent selected from -S-(C_1 - C_3 alkyl), -O-(C_1 - C_3 alkyl) (optionally substituted with 1, 2 or 3 F atoms), -O-(C_3 - C_6 cycloalkyl), -SO₂-(C_1 - C_3 alkyl), -CN, -COO-(C_1 - C_2 alkyl) and -OH); C_2 - C_6 alkenyl; -(CH₂)_q-Ar₂; or a group of formula (i) or (ii)



R^2 , R^3 and R^4 are each independently selected from hydrogen or C_1 - C_2 alkyl; R^5 , R^6 , R^7 and R^8 are at each occurrence independently selected from hydrogen or C_1 - C_2 alkyl; -X- is a bond, -CH₂-, -CH=CH-, -O-, -S-, or -SO₂-; -Y- is a bond, -CH₂- or -O-; -Z is hydrogen, -OH or -O-(C_1 - C_3 alkyl); p is 0, 1 or 2; q is 0, 1 or 2; r is 0 or 1; s is 0, 1, 2 or

3; t is 0, 1, 2 or 3; Ar₁ is phenyl, pyridyl, thiazolyl, benzothiophenyl or naphthyl; wherein said phenyl, pyridyl or thiazolyl group may be substituted with 1, 2 or 3 substituents each independently selected from halo, cyano, C₁-C₄ alkyl (optionally substituted with 1, 2 or 3 F atoms), -O-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms) and -S-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms) and/or with 1 substituent selected from pyridyl, pyrazole, phenyl (optionally substituted with 1, 2 or 3 halo substituents), benzyl and phenoxy (optionally substituted with 1, 2 or 3 halo substituents); and wherein said benzothiophenyl or naphthyl group may be optionally substituted with 1, 2 or 3 substituents each independently selected from halo, cyano, C₁-C₄ alkyl (optionally substituted with 1, 2 or 3 F atoms), -O-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms), and -S-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms); Ar₂ is naphthyl, pyridyl, thiazolyl, furyl, thiophenyl, benzothiophenyl, or phenyl, wherein said naphthyl, pyridyl, thiazolyl, furyl, thiophenyl, benzothiophenyl, or phenyl may be substituted with 1, 2 or 3 substituents each independently selected from halo, C₁-C₄ alkyl (optionally substituted with 1, 2 or 3 F atoms) and -O-(C₁-C₄ alkyl) (optionally substituted with 1, 2 or 3 F atoms); or a pharmaceutically acceptable salt thereof; provided that (a) the cyclic portion of the group of formula (i) must contain at least three carbon atoms and not more than seven ring atoms; (b) when -X- is -CH=CH-, then the cyclic portion of the group of formula (i) must contain at least five carbon atoms; and (c) when -Z is -OH or -O-(C₁-C₃ alkyl), then -X- is -CH₂-; and (d) when -Y- is -O- then p cannot be 0.

With respect to formula (IF) the term "C₁-C₆ alkyl" means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 6 carbon atoms.

With respect to formula (IF) the term "C₂-C₆ alkenyl" means a monovalent unsubstituted unsaturated straight-chain or branched-chain hydrocarbon radical having from 2 to 6 carbon atoms and containing at least one carbon-carbon double bond.

With respect to formula (IF) the term "C₃-C₆ cycloalkyl" means a monovalent unsubstituted saturated cyclic hydrocarbon radical having from 3 to 6 carbon atoms.

With respect to formula (IF) the term "C₁-C₆ alkylene" means a divalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 6 carbon atoms.

With respect to formula (IF) the term "halo" or "halogen" means F, Cl, Br or I.

5 With respect to formula (IF) the term "C₁-C₄ difluoroalkyl" means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 4 carbon atoms wherein two hydrogen atoms are substituted with two fluoro atoms. Preferably the two fluoro atoms are attached to the same carbon atom.

10 With respect to formula (IF) the term "C₁-C₄ trifluoroalkyl" means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 4 carbon atoms wherein three hydrogen atoms are substituted with three fluoro atoms. Preferably the three fluoro atoms are attached to the same carbon atom.

With respect to formula (IF) the term "phenoxy" means a monovalent unsubstituted phenyl radical linked to the point of substitution by an O atom.

15 With respect to formula (IF) the term "pyridyl" includes 2-pyridyl, 3-pyridyl and 4-pyridyl.

With respect to formula (IF) the term "furyl" includes 2-furyl and 3-furyl. 2-furyl is preferred.

20 With respect to formula (IF) the term "thiophenyl" includes 2-thiophenyl and 3-thiophenyl.

With respect to formula (IF) the term "thiazolyl" includes 2-thiazolyl, 4-thiazolyl and 5-thiazolyl.

With respect to formula (IF) the term "pyrazole" includes 1-pyrazole, 3-pyrazole and 4-pyrazole. 1-pyrazole is preferred.


25 With respect to formula (IF) the term "benzothiophenyl" includes 2-benzo[b]thiophenyl, 3-benzo[b]thiophenyl, 4-benzo[b]thiophenyl, 5-benzo[b]thiophenyl, 6-benzo[b]thiophenyl and 7-benzo[b]thiophenyl.

With respect to formula (IF) the term "naphthyl" includes 1-naphthyl, and 2-naphthyl. 1-naphthyl is preferred.

With respect to formula (IF), similar terms specifying different numbers of C atoms take an analogous meaning. For example the terms "C₁-C₄ alkyl" and "C₁-C₃ alkyl" mean a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 4 and 1 to 3 carbon atoms respectively. The term
 5 "C₁-C₄ alkyl" includes methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and tert-butyl. The term "C₁-C₃ alkyl" includes methyl, ethyl, n-propyl and iso-propyl.

With respect to formula (IF), it will be appreciated that when s is 2 or 3, then each R⁵ and/or each R⁶ can be different. In the same way when t is 2 or 3, then each R⁷ and/or each R⁸ can be different.


10 Preferred compounds of formula (IF) are those of formula (IF')



(IF')

wherein R¹, R², R³, R⁴ and Ar₁ have the values defined in formula (IF) above.

Preferred compounds of formula (IF) are those of formula (IF'')




(IF'')

15

wherein R¹, R², R³, R⁴ and Ar₁ have the values defined in formula (IF) above.

Preferred compounds of formula (IF) are those wherein R¹ is C₁-C₆ alkyl, C₂-C₆ alkenyl, -(CH₂)_m-CF₃, -(CH₂)_n-S-(C₁-C₃ alkyl), -CH₂-COO-(C₁-C₂ alkyl), -(C₁-C₅ alkylene)-O-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-O-(C₃-C₆ cycloalkyl), -(C₁-C₅ alkylene)-SO₂-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-OCF₃, -(C₁-C₆ alkylene)-OH, -(C₁-C₅ alkylene)-CN,
 20 -(CH₂)_q-Ar₂ or a group of formula (ia), (ib) or (ii)

48



R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , $-X$, $-Y$, p , q , r and s have the values defined above; m is 1, 2 or 3; n is 1, 2 or 3; t is 2, 3 or 4; $-Ar_1$ is phenyl, pyridyl, thiazolyl or naphthyl; wherein said phenyl, pyridyl or thiazolyl group may be substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl, cyano, C_1 - C_4 alkyl, $-O$ -(C_1 - C_4 alkyl), $-O$ -(C_1 - C_4 difluoroalkyl), $-O$ -(C_1 - C_4 trifluoroalkyl), $-S$ -(C_1 - C_4 alkyl), $-S$ -(C_1 - C_2 trifluoroalkyl) and/or with 1 substituent selected from pyridyl, pyrazole, phenyl (optionally substituted with 1, 2 or 3 halo substituents) and phenoxy (optionally substituted with 1, 2 or 3 halo substituents); and wherein said naphthyl group may be optionally substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl, cyano, C_1 - C_4 alkyl, $-O$ -(C_1 - C_4 alkyl), $-O$ -(C_1 - C_4 difluoroalkyl), $-O$ -(C_1 - C_4 trifluoroalkyl), $-S$ -(C_1 - C_4 alkyl), $-S$ -(C_1 - C_2 trifluoroalkyl); Ar_2 is naphthyl, pyridyl, thiazolyl, furyl, thiophenyl, benzothiophenyl, or phenyl, wherein said naphthyl, pyridyl, thiazolyl, furyl, thiophenyl, benzothiophenyl, or phenyl may be substituted with 1, 2 or 3 substituents each independently selected from halo, C_1 - C_4 alkyl, trifluoromethyl and $-O$ -(C_1 - C_4 alkyl).

Preferred compounds of formula (IF) are those wherein R^2 is hydrogen. In another preferred embodiment R^3 and R^4 are hydrogen. More preferably R^2 , R^3 and R^4 are hydrogen.

Preferred compounds of formula (IF) are those wherein each R^5 and R^6 is hydrogen. In another preferred embodiment each R^7 and R^8 is hydrogen. More preferably R^5 , R^6 , R^7 and R^8 are hydrogen.

Preferred compounds of formula (IF) are those wherein R^1 is C_1 - C_6 alkyl. More preferably R^1 is n-propyl, 1-methylethyl (i-propyl), 2-methylpropyl (i-butyl), 2-methylbutyl, 2,2-dimethylbutyl.

Preferred compounds of formula (IF) are those wherein R^1 is $-(C_4-C_5 \text{ alkylene})-$ OH. More preferably R^1 is 2,2-dimethyl-2-hydroxyethyl or 3,3-dimethyl-3-hydroxypropyl.

Preferred compounds of formula (IF) are those wherein R^1 is a group of formula
5 (i) and each R^5 and R^6 is hydrogen. More preferably each R^5 , R^6 , R^7 and R^8 is hydrogen.

Preferred compounds of formula (IF) are those wherein R^1 is a group of formula
(ii) and each R^5 and R^6 is hydrogen. More preferably each R^5 , R^6 , R^7 and R^8 is hydrogen.

Preferred compounds of formula (IF) are those wherein R^1 is a group of formula
(i), r is 0 or 1, s is 2, t is 1 or 2, $-Z$ is hydrogen and $-X-$ is $-O-$, $-S-$ or $-SO_2-$. More
10 preferably R^1 is a group of formula (i), r is 0 or 1, s is 2, t is 1 or 2, $-Z$ is hydrogen and $-X-$ is $-O-$, for example tetrahydro-2*H*-pyran-4-yl, tetrahydrofuran-3-yl or
(tetrahydrofuran-3-yl)methyl. Most preferably R^1 is a group of formula (i), r is 0, s is 2, t
is 1 or 2, $-Z$ is hydrogen and $-X-$ is $-O-$, for example tetrahydro-2*H*-pyran-4-yl or
tetrahydrofuran-3-yl.

15 Preferred compounds of formula (IF) are those wherein R^1 is a group of formula
(i), r is 0, s is 1, 2 or 3, t is 1, $-Z$ is hydrogen and $-X-$ is $-CH_2-$, for example cyclobutyl,
cyclopentyl or cyclohexyl.

Preferred compounds of formula (IF) are those wherein R^1 is a group of formula
(i), r is 1, s is 0, 1, 2 or 3, t is 1, $-Z$ is hydrogen and $-X-$ is $-CH_2-$.

20 Preferred compounds of formula (IF) are those wherein R^1 is a group of the
formula (ia). More preferably R^1 is a group of the formula (ia) and each R^5 , R^6 , R^7 and R^8
is hydrogen.

Preferred compounds of formula (IF) are those wherein R^1 is a group of the
formula (ib). More preferably R^1 is a group of the formula (ib), r is 1, t is 3, and each
25 R^7 and R^8 is hydrogen.

Preferred compounds of formula (IF) are those wherein R^1 is $-(CH_2)_m-CF_3$. More
preferably R^1 is $-(CH_2)_m-CF_3$ and m is 1, 2, or 3.

Preferred compounds of formula (IF) are those wherein R^1 is $-(CH_2)_n-S-(C_1-C_3$
alkyl). More preferably R^1 is $-(CH_2)_3-S-CH_3$.

30 Preferred compounds of formula (IF) are those wherein R^1 is $-CH_2-COO-(C_1-C_2$
alkyl). More preferably R^1 is $-CH_2-COOCH_3$.

Preferred compounds of formula (IF) are those wherein R¹ is -(C₁-C₅ alkylene)-O-(C₁-C₃ alkyl). More preferably R¹ is -(C₃-C₄ alkylene)-OCH₃.

Preferred compounds of formula (IF) are those wherein R¹ is -(C₁-C₅ alkylene)-O-(C₃-C₆ cycloalkyl). More preferably R¹ is -CH₂-CH₂-O-cyclobutyl.

5 Preferred compounds of formula (IF) are those wherein R¹ is -(C₁-C₅ alkylene)-SO₂-(C₁-C₃ alkyl).

Preferred compounds of formula (IF) are those wherein R¹ is -(C₁-C₅ alkylene)-OCF₃. More preferably R¹ is -CH₂-CH₂-OCF₃.

Preferred compounds of formula (IF) are those wherein R¹ is -(C₁-C₅ alkylene)-CN. More preferably R¹ is -(C₂-C₄ alkylene)-CN. Most preferably -CH₂-CH₂-CN or
10 -CH₂-C(CH₃)₂-CN.


Preferred compounds of formula (IF) are those wherein R¹ is -(CH₂)_q-Ar₂, and q is 1. More preferably R¹ is -(CH₂)_q-Ar₂, q is 1 and -Ar₂ is pyridyl, phenyl or phenyl substituted with 1, 2 or 3 substituents each independently selected from halo,
15 trifluoromethyl, C₁-C₄ alkyl or O-(C₁-C₄ alkyl).

Preferred compounds of formula (IF) are those wherein -Ar₁ is phenyl; phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl and C₁-C₄ alkyl and/or with 1 substituent selected from phenyl, phenyl substituted with 1, 2 or 3 halo substituents, pyridyl, pyrazole, phenoxy and phenoxy
20 substituted with 1, 2 or 3 halo substituents; pyridyl; or pyridyl substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl and C₁-C₄ alkyl and/or with 1 substituent selected from phenyl and phenyl substituted with 1, 2 or 3 halo substituents. More preferably -Ar₁ is phenyl or phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl and C₁-C₄ alkyl
25 and/or with 1 substituent selected from phenyl, phenyl substituted with 1, 2 or 3 halo substituents, pyridyl, pyrazole, phenoxy and phenoxy substituted with 1, 2 or 3 halo substituents. Most preferably -Ar₁ is phenyl substituted with 1 or 2 substituents each independently selected from halo, trifluoromethyl and C₁-C₄ alkyl and/or with 1 substituent selected from phenyl, phenyl substituted with 1, 2 or 3 halo substituents,
30 pyridyl, pyrazole, phenoxy and phenoxy substituted with 1, 2 or 3 halo substituents. Suitable -Ar₁ groups include, for example, 2-methylthiophenyl, 2-methylphenyl, 2-

fluorophenyl, 2-chlorophenyl, 2-isopropoxyphenyl, 2-trifluoromethylphenyl, 2-difluoromethoxyphenyl, 2-methoxyphenyl, 2-ethoxyphenyl, 2-(1,1'-biphenyl), 2-phenoxyphenyl, 2-benzylphenyl, 3-trifluoromethoxyphenyl, 3-chlorophenyl, 3-trifluoromethylphenyl, 3-methylphenyl, 3-trifluorothiomethoxyphenyl, 3-methoxyphenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 4-fluorophenyl, 3,5-dichlorophenyl, 3,5-dimethylphenyl, 3-trifluoromethyl-5-fluorophenyl, 3,5-difluorophenyl, 2,3-dichlorophenyl, 2,3-dimethylphenyl, 2-chloro-3-trifluoromethylphenyl, 2-chloro-3-methylphenyl, 2-methyl-3-chlorophenyl, 2,4-dichlorophenyl, 2,4-dimethyl, 2,4-difluorophenyl, 2-chloro-4-fluorophenyl, 2-trifluoromethyl-4-fluorophenyl, 2-fluoro-4-trifluoromethylphenyl, 2-methyl-4-chlorophenyl, 2-methoxy-4-fluorophenyl, 2-trifluoromethyl-5-fluorophenyl, 2,5-dimethylphenyl, 4-fluoro-[1,1'-biphenyl]-2-yl, 2-chloro-5-fluorophenyl, 2-(trifluoromethyl)-6-fluorophenyl, 2-chloro-6-fluorophenyl, 3,4-dichlorophenyl, and 3-chloro-4-fluorophenyl. In general when $-Ar_1$ is phenyl substituted with pyridyl, 3-pyridyl is preferred.


Preferred compounds of formula (IF) are those wherein $-Ar_1$ is pyridyl or pyridyl substituted with 1, 2 or 3 substituents each independently selected from halo, trifluoromethyl and C_1 - C_4 alkyl and/or with 1 substituent selected from phenyl and phenyl substituted with 1, 2 or 3 halo substituents. More preferably $-Ar_1$ is pyridyl substituted with 1 or 2 substituents each independently selected from halo, trifluoromethyl and C_1 - C_4 alkyl and/or with 1 substituent selected from phenyl and phenyl substituted with 1, 2 or 3 halo substituents. Suitable $-Ar_1$ groups include, for example, 3-phenyl-2-pyridyl. In general when $-Ar_1$ is a substituted pyridyl, substituted 2-pyridyl is preferred.

10. A compound of formula (IG)



(IG)

wherein -X- is -S- or -O-; each R is independently selected from H or C₁-C₄ alkyl; R¹ is H, C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, cyano, trifluoromethyl, trifluoromethoxy, -NR³R⁴, -CONR³R⁴, -COOR³ or a group of the formula (i)



- 5 R² is C₁-C₄ alkyl, phenyl or phenyl substituted with 1, 2 or 3 substituents each independently selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, nitro, hydroxy, cyano, halo, trifluoromethyl, trifluoromethoxy, benzyl, benzyloxy, -NR⁶R⁷, -CONR⁶R⁷, COOR⁶, -SO₂NR⁶R⁷ and -SO₂R⁶; R⁵ is selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, carboxy, nitro, hydroxy, cyano, halo, trifluoromethyl, trifluoromethoxy, benzyl, benzyloxy, -NR⁸R⁹, -CONR⁸R⁹, -SO₂NR⁸R⁹ and -SO₂R⁸; R³, R⁴, R⁶, R⁷, R⁸ and R⁹ are each independently selected from H or C₁-C₄ alkyl; and -Z- is a bond, -CH₂-, or -O-; or a pharmaceutically acceptable salt thereof.

With respect to formula (IG) the term "C₁-C₄ alkyl" means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 15 1 to 4 carbon atoms. Thus the term "C₁-C₄ alkyl" includes methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl.

With respect to formula (IG) the term "C₁-C₄ alkoxy" means a monovalent unsubstituted saturated straight-chain or branched-chain hydrocarbon radical having from 1 to 4 carbon atoms linked to the point of substitution by an O atom. Thus the term "C₁-C₄ alkoxy" includes methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy.

With respect to formula (IG) the term "halo" or "halogen" means F, Cl, Br or I.

Preferred compounds of formula (IG) are those wherein -X- is -S-.


Preferred compounds of formula (IG) are those wherein -X- is -O-.

25 Preferred compounds of formula (IG) are those wherein R² is phenyl.

Preferred compounds of formula (IG) are those wherein all R groups are hydrogen.


Preferred compounds of formula (IG) are those represented by the formula (IIG)

53



(IIG)

wherein R^1 is H, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, halo, cyano, trifluoromethyl, trifluoromethoxy, $-NR^3R^4$, $-CONR^3R^4$, $-COOR^3$ or a group of the formula (i)



(i)

R^5 is selected from C_1 - C_4 alkyl, C_1 - C_4 alkoxy, carboxy, nitro, hydroxy, cyano, halo, trifluoromethyl, trifluoromethoxy, benzyl, benzyloxy, $-NR^8R^9$, $-CONR^8R^9$, $-SO_2NR^8R^9$ and $-SO_2R^8$; R^3 , R^4 , R^8 and R^9 are each independently selected from H or C_1 - C_4 alkyl; $-Z-$ is a bond, $-CH_2-$, or $-O-$; or a pharmaceutically acceptable salt thereof.

Preferred compounds of formula (IG) or (IIG) are those wherein the substituent R^1 is in the three position of the pyridine ring as numbered in formula (IG) above. More preferably said substituent R^1 is H, C_1 - C_4 alkyl, halo, cyano, $-CONR^3R^4$, trifluoromethyl or a group of the formula (i). When R^1 is $-CONR^3R^4$, then R^3 and R^4 are both preferably H. When R^1 is C_1 - C_4 alkyl, then it is preferably methyl.

Preferred compounds of formula (IG) or (IIG) are those wherein the substituent R^1 is a group of the formula (i).

Preferred compounds of formula (IG) or (IIG) are those wherein R^1 is a group of the formula (i), $-Z-$ is a bond, and R^5 is H or halo.

Preferred compounds of formula (IG) or (IIG) are those wherein R^1 is a group of the formula (i), $-Z-$ is $-CH_2-$ or $-O-$, and R^5 is H.

Preferred compounds of formula (IG) or (IIG) are those wherein the substituent R^1 is in the five position of the pyridine ring as numbered in formula (IG) above. More preferably said substituent R^1 is selected from bromo, chloro or iodo.

Compounds within the scope of Formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above are inhibitors of norepinephrine reuptake. Certain compounds within the scope of Formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above are selective inhibitors of norepinephrine reuptake.

5 Biogenic amine transporters control the amount of biogenic amine neurotransmitters in the synaptic cleft. Inhibition of the respective transporter leads to a rise in the concentration of that neurotransmitter within the synaptic cleft. Compounds of Formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above and their pharmaceutically acceptable salts preferably exhibit a K_i value less than 500nM at the norepinephrine
10 transporter as determined using the scintillation proximity assay as described below. More preferred compounds of Formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above and their pharmaceutically acceptable salts exhibit a K_i value less than 100nM at the norepinephrine transporter. More preferred compounds of Formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above and their pharmaceutically acceptable salts exhibit a K_i value
15 less than 50nM at the norepinephrine transporter. Especially preferred compounds of Formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above and their pharmaceutically acceptable salts exhibit a K_i value less than 20nM at the norepinephrine transporter. Preferably, these compounds selectively inhibit the norepinephrine transporter relative to the serotonin and dopamine transporters by a factor of at least five, more preferably by a
20 factor of at least ten.

In addition, the compounds of Formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above of the present invention are preferably acid stable. Advantageously, they have a reduced interaction (both as substrate and inhibitor) with the liver enzyme Cytochrome P450 (CYP2D6). That is to say, they preferably exhibit less than 75% metabolism via the
25 CYP2D6 pathway according to the CYP2D6 substrate assay described below and they preferably exhibit an IC_{50} of $>6\mu M$ according to the CYP2D6 inhibitor assay described below.

While all compounds exhibiting norepinephrine reuptake inhibition are useful for the methods of the present invention, certain are preferred. It is preferred that the
30 norepinephrine reuptake inhibitor is selective for the reuptake of norepinephrine over the

reuptake of other neurotransmitters. It is also preferred that the norepinephrine reuptake inhibitor does not exhibit significant direct agonist or antagonist activity at other receptors. It is especially preferred that the norepinephrine reuptake inhibitor be selected from atomoxetine, reboxetine, (S,S)-reboxetine, (R)-N-methyl-3-(2-methyl-thiophenoxy)-
5 3-phenylpropylamine, and compounds of Formulae (I), (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above.

The present invention encompasses pharmaceutical compositions comprising the compounds disclosed herein, or pharmaceutically acceptable salts thereof, together with a pharmaceutically acceptable carrier, diluent, or excipient.

10 It will be understood by the skilled reader that most or all of the compounds used in the present invention are capable of forming salts, and that the salt forms of pharmaceuticals are commonly used, often because they are more readily crystallized and purified than are the free bases. In all cases, the use of the pharmaceuticals described above as salts is contemplated in the description herein, and often is preferred, and the
15 pharmaceutically acceptable salts of all of the compounds are included in the names of them.

Many of the compounds used in this invention are amines, and accordingly react with any of a number of inorganic and organic acids to form pharmaceutically acceptable acid addition salts. Since some of the free amines of the compounds of this invention are
20 typically oils at room temperature, it is preferable to convert the free amines to their pharmaceutically acceptable acid addition salts for ease of handling and administration, since the latter are routinely solid at room temperature. Acids commonly employed to form such salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids, such as p-
25 toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-bromophenylsulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid and the like. Examples of such pharmaceutically acceptable salts thus are the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate,
30 caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-

1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, b-hydroxybutyrate, glycollate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the
5 like. Preferred pharmaceutically acceptable salts are those formed with hydrochloric acid.

Pharmaceutically acceptable salts of the compounds of Formulae (IA), (IB), (IC), (ID) (IE), (IF) and (IG) above include acid addition salts, including salts formed with inorganic acids, for example hydrochloric, hydrobromic, nitric, sulphuric or phosphoric
10 acids, or with organic acids, such as organic carboxylic or organic sulphonic acids, for example, acetoxybenzoic, citric, glycolic, *o*- mandelic-l, mandelic-dl, mandelic d, maleic, mesotartaric monohydrate, hydroxymaleic, fumaric, lactobionic, malic, methanesulphonic, napsylic, naphthalenedisulfonic, naphtoic, oxalic, palmitic, phenylacetic, propionic, pyridyl hydroxy pyruvic, salicylic, stearic, succinic, sulphanilic,
15 tartaric, 2-hydroxyethane sulphonic, toluene-p-sulphonic, and xinafoic acids.

In addition to the pharmaceutically acceptable salts, other salts can serve as intermediates in the purification of compounds, or in the preparation of other, for example pharmaceutically acceptable, acid addition salts, or are useful for identification, characterization, or purification.

20 The present invention encompasses the administration of a composition that exhibits (preferably selective) norepinephrine reuptake inhibitor activity. The composition can comprise one or more agents that, individually or together, inhibit norepinephrine reuptake preferably in a selective manner.

25 Dosages

The dosages of the drugs used in the methods of the present invention must, in the final analysis, be set by the physician in charge of the case using knowledge of the drugs, the properties of the drugs alone or in combination as determined in clinical trials, and the characteristics of the patient including diseases other than that for which the physician is
30 treating the patient. General outlines of the dosages, and some preferred dosages, are as follows:

intravenous, or intradermal administration), intra-pulmonary, vaginal, rectal, intranasal, ophthalmic, or intraperitoneal administration, or by an implantable extended release device. Oral administration is preferred. The route of administration can be varied in any way, limited by the physical properties of the drugs, the convenience of the patient and the caregiver, and other relevant circumstances (*Remington's Pharmaceutical Sciences* (1990) 18th Edition, Mack Publishing Co.).

The pharmaceutical compositions are prepared in a manner well known in the pharmaceutical art. The carrier or excipient can be a solid, semi-solid, or liquid material that can serve as a vehicle or medium for the active ingredient. Suitable carriers or excipients are well known in the art. The pharmaceutical composition can be adapted for oral, inhalation, parenteral, or topical use and can be administered to the patient in the form of tablets, capsules, aerosols, inhalants, suppositories, solutions, suspensions, or the like.

The compounds of the present invention can be administered orally, for example, with an inert diluent or capsules or compressed into tablets. For the purpose of oral therapeutic administration, the compounds can be incorporated with excipients and used in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums and the like. These preparations should contain at least 4% of the compound of the present invention, the active ingredient, but can be varied depending upon the particular form and can conveniently be between 4% to about 70% of the weight of the unit. The amount of the compound present in compositions is such that a suitable dosage will be obtained. Preferred compositions and preparations according to the present invention can be determined by a person skilled in the art.

The tablets, pills, capsules, troches, and the like can also contain one or more of the following adjuvants: binders such as microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch or lactose, disintegrating agents such as alginic acid, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; and sweetening agents such as sucrose or saccharin can be added or a flavoring agent such as peppermint, methyl salicylate or orange flavoring. When the dosage unit form is a capsule, it can contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or a fatty oil.

Atomoxetine:

In adults and older adolescents: from about 5 mg/day to about 200 mg/day; preferably in the range from about 60 to about 150 mg/day; more preferably from about 60 to about 130 mg/day; and still more preferably from about 50 to about 120 mg/day;

In children and younger adolescents: from about 0.2 to about 3.0 mg/kg/day; preferably in the range from about 0.5 to about 1.8 mg/kg/day;

Reboxetine: Racemic reboxetine can be administered to an individual in an amount in the range of from about 2 to about 20 mg per patient per day, more preferably from about 4 to about 10 mg/day, and even more preferably from about 6 to about 10 mg/day. Depending on the formulation, the total daily dosage can be administered in smaller amounts up to two times per day. A preferred adult daily dose of optically pure (S,S) reboxetine can be in the range of from about 0.1 mg to about 10 mg, more preferably from about 0.5 mg to about 8 to 10 mg, per patient per day. The effective daily dose of reboxetine for a child is smaller, typically in the range of from about 0.1 mg to about 4 to about 5 mg/day. Treatments using compositions containing optically pure (S,S)-reboxetine are about 5 to about 8.5 times more effective in inhibiting the reuptake of norepinephrine than compositions containing a racemic mixture of (R,R)- and (S,S)-reboxetine, and therefore lower doses can be employed. PCT International Publication No. WO 01/01973 contains additional details concerning the dosing of (S,S) reboxetine.

Compounds of formula I: from about 0.01 mg/kg to about 20 mg/kg; preferred daily doses are from about 0.05 mg/kg to 10 mg/kg; more preferably from about 0.1 mg/kg to about 5 mg/kg;

Compounds of formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above: from about 5 to about 500 mg, more preferably from about 25 to about 300 mg, of the active ingredient per patient per day.

Administration

The compounds disclosed herein can be administered by various routes, for example systemically via oral (including buccal or sublingual), topical (including buccal, sublingual, or transdermal), parenteral (including subcutaneous, intramuscular,

Other dosage unit forms can contain other various materials that modify the physical form of the dosage unit, for example, as coatings. Thus, tablets or pills can be coated with sugar, shellac, or other coating agents. A syrup can contain, in addition to the present compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

A formulation useful for the administration of R-(-)-N-methyl 3-((2-methylphenyl)oxy)-3-phenyl-1-aminopropane hydrochloride (atomoxetine) comprises a dry mixture of R-(-)-N-methyl 3-((2-methylphenyl)oxy)-3-phenyl-1-aminopropane hydrochloride with a diluent and lubricant. A starch, such as pregelatinized corn starch, is a suitable diluent and a silicone oil, such as dimethicone, a suitable lubricant for use in hard gelatin capsules. Suitable formulations are prepared containing about 0.4 to 26% R-(-)-N-methyl 3-((2-methylphenyl)oxy)-3-phenyl-1-aminopropane hydrochloride, about 73 to 99% starch, and about 0.2 to 1.0% silicone oil. Tables 1 and 2 illustrate particularly preferred formulations:

Table 1

Ingredient (%)	2.5 mg	5 mg	10 mg	18 mg	20 mg	25 mg	40 mg	60 mg
R-(-)-N-methyl 3- ((2-methylphenyl)oxy)-3- phenyl-1- aminopropane hydrochloride	1.24	2.48	4.97	8.94	9.93	12.4 2	19.8 7	22.1 2
Dimethicone	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Pregelatinized Starch	98.2 6	97.0 2	94.5 3	90.5 6	89.5 7	87.0 8	79.6 3	77.3 8

Table 2

Ingredient (mg/capsule)	2.5 mg	5 mg	10 mg	18 mg	20 mg	25 mg	40 mg	60 mg
R-(-)-N-methyl 3- ((2-meth- ylphenyl)oxy)-3- phenyl-1- aminopropane hydrochloride	2.86	5.71	11.4 3	20.5 7	22.8 5	28.5 7	45.7 1	68.5 6
Dimethicone	1.15	1.15	1.15	1.15	1.15	1.15	1.15	1.55
Pregelatinized Starch	225. 99	223. 14	217. 42	208. 28	206. 00	200. 28	183. 14	239. 89
Capsule Fill Weight (mg)	230	230	230	230	230	230	230	310
Capsule Size	3	3	3	3	3	3	3	2

- 5 For the purpose of parenteral therapeutic administration, the compounds of the present invention can be incorporated into a solution or suspension. These preparations typically contain at least 0.1% of a compound of the invention, but can be varied to be between 0.1 and about 90% of the weight thereof. The amount of the compound of formula I present in such compositions is such that a suitable dosage will be obtained.
- 10 The solutions or suspensions can also include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylene diaminetetraacetic acid; buffers such as
- 15 acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Preferred compositions and preparations are able to be determined by one skilled in the art.

The compounds of the present invention can also be administered topically, and when done so the carrier can suitably comprise a solution, ointment, or gel base. The base, for example, can comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bees wax, mineral oil, diluents such as water and alcohol, and emulsifiers, and stabilizers. Topical formulations can contain a concentration of the compound, or its pharmaceutical salt, from about 0.1 to about 10% w/v (weight per unit volume).


The compositions are preferably formulated in a dosage unit form, i.e., physically discrete units suitable as unitary doses for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier, diluent, or excipient.

The following examples are provided to illustrate various aspects of the present invention, and should not be construed to be limiting thereof in any way.

Preparation of Compounds of Formula (IA)

Compounds of formula (IA) may be prepared by conventional organic chemistry techniques and also by solid phase synthesis. In the present specification the abbreviation "boc" refers to the N-protecting group t-butyloxycarbonyl. In the present specification the abbreviation "TFA" refers to trifluoroacetic acid. In the present specification the abbreviation "DMF" refers to dimethylformamide. In the present specification the abbreviation "SPE" refers to solid phase extraction. In the present specification the abbreviation "ACE-Cl" refers to α -chloroethyl chloroformate.

When R₈ is H, a suitable three-step conventional synthesis is outlined in **Scheme 1A** shown below.

**Scheme 1A**


A Boc-protected 4-piperidone (IIA) is reductively aminated with an amine to
 5 provide a 4-amino-piperidine (IIIAa or IIIAb). A second reductive amination with an

aldehyde or ketone provides a boc-protected compound of formula (IA) (IVA). The boc group is removed under acidic conditions to provide a compound of formula (IA) (where R₈ is H). If desired, the compound of formula (IA) (where R₈ is H) may be converted to a suitable salt by addition of a suitable quantity of a suitable acid. In the schemes above
5 (and below) R₁ to R₇, R₉, R₁₀ and n are as previously defined, m is 0, 1 or 2 and R₁₁ and R₁₂ are chosen such that R₁₁-CH-R₁₂ = R₁.

Although the boc N-protecting group is used in the above illustration, it will be appreciated that other N-protecting groups (for example acetyl, benzyl or benzyloxycarbonyl) could also be used together with a deprotection step appropriate for the
10 N-protecting group used. Similarly, other reducing agents (for example NaBH₄ or LiAlH₄) may be used in the reductive amination steps and other acids (for example HCl) may be used in the deprotection step.


As an alternative to the second reductive amination step, compound IIIAa or IIIAb may be subjected to an alkylation step as shown in **Scheme 1B** below (L represents a
15 suitable leaving group – for example Br or tosyl).

64

**Scheme 1B**

Once again, N-protection other than boc may also be used together with a suitable
 5 deprotection step. Similarly, bases other than potassium carbonate (e.g NaH) may be used
 for the alkylation step

Using essentially the same chemical reactions as in the first scheme above, the
 compounds of formula (IA) (where R8 is H) may also be prepared by a solid phase
 parallel synthesis technique as outlined in **Scheme 1C** shown below.



Scheme 1C


A piperidone hydrate is attached to a polystyrene resin to provide a resin bound piperidone (VA). Aliquots are reductively aminated to provide a resin bound secondary amine (VIA) that can undergo a further reductive amination with an aldehyde or ketone to give the tertiary amine (VIIA). Acidic cleavage from the resin and SPE provides

compounds of formula (IA) (where R₈ is H) which may be purified by ion exchange methods using, for example, the SCX-2 ion exchange resin.

Although NaBH(OAc)₃ is used in the above illustration, it will be appreciated that other reducing agents (for example NaBH₄ or LiAlH₄) may be used in the reductive amination steps and other acids (for example HCl) may be used in the deprotection step. Solid phase resins other than the p-nitrophenylcarbonate-polystyrene resin illustrated above may also be employed.

When R₈ is C₁-C₄alkyl, a conventional synthetic route is outlined in **Scheme 1D** shown below.

10




Scheme 1D

A benzyl-protected 4-piperidone (VIII A) is alkylated with an alkyl lithium reagent to provide a 4-amino-piperidinol (IX A). Treatment with an alkylnitrile or alkylamide under strongly acidic conditions provides a secondary amide (X A) which may be deprotected, boc-protected and reduced to provide a secondary amine (XIA). Alkylation of the secondary amine (XIA) followed by removal of the boc group provides a compound of formula (IA) (where R₈ is C₁-C₄alkyl). In the scheme above L is a leaving group as previously defined and R₁₃ is chosen such that R₁₃-CH₂ = R₁.

Although the benzyl and boc N-protecting groups are used in the above illustration, it will be appreciated that other N-protecting groups could also be used in their place together with deprotection steps appropriate for those N-protecting groups. Similarly, other reducing agents may be used in the amidocarbonyl reduction step and other organometallics or bases may be used in the respective alkylation steps.

Preparation of Compounds of Formula (IB)


A general scheme outlining the synthetic routes to compounds of Formulae (IB) wherein Y is OH is shown below (**Scheme 1B**). For clarity, Ar₂ is shown as phenyl and R_y and R_z are shown as H. It will be appreciated that analogous methods could be applied for other possible identities of Ar₂, R_y and R_z.




Scheme 1B

Compounds of Formulae (IB) can be prepared by conventional organic chemistry techniques from an *N*-benzyl-ketomorpholine of type **1B** by addition of a suitable organometallic derivative (method A), or *via* the addition of a suitable organometallic reagent to an epoxide of type **2B** (method B), as outlined in **Scheme 1B**.


The racemic intermediates of type **1B** can be obtained as outlined in **Scheme 2B** by condensation of an *N*-benzyl cyanomorpholine **5B** (*J. Med. Chem.* 1993, 36, pp 683 – 689) with a suitable aryl organometallic reagent followed by acid hydrolysis. Chiral HPLC separations of the racemic *N*-benzyl-aryl-ketomorpholine of type **1B** gives the required single enantiomer, i.e., the (2*S*)-*N*-benzyl-aryl-ketomorpholine of type **6B** (**Scheme 2B**).

**Scheme 2B**

Condensation of a chiral (2*S*)-*N*-benzyl-aryl-ketomorpholine of type **6B** with a commercially available benzylic magnesium halide or a benzylic magnesium halide prepared using standard Grignard techniques from the corresponding halo-benzylic derivative gives a tertiary alcohol of type **3B** without any observed epimerisation of the existing asymmetric center (ee's/de's determinations can be carried out using chiral HPLC) and with very high overall diastereoisomeric excesses (see **Scheme 3B**). The final compounds of type **4B** can be obtained after cleavage of the *N*-benzyl protecting group on a compound of type **3B**. The deprotection can be done using catalytic palladium hydrogenolysis, or carbamate exchange with ACE-Cl (1-Chloroethyl chloroformate), giving intermediates of type **7B**, followed by methanolysis as shown in **Scheme 3B**.



The intermediates **3B** can be further elaborated using for example organometallic type couplings between an ortho bromide derivative of type **8B** and an arylboronic acid as shown in **Scheme 4B**. For clarity, Ar₁ and its substituent (R₁) are shown as phenyl and substitution occurs at the 2-position. It will be appreciated that analogous methods could be applied for other possible identities of Ar₁ and R₁ and other possible substitution positions. This approach can also be carried out by solid phase synthetic methods as described in more detail in the specific examples below.




10

15

An alternative route for the preparation of the compounds of Formulae (IB) is method B (see **Scheme 1B**). Formation of the intermediate epoxides of type **2B** from racemic *N*-benzyl-ketomorpholines of type **1B**, can be done using for example trimethyl sulfoxonium iodide and a suitable base, for example sodium hydride. Condensation of **2B** with a commercially available aryl organometallic, or an aryl organometallic prepared

from the corresponding halo aryl derivative, gives the intermediates of type **3B**, as mixtures of diastereoisomers. Final deprotections can be done as described above (see **Scheme 3B**). Final compounds made using method B can be purified using chiral HPLC.

- Compounds of Formula (IB) of the present invention wherein Y is OR and R is C₁-C₄ alkyl, can be synthesized by standard alkylation of intermediates of type **3B** prior to deprotection of the morpholine N-atom as shown in **Scheme 5B**. Suitable strong bases will be known to the person skilled in the art and include, for example, sodium hydride. Similarly, suitable alkylating agents will be known to the person skilled in the art and include, for example, C₁-C₄ alkyl halides such as methyl iodide.




Scheme 5B

Preparation of Compounds of Formula (IC)


- Compounds of formula (IC) may be prepared by conventional organic chemistry techniques from N-benzyl-cyanomorpholine **1C** (Route A) or N-benzyl-morpholinone **2C** (Route B) as outlined in **Scheme 1C** below: For clarity, X is shown as phenyl and R' and R¹ are shown as H. It will be appreciated that analogous methods could be applied for other possible identities of X, R' and R¹.

71



Scheme 1C

More detail of Route A is given in Scheme 2C:



Scheme 2C


5

The amino alcohol **4Ca** can be obtained by reaction of N-benzyl-cyanomorpholine **1C** with a Grignard reagent, followed by acid hydrolysis to give racemic phenyl ketone **3C** which may be separated on chiral HPLC. (2S)-Phenyl ketone **3Ca** may then be reduced with DIP-Cl to give **4Ca** in high diastereomeric excess. The amino alcohol **4Ca** is converted into benzyl bromide **5Ca**, to give the desired N-substituted aryl thio morpholines after displacement with the requisite aryl thiol. N-substituted aryloxy morpholines may be obtained in an analogous manner by displacement with the requisite hydroxyaryl compound. Alternatively, N-substituted aryloxy morpholines may be

10

obtained by addition of a strong base, such as sodium hydride, to the amino alcohol **4Ca** to form a nucleophilic alkoxide followed by an S_NAr reaction with an Ar group substituted with a suitable leaving group (e.g. F). Deprotection of the tertiary amine gives the final products.

5 Detail of route B is given in **Scheme 3C**:




Scheme 3C

Treatment of *N*-benzyl morpholinone **2C** with a strong base such as lithium diisopropylamide at low temperature followed by addition of benzaldehyde gives aldol adducts **6Ca-6Cd** as a 2:1 mixture of diastereomer pairs **6Ca,6Cb** and **6Cc,6Cd**, which may be separated using conventional chromatographic techniques. Reduction with a borane reagent at elevated temperatures gives diastereomeric amino alcohol pairs **4Ca,4Cb** and **4Cc,4Cd** respectively.

Amino alcohol pair **4Ca,4Cb** may be converted to bromide **5Ca,5Cb** and further to racemic aryl thio morpholines as outlined in **Scheme 4C**. Amino alcohol pair **4Cc,4Cd** may be converted into the corresponding mesylate. Displacement with the requisite thiol, followed by removal of the nitrogen protecting group furnishes aryl thiol morpholines as racemic mixtures of two diastereomers. The racemic aryl thiol morpholines may be separated into enantiomerically pure products using chiral HPLC technology. *N*-substituted aryloxy morpholines may be obtained in an analogous manner by displacement with the requisite hydroxyaryl compound.


73



Scheme 4C

Aryl-substituted morpholines **33C**, **35C**, **37C** may be obtained from morpholinone **2C** as outlined in Scheme 5C:


5



Scheme 5C

An alternative route to **9C** is outlined in Scheme 6C. This route makes use of a chiral auxiliary and gives **9C** in enantiomerically pure form.

74




Scheme 6C

Preparation of Compounds of Formula (ID)

- 5 Compounds of formula (ID) may be prepared using the following methods. General schemes outlining the synthetic routes used to prepare racemic products are given below. All active racemates may be separated into single enantiomers using chiral HPLC and may be readily converted into suitable salts.


10 Compounds of formula (ID) wherein Ar is (i) and R^{2c} is H may be prepared as shown in **Scheme 1D** below:



Quinolin-2-one **1D** or its corresponding 4-oxo and 4-thio derivatives can be N-arylated using modified conditions to those reported by Buchwald, (*J. Am. Chem. Soc.*, 123, **2001**, p. 7727). For example the quinolin-2-one **1D** is reacted with 3 equivalents of Ar-Br wherein Ar is (i) and R^{2c} is H, 0.2 equivalents of trans-cyclohexanediamine, 0.2 equivalent of copper iodide (CuI), 2.1 equivalents of potassium carbonate (K₂CO₃), in an organic solvent such as 1,4-dioxane at a temperature of 125°C overnight. The resulting N-arylated quinolin-2-one **2D** can be alkylated by treatment with a strong base such as lithium hexamethyldisilazide (LiHMDS) at temperatures of -78°C in a suitable organic solvent such as tetrahydrofuran (THF), followed by the addition of an alkyl halide such as alkyl iodide to give the corresponding 3-alkylated-N-arylated quinolin-2-one derivative **3D**. Using the same alkylating conditions above with a 1,2-dihaloethane, such as 1-

bromo-2-chloroethane, or a 1,3-dihalopropane, such as 1-bromo-3-chloropropane, as alkylating agents provides **4D** or **5D** wherein n is 2 or 3 respectively. These halo analogues were chosen as ideal precursors to the desired amine products. For instance, treatment of **4D** or **5D** with aqueous methylamine, in the presence of a catalytic amount of a suitable iodide, such as potassium iodide (KI), in ethanol at 100°C provided the racemic amine products **6D** and **7D** respectively, in moderate yields.

Compounds of formula (ID) wherein Ar is (i), R^{2c} is H and n is 3 may be prepared using alternative chemistry as shown in **Scheme 2D**.



Scheme 2D

10

Quinolin-2-ones **2D** and **3D** can be alkylated using the aforementioned alkylating procedure using an allyl halide e.g. allyl bromide as the alkylating agent to give the corresponding 3-allyl-N-arylated-quinolin-2-ones **11D**. Said allyl analogues could then be converted to the corresponding primary alcohols **12D** by a hydroboration procedure involving a suitable borane, such as 9-BBN in a suitable solvent such as THF. Oxidative work up using for example reaction conditions such as aqueous hydrogen peroxide in a solvent such as ethanol, in the presence of a suitable base, such as sodium hydroxide, gave moderate to good yields of alcohol products after column chromatography


15

purification. The alcohols were cleanly converted into their mesylates, by reaction of a mesyl halide such as mesyl chloride in the presence of a suitable base such as triethylamine in a suitable solvent such as THF at a suitable temperature such as 0°C to room temperature. The resulting mesylates are used directly in the amination step

described above in **Scheme 1D** to provide good yields of the final racemic targets **13D**.

In order to prepare a range of N-arylated analogues advanced intermediates were prepared that could undergo N-arylations with a range of substituted aryl halides, such as aryl bromides or iodides, 2 and 3-halothiophenes, 2 and 3-halofurans or 2 and 3-halopyrroles. The synthetic route used to prepare intermediates **19D** is shown below in

Scheme 3D.




Scheme 3D

Compounds of formula (ID) wherein n is 3 may be prepared as shown in **Scheme 3D**. This method is particularly suitable for compounds wherein Ar is (i) and R^{2c} is H or Ar is (ii), wherein -Y- is -S-.

Quinolin-2-one **1D** can be protected using a suitable amide-protecting group such as those described in T.W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1991, hereafter referred to as "Greene". For example quinolin-2-one **1D** can be protected with a 4-methoxybenzyl group. The protection reaction can be carried out for example using a suitable base, such as sodium hydride in a suitable solvent, such as dimethylformamide, followed by reaction with a 4-methoxybenzyl halide, such as 4-methoxybenzyl chloride, to give the corresponding N-protected derivative **14D** in good yield. This intermediate can be converted directly to the allyl analogue **16Da**, wherein R¹ = H, in a manner described earlier or converted into the alkyl analogue **15D** which can be subsequently alkylated with a allyl halide to give the allyl analogue **16Db**, wherein R¹ is C₁-C₄ alkyl. Using the same hydroboration, mesylation and amination sequence described in **Scheme 2D** provided both amines **18Da-b**. Deprotection of protected quinolin-2-one could be achieved using any suitable deprotection conditions as those shown in Greene. For example, the 4-methoxybenzyl group could be cleaved cleanly using trifluoroacetic acid and anisole at 65°C. The resultant product could be selectively protected on the secondary amine with a suitable nitrogen protecting group as those described in Greene. For example, the secondary amine can be protected with a Boc group. The reaction can be carried out with Boc anhydride in a suitable solvent such as THF to provide multi gram quantities of **19Da-b**. Reaction of **19Da-b** with various aryl bromides using the previously described N-arylation conditions, deprotection using suitable deprotecting conditions such as those described in Greene gave a range of final racemic targets **21Da-b** or **22Da-b**. For example, for compounds protected with a Boc group they can be deprotected in the presence of trifluoroacetic acid (TFA) in a suitable organic solvent such as dichloromethane (DCM).


Intermediates **19Da-b** wherein R³ is a halo group, for example chloro or bromo, can be used to provide compounds of formula (ID) wherein R³ is a phenyl group, such as compound **24D**, via a Suzuki coupling, see **Scheme 4D** below.




Intermediates **19Da-b**, wherein R^3 is for example bromo can be N-protected with a suitable amide protecting group for example 4-methoxybenzyl as described in **Scheme 3D** above and then coupled with phenylboronic acid under Suzuki conditions to provide the phenyl analogues **23D**. Deprotection of the 4-methoxybenzyl group with TFA, followed by protection of the resulting secondary amine with a suitable nitrogen protecting group such as Boc followed by subsequent N-arylation and Boc deprotection using the previously described methodology gave the final target **24D**.

It will be appreciated that compounds of formula (IDa) wherein R^3 is bromo or chloro can be prepared as shown in **Schemes 1D to 4D** above starting from the corresponding haloquinolin-2-ones. Alternatively, they can be prepared from the corresponding quinolin-2-one **1D** wherein R^3 is hydrogen as mentioned above including an extra step comprising the halogenation of a suitable intermediate at some stage of the synthesis. For example quinolin-2-one **1D** in **Scheme 2D** can be halogenated using N-chlorosuccinimide in a suitable solvent such as DMF at a suitable temperature such as room temperature to give the corresponding 6-chloro-quinolin-2-one **1D** wherein R^3 is Cl.

Alternatively intermediates (**19Da-b**) wherein R^3 is H in **Scheme 3D** can be halogenated in the presence of N-chloro and N-bromosuccinimide in a suitable solvent such as DMF to give the corresponding 6-chloro and 6-bromoquinolin-2-ones (**20Da-c**).

**20Da-c**


It will be appreciated that **Schemes 1D to 4D** above relate to methods for the preparation of compounds of formula (ID) wherein Ar is (i) and R^{2c} is hydrogen. Compounds of formula (ID) wherein Ar is (i) and R^{2c} can be other than hydrogen, can be prepared using any of the general methods mentioned above, starting from the corresponding N-arylated quinolin-2-one **27D**. A general method for preparing said intermediates is illustrated in **Scheme 5D**. Commercially available 3-(2-Bromo-phenyl)-propionic acids **25D** can be converted to amide **26D** using standard amide coupling conditions and converted to the N-arylated quinolin-2-ones **27D** by an intramolecular, palladium catalysed cyclisation according to the method of Buchwald et al (Tetrahedron, 1996, 52, p. 7525).

**Scheme 5D**

Preparation of Compounds of Formula (IE)

Compounds of formula (IE) may be prepared by conventional organic chemistry techniques and also by solid phase synthesis. Compounds of formula (IE) can be prepared via the 3-aminopyrrolidine intermediate of formula (IVE) as illustrated in the

5 Scheme 1E below:

**Scheme 1E**

Commercially available 3-hydroxypyrrolidine of formula (IIIE) wherein R² is hydrogen, can be protected using a suitable nitrogen-protecting group such as those


10 described in T.W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1991, hereafter referred to as "Greene". For example 3-R-

hydroxypyrrolidine (IIIE) can be protected with a tert-butoxycarbonyl group, (boc). The protection reaction can be carried out for example using Boc anhydride in a suitable solvent such as for example tetrahydrofuran (THF) or dichloromethane (DCM) in the presence of a base such as triethylamine (TEA) or 4-(dimethylamino)pyridine (DMAP).

- 5 It will be appreciated that for compounds of formula (IE) wherein R^2 is C_1 - C_2 alkyl, the 3-hydroxypyrrolidine of formula (IIIE) can be prepared from the readily available 3-pyrrolidinone via addition of the appropriate C_1 - C_2 alkyl organometallic. The hydroxy group of the N-protected-3-hydroxypyrrolidine can be converted into a suitable leaving group (L) such as for example chloride, bromide, iodide or mesylate. For example the N-
- 10 protected-hydroxypyrrolidine can be converted to the mesylate in the presence of mesyl chloride and a suitable base such as triethylamine in a solvent such as DCM. Said mesylate is subsequently displaced with the corresponding azide in a suitable solvent such as dimethylformamide (DMF) or dimethylsulphoxide (DMSO). This azide intermediate can be converted to the corresponding N-protected-aminopyrrolidine of formula (IVE) via
- 15 hydrogenation in the presence of a suitable catalyst such as Palladium on charcoal and in a suitable solvent such as methanol or ethanol.


- For compounds of formula (IE) wherein R^4 is H, intermediate (IVE) can be alkylated via reductive alkylation with a ketone of formula R^3 -CO-Ar₁ wherein R^3 and Ar₁ have the values for formula (IE) above. The reductive alkylation can be carried out
- 20 for example as a hydrogenation reaction in the presence of a suitable catalyst such as Palladium on charcoal and a suitable solvent such as for example ethanol. Alternatively, said reductive alkylation can be carried out in the presence of a suitable borane such as sodium triacetoxyborohydride, NaBH(OAc)₃ and optionally in the presence of a suitable acid such as acetic acid, in a suitable solvent such as for example dichloroethane (DCE).

- 25 Alternatively, intermediate of formula (VE) wherein R^4 is H can be prepared as shown in **Scheme 2E** below by reductive alkylation of readily available 3-aminopyrrolidine of formula (VIE) wherein R^2 has the values defined for formula (IE) above, followed by the protection of the nitrogen in the pyrrolidine ring using a suitable protecting group such as those defined in Greene.

**Scheme 2E**


For example the reductive alkylation can be carried out in the presence of a ketone of formula Ar_1-CO-R^3 wherein Ar_1 and R^3 have the values defined for formula (IE) above. Initial condensation of the amino pyrrolidine with the ketone is undertaken in the presence of a suitable acid such as p-toluenesulphonic acid, in a suitable solvent such as toluene. The resultant imino pyrrolidine intermediate can then be protected with for example a boc group. The reaction can be carried out in the presence of boc anhydride and a suitable base such as DMAP, in a suitable solvent such as DCM. Said imine is reduced via hydrogenation in the presence of a suitable catalyst such as palladium on charcoal, in a suitable solvent such as ethanol to give the corresponding amine of formula (VE).

Intermediate of formula (VE) can be converted to compounds of formula (VIII E) via reductive alkylation with an aldehyde of formula R^9-CHO , wherein R^9 is chosen such that $R^9-CH_2 = R^1$ and R^1 has the values defined for formula (IE) above. The reductive alkylation can be carried out using standard methods, for instance as those mentioned above with the ketone Ar_1-CO-R^3 .

**Scheme 3E**


For example a compound of formula (VE) can be alkylated with R^9 -CHO in the presence of a suitable borane, such as $\text{NaBH}(\text{OAc})_3$, optionally in the presence of an acid such as acetic acid, in the presence of a suitable solvent such as dichloroethane (DCE).

For compounds of formula (IE) wherein R^3 and R^4 are hydrogen the alkylation of
 5 intermediate (VE) can be carried out with a compound of formula $\text{Ar}_1\text{CH}_2\text{L}_1$ wherein L_1 is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding intermediate of formula (VIII E)_a. It will be appreciated that the same reaction can be carried out using $\text{Ar}_1\text{-CR}^3\text{R}^4\text{-L}_1$ wherein R^3 and R^4 are $\text{C}_1\text{-C}_2$ alkyl.



Scheme 4E


Compounds of formula (IE) wherein R^1 is $-\text{CH}_2\text{-COO-(C}_1\text{-C}_2\text{ alkyl)}$ can be prepared by reacting intermediate (VE) with a compound of formula $\text{L}_2\text{-CH}_2\text{-COO-(C}_1\text{-C}_2$
 15 alkyl) wherein L_2 is a suitable leaving group such as for example bromo, chloro or iodo. Said reaction can be carried out in the presence of a suitable base such as sodium hydride, in a suitable solvent such as dimethylformamide.



Scheme 5E


Compounds of formula (IE) wherein R^1 is $-(\text{CH}_2)_m\text{-CF}_3$ can be prepared by reacting intermediate (VE) with a compound of formula $\text{HOOC-(CH}_2)_{m_1}\text{-CF}_3$, wherein m_1

is 0, 1 or 2. The acid may be activated as its anhydride or acyl chloride, and is reacted in the presence of a suitable base such as triethylamine and a catalytic amount of DMAP, in a suitable solvent such as DCM. The resulting amide can be reduced to the amine of formula (VIII)_e in the presence of a suitable borane. For example, for compounds wherein m is 1, the reduction can be carried out in the presence of BH₃-Me₂S borane-dimethyl sulphide complex, in a suitable solvent such as THF.



Scheme 6E


Compounds of formula (IE) wherein R¹ is -(C₁-C₆ alkylene)-OH can be prepared by reacting intermediate (VE) with an epoxide. For example for compounds wherein R¹ is -CH₂-C(CH₃)₂-OH, the intermediate of formula (VE) is reacted with 2,2-dimethyloxirane, in a suitable solvent such as aqueous ethanol.



Scheme 7E

Alternatively compounds of formula (IE) wherein R¹ is -(C₁-C₆alkylene)-OH can be prepared by reacting intermediate (VE) with an ω-haloalkanoate, such as methylbromoacetate, in the presence of a base such a sodium hydrogen carbonate in a solvent such as acetonitrile. The intermediate ester is then reacted with 2 equivalents of methyl magnesium bromide in THF to yield the tertiary alcohol(VIII)_d:


86



Scheme 8E

It will be appreciated that the **Scheme 8E** above applies to alkylene chains longer than -CH₂-.
5


Compounds of formula (IE) wherein R¹ is -C₂-C₆ alkenyl, -(CH₂)_n-S-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-O-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-O-(C₃-C₆ cycloalkyl), -(C₁-C₅ alkylene)-SO₂-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-OCF₃, or -(C₁-C₅ alkylene)-CN, can be prepared via alkylation of intermediate (VE) with a compound of formula L₂-C₂-C₆ alkenyl, L₂-(CH₂)_n-S-(C₁-C₃ alkyl), L₂-(C₁-C₅ alkylene)-O-(C₁-C₃ alkyl), L₂-(C₁-C₅ alkylene)-O-(C₃-C₆ cycloalkyl), L₂-(C₁-C₅ alkylene)-SO₂-(C₁-C₃ alkyl), L₂-(C₁-C₅ alkylene)-OCF₃, or L₂-(C₁-C₅ alkylene)-CN respectively, wherein L₂ is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding
10 intermediate of formula (VIII E)_e.
15



Scheme 9E


Compounds of formula (IE) wherein R¹ is a group of formula (i) can be prepared using the synthesis illustrated in **Scheme 10E** for compounds wherein R¹ is 4-tetrahydropyranyl. The compound of formula (IVE) can be alkylated via reductive alkylation using standard methods, as those mentioned above with the ketone Ar₁-CO-R³. For example compound of formula (IVE) can be alkylated with 4-tetrahydropyranone in
20

the presence of a suitable borane, such as sodium borohydride or $\text{NaBH}(\text{OAc})_3$, optionally in the presence of an acid such as acetic acid, in the presence of a suitable solvent such as dichloroethane (DCE). Then, the secondary amine can be alkylated with a compound of formula $\text{Ar}_1\text{CH}_2\text{L}_1$ wherein L_1 is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding intermediate of formula (VIII)_f. It will be appreciated that as mentioned above the same reaction can be carried out using $\text{Ar}_1\text{-CR}^3\text{R}^4\text{-L}_1$ wherein R^3 and R^4 are $\text{C}_1\text{-C}_2$ alkyl.




Scheme 10E

It will be appreciated that for compounds of formula (IE) wherein R^1 is a group of formula (i) and r is 1 then the reductive amination can be carried out using the same reaction conditions but using the corresponding homologous aldehyde of formula



instead of the corresponding 4-tetrahydropyranone. Alternatively, compounds of formula (IE) wherein R^1 is a group of formula (i) and r is 1 can be prepared via formation of an amide, followed by reduction of this amide bond to the corresponding amine as shown in Scheme 11E below:


88

**Scheme 11E**

The coupling reaction can be carried out using standard methods known in the art.


- 5 The reduction of the amide bond can also be carried by general methods known in the art for example using the same reduction conditions as those used in **Scheme 6**, such as in the presence of $\text{BH}_3\text{-Me}_2\text{S}$ (borane-dimethyl sulphide complex), in a suitable solvent such as THF.

- 10 Alternatively, compounds of formula (IE) wherein R^1 is a group of formula (i) wherein r is 0 can be prepared by a process illustrated in **Scheme 12E** for compounds wherein $-\text{Z}$ is hydrogen, s is 1, t is 2, each R^5 , R^6 , R^7 and R^8 are hydrogen and $-\text{X}-$ is $-\text{O}-$, (i.e. R^1 is 2-tetrahydrofuranyl). The compound of formula (IVE) can be alkylated with a compound of formula:



- 15 wherein L_4 is a suitable leaving group such as chloro, bromo, iodo, mesylate or tosylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding secondary amine which can be subsequently alkylated with a compound of formula $\text{Ar}_1\text{CH}_2\text{L}_1$ wherein L_1 is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as
- 20 potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding


intermediate of formula (VIII)_f. It will be appreciated that as mentioned above the same reaction can be carried out using Ar₁-CR³R⁴-L₁ wherein R³ and R⁴ are C₁-C₂ alkyl.



Scheme 12E

The tetrahydrofuranyl intermediates can be prepared from the corresponding 3-hydroxytetrahydrofuran, wherein the hydroxy group is converted into the leaving group using standard methods.

Compounds of formula (IE) wherein R¹ is a group of formula (i) and -X- is -SO₂- can be prepared from the corresponding intermediates (VIII)_f wherein the thioether is oxidized to the corresponding sulfoxide as shown in Scheme 13E below:




Scheme 13E

15

Compounds of formula (IE) wherein R¹ is a group of formula (ii) can be prepared using the synthesis illustrated in Scheme 14E for compounds wherein R¹ is oxabicyclo[3,2,1]octan-3-yl. The compound of formula (IVE) can be alkylated via reductive alkylation using standard methods, as those mentioned above with the ketone Ar₁-CO-R³. For example compound of formula (IVE) can be alkylated with oxabicyclo[3,2,1]octan-3-one in the presence of a suitable borane, such as sodium

20


borohydride or $\text{NaBH}(\text{OAc})_3$, optionally in the presence of an acid such as acetic acid, in the presence of a suitable solvent such as dichloroethane (DCE). Then, the secondary amine can be alkylated with a compound of formula $\text{Ar}_1\text{CH}_2\text{L}_1$ wherein L_1 is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding intermediate of formula (VIII E)_i. It will be appreciated that as mentioned above the same reaction can be carried out using $\text{Ar}_1\text{-CR}^3\text{R}^4\text{-L}_1$ wherein R^3 and R^4 are $\text{C}_1\text{-C}_2$ alkyl.



Scheme 14E


The oxabicyclo[3,2,1]octan-3-one intermediate is prepared according to the method described in A E Hill, G Greenwood and H M R Hoffmann JACS 1973, 95, 1338.

It will be appreciated that for compounds of formula (IE) wherein R^1 is a group of formula (i) and r is 1 then the reductive amination can be carried out using the same reaction conditions but using the corresponding homologous aldehyde of formula




instead of the corresponding oxabicyclo[3,2,1]octan-3-one.


Compounds of formula (IE) wherein Ar_1 is a substituted or unsubstituted pyridyl group can be prepared by a process illustrated in Scheme 15E for compounds wherein R^3 and R^4 are hydrogen and Ar_1 is 3-phenylpyrid-2-yl.

**Scheme 15E**

The compound of formula (IVE) can be alkylated via reductive alkylation using standard methods, as those mentioned above with the ketone $\text{Ar}_1\text{-CO-R}^3$. For example compound of formula (IVE) can be alkylated with an aldehyde of formula:




in the presence of a suitable borane, such as sodium borohydride or $\text{NaBH}(\text{OAc})_3$, optionally in the presence of an acid such as acetic acid, in the presence of a suitable solvent such as dichloroethane (DCE). Then, the secondary amine can be alkylated using the general methods described above for the incorporation of R^1 . The intermediate aldehyde can be prepared via reduction of readily available methyl 3-phenyl picolinate to the corresponding alcohol and subsequent oxidation to the aldehyde as shown in **Scheme 16E** below.

**Scheme 16E**

The reduction step can be carried out in the presence of a suitable reducing agent such as lithium borohydride in a suitable solvent such as tetrahydrofuran. The oxidation to the aldehyde can be carried out under Swern conditions such as oxalyl chloride and DMSO in DCM.


- 5 Compounds of formula (IE) wherein Ar_1 is a substituted or unsubstituted phenyl group can be prepared by a process illustrated in **Scheme 17E** for compounds wherein R^3 and R^4 are hydrogen and Ar_1 is 2-(3-pyridyl)phenyl.



10


Scheme 17E

The compound of formula (IV E) can be alkylated via reductive alkylation using standard methods, as those mentioned above with the ketone $\text{Ar}_1\text{-CO-R}^3$. For example compound of formula (IV E) can be alkylated with an aldehyde of formula:




- 15 in the presence of a suitable borane, such as sodium borohydride or NaBH(OAc)_3 , optionally in the presence of an acid such as acetic acid, in the presence of a suitable solvent such as dichloroethane (DCE). Then, the secondary amine can be alkylated using the general methods described above for the incorporation of R^1 . The intermediate aldehyde can be prepared from the commercially available 2-formyl phenyl boronic acid
- 20 via palladium coupling in the presence of 3-bromopyridine, a suitable palladium catalyst such as $\text{Pd(PPh}_3)_4$ and a suitable base such as potassium carbonate in a suitable solvent such as acetonitrile, as shown in **Scheme 18E** below.

93

**Scheme 18E**


Compounds of formula (IE) wherein Ar₁ is a phenyl group substituted with a 1-pyrazole group can be prepared by a process illustrated in **Scheme 19E**.

**Scheme 19E**


The pyrazole group can be incorporated by reacting a compound of formula (VIIIIE)_{m'}, wherein L₅ is a suitable leaving group such as bromo, chloro or iodo, with pyrazole in the presence of a suitable base such as potassium carbonate and a catalytic amount of copper iodide in a suitable solvent such as for example DMF. The compound of formula (VIIIIE)_{m'} can be prepared by any of the methods mentioned above for compounds wherein Ar₁ is a phenyl group substituted with a halogen atom such as chloro, bromo or iodo.

It will be appreciated that any of the intermediates (VIIIIE), (VIIIIE)_{a-m} are then deprotected using suitable deprotecting conditions such as those discussed in Greene, to give the corresponding compounds of formula (IE). For example if the protecting group is a boc group, the deprotection reaction can be carried out in trifluoroacetic acid in a suitable solvent such as DCM. Alternatively the reaction can be carried out in ethanolic hydrochloric acid.

94

**Scheme 20E**

Compounds of formula (IE) wherein R³ and R⁴ are both hydrogen may also be prepared by solid phase synthesis by the route shown below in **Scheme 21E** below.

**Scheme 21E**

The sequence is preferably performed on a polystyrene resin. The process may be run in a combinatorial fashion such that all possible compounds from sets of precursors Ar₁CHO and R⁹CHO may be prepared, wherein R⁹ is chosen such that R⁹-CH₂ = R¹, and R¹ and Ar₁ have the values defined above for formula (IE). The sequence is performed without characterisation of the resin-bound intermediates. In step (i) 3-trifluoroacetamido-pyrrolidine is bound to a solid support by reaction with 4-nitrophenyl carbonate activated polystyrene resin in the presence of a base, such as N,N-diisopropylethylamine, in a solvent such as DMF. In step (ii), the trifluoroacetamido protecting group is cleaved by hydrolysis with a base such as aqueous lithium hydroxide. In step (iii) the primary amine is then condensed with a substituted benzaldehyde in the presence of a dehydrating agent, such as trimethylorthoformate, to form the intermediate imine. In step (iv) the imine is

reduced with a borane reducing agent, such as sodium cyanoborohydride, in a solvent such as DMF, containing acetic acid. In step (v) the resultant secondary amine is then reductively alkylated with an aldehyde in the presence of a reducing agent such as sodium triacetoxyborohydride in a solvent, such as DMF. In step (vi) the desired product is
5 finally cleaved from the resin with acid, such as aqueous trifluoroacetic acid.


Preparation of Compounds of Formula (IF)

Compounds of formula (IF) may be prepared by conventional organic chemistry techniques and also by solid phase synthesis.

10 Compounds of formula (IF') can be prepared by the general methods illustrated below. It will be appreciated that the same methods can be used for compounds of formula (IF'') with the only difference that the nitrogen atom of the quinuclidines does not need to be protected as it is already a tertiary amine as it is explained in more detail below with reference to **Scheme 1F**.

15 Compounds of formula (IF') can be prepared via the 3-aminopiperidine intermediate of formula (IVF) as illustrated in **Scheme 1F** below:

96



Scheme 1 F


Commercially available 3-hydroxypiperidine of formula (IIIF) wherein R² is hydrogen, can be protected using a suitable nitrogen-protecting group such as those described in T.W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1991, hereafter referred to as "Greene". For example 3-*R*-hydroxypiperidine (IIIF) can be protected with a tert-butoxycarbonyl group, (boc). The protection reaction can be carried out for example using Boc anhydride in a suitable solvent such as for example tetrahydrofuran (THF) or dichloromethane (DCM) in the presence of a base such as triethylamine (TEA) or 4-(dimethylamino)pyridine (DMAP). It will be appreciated that for compounds of formula (IF) wherein R² is C₁-C₂ alkyl, the 3-

hydroxypiperidine of formula (IIIF) can be prepared from the readily available 3-pyrrolidinone via addition of the appropriate C₁-C₂ alkyl organometallic.

The hydroxy group of the N-protected-3-hydroxypiperidine can be converted into a suitable leaving group (L) such as for example chloride, bromide, iodide or mesylate. For example the N-protected-hydroxypiperidine can be converted to the mesylate in the presence of mesyl chloride and a suitable base such as triethylamine in a solvent such as DCM. Said mesylate is subsequently displaced with the corresponding azide in a suitable solvent such as dimethylformamide (DMF) or dimethylsulphoxide (DMSO). This azide intermediate can be converted to the corresponding N-protected-aminopiperidine of formula (IV) via hydrogenation in the presence of a suitable catalyst such as Palladium on charcoal and in a suitable solvent such as methanol or ethanol.

For compounds of formula (IF) wherein R⁴ is H, intermediate (IVF) can be alkylated via reductive alkylation with a ketone of formula R³-CO-Ar₁ wherein R³ and Ar₁ have the values for formula (IF) above. The reductive alkylation can be carried out for example as a hydrogenation reaction in the presence of a suitable catalyst such as Palladium on charcoal and a suitable solvent such as for example ethanol. Alternatively, said reductive alkylation can be carried out in the presence of a suitable borane such as sodium triacetoxyborohydride, NaBH(OAc)₃ and optionally in the presence of a suitable acid such as acetic acid, in a suitable solvent such as for example dichloroethane (DCE).


Alternatively, intermediate of formula (VF) wherein R⁴ is H can be prepared as shown in **Scheme 2F** below by reductive alkylation of readily available 3-aminopiperidine of formula (VIF) wherein R² has the values defined for formula (IF) above, followed by the protection of the nitrogen in the piperidine ring using a suitable protecting group such as those defined in Greene.



Scheme 2F

For example the reductive alkylation can be carried out in the presence of a ketone of formula $\text{Ar}_1\text{-CO-R}^3$ wherein Ar_1 and R^3 have the values defined for formula (IF) above. Initial condensation of the amino piperidine with the ketone is undertaken in the presence of a suitable acid such as p-toluenesulphonic acid, in a suitable solvent such as toluene. The resultant imino piperidine intermediate can then be protected with for example a boc group. The reaction can be carried out in the presence of boc anhydride and a suitable base such as DMAP, in a suitable solvent such as DCM. Said imine is reduced via hydrogenation in the presence of a suitable catalyst such as palladium on charcoal, in a suitable solvent such as ethanol to give the corresponding amine of formula (VF).

Intermediate of formula (VF) can be converted to compounds of formula (VIIF) via reductive alkylation with an aldehyde of formula $\text{R}^9\text{-CHO}$, wherein R^9 is chosen such that $\text{R}^9\text{-CH}_2 = \text{R}^1$ and R^1 has the values defined for formula (IF) above. The reductive alkylation can be carried out using standard methods, for instance as those mentioned above with the ketone $\text{Ar}_1\text{-CO-R}^3$.




Scheme 3F

For example a compound of formula (VF) can be alkylated with $\text{R}^9\text{-CHO}$ in the presence of a suitable borane, such as NaBH(OAc)_3 , optionally in the presence of an acid such as acetic acid, in the presence of a suitable solvent such as dichloroethane (DCE).


For compounds of formula (IF) wherein R^3 and R^4 are hydrogen the alkylation of intermediate (VF) can be carried out with a compound of formula $\text{Ar}_1\text{CH}_2\text{L}_1$ wherein L_1 is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding intermediate of formula (VIIF)_a. It will be appreciated that the same reaction can be carried out using $\text{Ar}_1\text{-CR}^3\text{R}^4\text{-L}_1$ wherein R^3 and R^4 are $\text{C}_1\text{-C}_2$ alkyl.

99



Scheme 4F


Compounds of formula (IF) wherein R^1 is $-\text{CH}_2\text{-COO}-(\text{C}_1\text{-C}_2 \text{ alkyl})$ can be prepared by reacting intermediate (VF) with a compound of formula $\text{L}_2\text{-CH}_2\text{-COO}-(\text{C}_1\text{-C}_2 \text{ alkyl})$ wherein L_2 is a suitable leaving group such as for example bromo, chloro or iodo. Said reaction can be carried out in the presence of a suitable base such as sodium hydride, in a suitable solvent such as dimethylformamide.




Scheme 5F

Compounds of formula (IF) wherein R^1 is $-(\text{CH}_2)_m\text{-CF}_3$ can be prepared by reacting intermediate (VF) with a compound of formula $\text{HOOC}-(\text{CH}_2)_{(m-1)}\text{-CF}_3$. The acid may be activated as its anhydride or acyl chloride, and is reacted in the presence of a suitable base such as triethylamine and a catalytic amount of DMAP, in a suitable solvent such as DCM. The resulting amide can be reduced to the amine of formula (VIII F)_c in the presence of a suitable borane. For example, for compounds wherein m is 1, the reduction can be carried out in the presence of $\text{BH}_3\text{-Me}_2\text{S}$ borane-dimethyl sulphide complex, in a suitable solvent such as THF.


100

**Scheme 6F**

Compounds of formula (IF) wherein R¹ is -(C₁-C₆ alkylene)-OH can be prepared by reacting intermediate (VF) with an epoxide. For example for compounds wherein R¹ is -CH₂-C(CH₃)₂-OH, the intermediate of formula (VF) is reacted with 2,2-dimethyloxirane, in a suitable solvent such as aqueous ethanol.


**Scheme 7F**

Alternatively compounds of formula (IF) wherein R¹ is -(C₁-C₆alkylene)-OH can be prepared by reacting intermediate (VF) with an ω-haloalkanoate, such as methylbromoacetate, in the presence of a base such a sodium hydrogen carbonate in a solvent such as acetonitrile. The intermediate ester is then reacted with 2 equivalents of methyl magnesium bromide in THF to yield the tertiary alcohol(VIII F)_d:

**Scheme 8F**

It will be appreciated that the **Scheme 8F** above applies to alkylene chains longer than -CH₂-.


Compounds of formula (IF) wherein R¹ is -C₂-C₆ alkenyl, -(CH₂)_n-S-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-O-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-O-(C₃-C₆ cycloalkyl), -(C₁-C₅ alkylene)-SO₂-(C₁-C₃ alkyl), -(C₁-C₅ alkylene)-OCF₃, or -(C₁-C₅ alkylene)-CN, can be prepared via alkylation of intermediate (VF) with a compound of formula L₂-C₂-C₆ alkenyl, L₂-(CH₂)_n-S-(C₁-C₃ alkyl), L₂-(C₁-C₅ alkylene)-O-(C₁-C₃ alkyl), L₂-(C₁-C₅ alkylene)-O-(C₃-C₆ cycloalkyl), L₂-(C₁-C₅ alkylene)-SO₂-(C₁-C₃ alkyl), L₂-(C₁-C₅ alkylene)-OCF₃, or L₂-(C₁-C₅ alkylene)-CN respectively, wherein L₂ is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding intermediate of formula (VIIF)_c.



Scheme 9F

Compounds of formula (IF) wherein R¹ is a group of formula (i) can be prepared using the synthesis illustrated in **Scheme 10F** for compounds wherein R¹ is 4-tetrahydropyranyl. The compound of formula (IVF) can be alkylated via reductive alkylation using standard methods, as those mentioned above with the ketone Ar₁-CO-R³. For example a compound of formula (IVF) can be alkylated with 4-tetrahydropyranone in the presence of a suitable borane, such as sodium borohydride or NaBH(OAc)₃, optionally in the presence of an acid such as acetic acid, in the presence of a suitable solvent such as dichloroethane (DCE). Then, the secondary amine can be alkylated with a compound of formula Ar₁CH₂L₁ wherein L₁ is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding intermediate of


formula (VIII F)_f. It will be appreciated that as mentioned above the same reaction can be carried out using Ar₁-CR³R⁴-L₁ wherein R³ and R⁴ are C₁-C₂ alkyl.



5


Scheme 10F

It will be appreciated that for compounds of formula (IF) wherein R¹ is a group of formula (i) and r is 1 then the reductive amination can be carried out using the same reaction conditions but using the corresponding homologous aldehyde of formula



- 10 instead of the corresponding 4-tetrahydropyranone. Alternatively, compounds of formula (IF) wherein R¹ is a group of formula (i) and r is 1 can be prepared via formation of an amide, followed by reduction of this amide bond to the corresponding amine as shown in **Scheme 11F** below:


103

**Scheme 11F**

The coupling reaction can be carried out using standard methods known in the art.


- 5 The reduction of the amide bond can also be carried out by general methods known in the art for example using the same reduction conditions as those used in **Scheme 6F**, such as in the presence of $\text{BH}_3\text{-Me}_2\text{S}$ (borane-dimethyl sulphide complex), in a suitable solvent such as THF.

- 10 Alternatively, compounds of formula (IF) wherein R^1 is a group of formula (i) wherein r is 0 can be prepared by a process illustrated in **Scheme 12F** for compounds wherein $-\text{Z}$ is hydrogen, s is 1, t is 2, each R^5 , R^6 , R^7 and R^8 are hydrogen and $-\text{X}-$ is $-\text{O}-$, (i.e. R^1 is tetrahydrofuran-3-yl). The compound of formula (IVF) can be alkylated with a compound of formula:



- 15 wherein L_4 is a suitable leaving group such as chloro, bromo, iodo, mesylate or tosylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding secondary amine which can be subsequently alkylated with a compound of formula $\text{Ar}_1\text{CH}_2\text{L}_1$ wherein L_1 is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as
- 20 potassium carbonate and a suitable solvent such as acetonitrile, to give the corresponding


intermediate of formula (VIIF)_f. It will be appreciated that as mentioned above the same reaction can be carried out using Ar₁-CR³R⁴-L₁ wherein R³ and R⁴ are C₁-C₂ alkyl.



Scheme 12F

The tetrahydrofuranyl intermediates can be prepared from the corresponding 3-hydroxytetrahydrofuran, wherein the hydroxy group is converted into the leaving group using standard methods.

Compounds of formula (IF) wherein R¹ is a group of formula (i) and -X- is -SO₂- can be prepared from the corresponding intermediates (VIIF)_f wherein the thioether is oxidized to the corresponding sulfoxide as shown in Scheme 13F below:




Scheme 13F

Compounds of formula (IF) wherein R¹ is a group of formula (ii) can be prepared using the synthesis illustrated in Scheme 14F for compounds wherein R¹ is oxabicyclo[3,2,1]octan-3-yl. The compound of formula (IVF) can be alkylated via reductive alkylation using standard methods, as those mentioned above with the ketone Ar₁-CO-R³. For example compound of formula (IVF) can be alkylated with oxabicyclo[3,2,1]octan-3-one in the presence of a suitable borane, such as sodium borohydride or NaBH(OAc)₃, optionally in the presence of an acid such as acetic acid, in

the presence of a suitable solvent such as dichloroethane (DCE). Then, the secondary amine can be alkylated with a compound of formula $\text{Ar}_1\text{CH}_2\text{L}_1$ wherein L_1 is a suitable leaving group such as chloro, bromo, iodo or mesylate, in the presence of a suitable base such as potassium carbonate and a suitable solvent such as acetonitrile, to give the

5 corresponding intermediate of formula (VIIF)_j. It will be appreciated that as mentioned above the same reaction can be carried out using $\text{Ar}_1\text{-CR}^3\text{R}^4\text{-L}_1$ wherein R^3 and R^4 are C₁-C₂ alkyl.




Scheme 14F

10

15

The oxabicyclo[3,2,1]octan-3-one intermediate is prepared according to the method described in A E Hill, G Greenwood and H M R Hoffmann JACS 1973, 95, 1338. It will be appreciated that for compounds of formula (IF) wherein R^1 is a group of formula (i) and r is 1 then the reductive amination can be carried out using the same


reaction conditions but using the corresponding homologous aldehyde of formula




instead of the corresponding oxabicyclo[3,2,1]octan-3-one.

Compounds of formula (IF) wherein Ar_1 is a substituted or unsubstituted pyridyl group can be prepared by a process illustrated in **Scheme 15F** for compounds wherein R^3 and R^4 are hydrogen and Ar_1 is 3-phenylpyrid-2-yl.


20

**Scheme 15F**

The compound of formula (IVF) can be alkylated via reductive alkylation using standard methods, as those mentioned above with the ketone $\text{Ar}_1\text{-CO-R}^3$. For example compound of formula (IVF) can be alkylated with an aldehyde of formula:




in the presence of a suitable borane, such as sodium borohydride or NaBH(OAc)_3 , optionally in the presence of an acid such as acetic acid, in the presence of a suitable solvent such as dichloroethane (DCE). Then, the secondary amine can be alkylated using the general methods described above for the incorporation of R^1 . The intermediate aldehyde can be prepared via reduction of readily available methyl 3-phenyl picolinate to the corresponding alcohol and subsequent oxidation to the aldehyde as shown in **Scheme 16F** below.

**Scheme 16F**

The reduction step can be carried out in the presence of a suitable reducing agent such as lithium borohydride in a suitable solvent such as tetrahydrofuran. The oxidation to the aldehyde can be carried out under Swern conditions such as oxalyl chloride and DMSO in DCM.


- 5 Compounds of formula (IF) wherein Ar_1 is a substituted or unsubstituted phenyl group can be prepared by a process illustrated in **Scheme 17F** for compounds wherein R^3 and R^4 are hydrogen and Ar_1 is 2-(3-pyridyl)phenyl.



10


Scheme 17F

The compound of formula (IVF) can be alkylated via reductive alkylation using standard methods, as those mentioned above with the ketone $\text{Ar}_1\text{-CO-R}^3$. For example compound of formula (IVF) can be alkylated with an aldehyde of formula:




- 15 in the presence of a suitable borane, such as sodium borohydride or NaBH(OAc)_3 , optionally in the presence of an acid such as acetic acid, in the presence of a suitable solvent such as dichloroethane (DCE). Then, the secondary amine can be alkylated using the general methods described above for the incorporation of R^1 . The intermediate aldehyde can be prepared from the commercially available 2-formyl phenyl boronic acid
- 20 via palladium coupling in the presence of 3-bromopyridine, a suitable palladium catalyst such as $\text{Pd(PPh}_3)_4$ and a suitable base such as potassium carbonate in a suitable solvent such as acetonitrile, as shown in **Scheme 18F** below.

108

**Scheme 18F**


Compounds of formula (IF) wherein Ar₁ is a phenyl group substituted with a 1-pyrazole group can be prepared by a process illustrated in **Scheme 19F**.

**Scheme 19F**


The pyrazole group can be incorporated by reacting a compound of formula (VIIIIF)_{m'}, wherein L₅ is a suitable leaving group such as bromo, chloro or iodo, with pyrazole in the presence of a suitable base such as potassium carbonate and a catalytic amount of copper iodide in a suitable solvent such as for example DMF. The compound of formula (VIIIIF)_m can be prepared by any of the methods mentioned above for compounds wherein Ar₁ is a phenyl group substituted with a halogen atom such as chloro, bromo or iodo.

It will be appreciated that any of the intermediates (VIIIIF), (VIIIIF)_{a-m} are then deprotected using suitable deprotecting conditions such as those discussed in Greene, to give the corresponding compounds of formula (IF). For example if the protecting group is a boc group, the deprotection reaction can be carried out in trifluoroacetic acid in a suitable solvent such as DCM. Alternatively the reaction can be carried out in ethanolic hydrochloric acid.

109

**Scheme 20F**

Compounds of formula (IF) wherein R³ and R⁴ are both hydrogen may also be prepared by solid phase synthesis by the route shown below as **Scheme 21F**.


**Scheme 21F**

The sequence is preferably performed on a polystyrene resin. The process may be run in a combinatorial fashion such that all possible compounds from sets of precursors Ar₁CHO and R⁹CHO may be prepared, wherein R⁹ is chosen such that R⁹-CH₂ = R¹, and R¹ and Ar₁ have the values defined above for formula (IF). The sequence is performed without characterisation of the resin-bound intermediates. In step (i) 3-trifluoroacetamido-piperidine is bound to a solid support by reaction with 4-nitrophenyl carbonate activated polystyrene resin in the presence of a base, such as N,N-diisopropylethylamine, in a solvent such as DMF. In step (ii), the trifluoroacetamido protecting group is cleaved by hydrolysis with a base such as aqueous lithium hydroxide. In step (iii) the primary amine is then condensed with a substituted benzaldehyde in the presence of a dehydrating agent, such as trimethylorthoformate, to form the intermediate imine. In step (iv) the imine is

- reduced with a borane reducing agent, such as sodium cyanoborohydride, in a solvent such as DMF, containing acetic acid. In step (v) the resultant secondary amine is then reductively alkylated with an aldehyde in the presence of a reducing agent such as sodium triacetoxyborohydride in a solvent, such as DMF. In step (vi) the desired product is
- 5 finally cleaved from the resin with acid, such as aqueous trifluoroacetic acid.

Preparation of Compounds of Formula (IG)

- Compounds of formula (IG) may be prepared by conventional organic chemistry techniques from N-protected-2-cyanomorpholines as outlined in Error! Reference source not found.G below, wherein R and R² have the values defined for formula (IG) above and P is a suitable nitrogen protecting group such as those described in T.W. Greene,
- 10 "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1991, hereafter referred to as "Greene". For example a suitable nitrogen protecting group is a benzyl group:




15

Scheme 1G

The phenyl ketone (IIIG) can be obtained by reaction of N-protected-2-cyanomorpholine with a Grignard reagent, followed by acid hydrolysis to give the racemic phenyl ketone which may be separated on chiral HPLC.

- 20 Compounds of formula (IG) can be prepared from the N-protected morpholine ketone intermediate of formula (IIIG), as illustrated in Error! Reference source not found.G below:

111

**Scheme 2G**

The ketone is stereoselectively reduced to the corresponding (2S) or (2R) alcohol of formula (IVG) or (IVG)_a using standard methods known in the art. For example it can be reduced in the presence of [(-)-B-chlorodiisopinocampheylborane] in a suitable solvent such as tetrahydrofuran (THF) to provide the (2S) alcohol.


The resulting alcohol is then transformed into a suitable leaving group L. Suitable leaving groups include halo groups, such as bromo, chloro or iodo and sulfonate groups, such as mesylate. When L is a halo group, the alcohol used will be the (2S) enantiomer (IVG) and it will be reacted with inversion of stereochemistry. For example, when L is bromo, the bromination reaction can be carried out in the presence of a brominating agent

such as triphenylphosphine dibromide, in a suitable solvent such as chloroform. When L is a mesylate group, the alcohol used will be the (2R) enantiomer (IVG)_a and it will be reacted with retention of stereochemistry in the presence of mesylate chloride and a suitable base.

5 The resulting intermediate of formula (VG) can then be converted into the corresponding methylethanethioate of formula (VIG) via displacement of the leaving group with a suitable thiolacetate salt such as potassium thiolacetate in the presence of a suitable solvent such as a mixture of dimethylformamide (DMF) and tetrahydrofuran (THF).

10 The methanethiol intermediate of formula (VIIG) can be prepared via reaction of the methylethanethioate (VIG) with a suitable thiomethoxide such as sodium thiomethoxide in the presence of a suitable solvent such as methanol (one can use a variety of bases but thiomethoxide is preferred because it also acts as a reducing agent and prevents oxidation of thiol hence inhibiting dimerisation; Ref: O.B.Wallace &
15 D.M.Springer, *Tetrahedron Letters*, 1998, 39 (18), pp2693-2694).

The pyridyl portion of the molecule is incorporated via general methods known in the art. A particularly useful method is the reaction of the methanethiol (VIIG) with a compound of the formula




20 wherein R¹ has the values defined above and L₁ is a suitable leaving group such as fluoro, bromo, chloro, iodo or mesylate, in the presence of suitable base such as sodium hydride, cesium fluoride or sodium methoxide, in a suitable solvent such as DMF.

Compounds of formula (IG) wherein -X- is -O- can be prepared in an analogous fashion by reaction of the (2S) alcohol of formula (IVG) with a compound of formula
25 (VIII G) above.

The final step for the preparation of compounds of formula (IG) comprises deprotection of the morpholine ring. Conditions for the deprotection depend on the protecting group chosen. Suitable deprotecting conditions can be found in Greene. For

example when the nitrogen protecting group is a benzyl group, the deprotection reaction can be carried out in the presence of polymer supported diisopropylamine (PS-DIEA) and 1-chloroethyl chloroformate (ACE-Cl) in a suitable solvent such as dichloromethane, followed by reaction with methanol to give compounds of formula (IG).


- 5 Compounds of formula (IG) can alternatively be prepared by the derivatisation of a suitable substituent in the pyridyl ring to give the desired substituent R^1 as shown in **Scheme 3G** below. For example compounds of formula (IG) wherein $-R^1$ is $-CF_3$ can be prepared via reaction of the intermediate (IXG)' wherein L_2 is introduced into the molecule in place of R^1 in formula (VIIG) as shown in Error! Reference source not found.G above. The group L_2 is a suitable leaving group such as for example iodo, bromo, chloro or fluoro. The leaving group is converted into a trifluoromethyl group via reaction in the presence of copper iodide, a suitable base such as for example potassium fluoride, and a suitable source of a trifluoromethyl group such as for example (trifluoromethyl)trimethylsilane, in a suitable solvent such as for example a mixture of
- 10 DMF and N-methyl-pyrrolidinone (NMP). The resulting compound of formula (XG) is deprotected using the methodology described above.
- 15



Scheme 3G

- Compounds of formula (IG) wherein $-X-$ is $-S-$ can alternatively be prepared directly from the intermediate methylethanethioate of formula (VIG) as illustrated in Error! Reference source not found.G below.
- 20


114

**Scheme 4G**

The reaction can be carried out via general methods known in the art. For example, the intermediate (VIG) can be reacted with a compound of formula (VIII), wherein R¹ and L₁ have the values defined above, in the presence of a suitable base such as sodium methoxide, in a suitable solvent such as for example DMF.

The resulting compound of formula (IXG) wherein -X- is -S- is then deprotected using the methods described above for Error! Reference source not found.G to give a compound of formula (IG) wherein -X- is -S-. This method is particularly useful when L₁ and R¹ are halogen groups such as for example fluoro and bromo respectively. Alternatively, the reaction can be carried out in the presence of a suitable base such as sodium hydroxide in a suitable solvent such as a mixture of ethanol and water. This method is particularly useful when L₁ is a halogen group and -R¹ is -CN or -CONR³R⁴, wherein R³ and R⁴ have the values defined for formula (IG) above.

Compounds of formula (IG) wherein -X- is -S- can also be prepared via an alternative method using the intermediate of formula (VG) as illustrated below in Error! Reference source not found.G.


**Scheme 5G**

The leaving group of intermediate (VG) is displaced with a suitable thiol of formula (XIG) wherein R^1 has the values defined for formula (IG) above, in the presence of a suitable base such as potassium carbonate, in a suitable solvent such as DMF. The resulting intermediate of formula (IXG) wherein -X- is -S- is then deprotected as described in Error! Reference source not found.G above.

The intermediate of formula (VIII G) above (including analogs wherein L_2 is introduced in place of R^1) often commercially available. This is the case for intermediates wherein L_1 is a halogen group and R^1 (or L_2) has the values selected from H, methyl, halo, cyano, trifluoromethyl, NH_2 , CO_2H , $CONH_2$, SO_2H , SO_2NHCH_3 , $NCOCCl_3$ and NSO_2Ph .

Intermediates of formula (VIII G) wherein R^1 is a group of formula (i) can readily be prepared via methods known in the art. We illustrate below 3 methods for the preparation of compounds of formula (VIII G) wherein R^1 is a group of formula (i) and -Z- has the value of a bond (Error! Reference source not found.G), $-CH_2-$ (Error! Reference source not found.G) or $-O-$ (Error! Reference source not found.G). It will be appreciated that these methods are only illustrative as there are many other alternative methods known in the art which can be used.

As mentioned above, intermediates of formula (VIII G) wherein R^1 is a group of formula (i) and -Z- is a bond can be prepared via palladium coupling as illustrated in Error! Reference source not found.G below.




Scheme 6G

The reaction is carried out via reaction of readily available pyridines of formula (XIIG) wherein L_1 has the values mentioned above and L_3 is a suitable leaving group such as for example a halogen group such as bromo or chloro, with the corresponding phenylboronic acid of formula (XIIIG), in the presence of a suitable palladium catalyst

such as for example palladium acetate, a suitable ligand such as triphenylphosphine, in a suitable solvent such as acetonitrile. Alternative palladium catalysts are known in the art, for example bis(benzonitrile)palladium(II)dichloride can be used in the presence of a suitable ligand such as for example bis(diphenylphosphine)butane and a suitable base
 5 such as sodium carbonate in a suitable solvent such as for example ethanol, to give good yields of intermediate of formula (VIII G) wherein R^1 is a group of formula (i) and -Z- is a bond.

Intermediates of formula (VIII G) wherein R^1 is a group of formula (i) and -Z- is -CH₂- can be prepared by the method illustrated in Error! Reference source not found. G
 10 below.




Scheme 7G

Readily available pyridine compounds of formula (XIV G) wherein L_1 has the values mentioned above (preferably fluoro) are reacted with suitable benzaldehydes of formula (XV G), wherein R^5 has the value defined for formula (IG) above, in the presence of a suitable base such as for example n-butyllithium or lithium diisopropylamide, in a suitable solvent such as THF, to give the alcohol of formula (XVIG). Said alcohol is then
 20 reduced to give the corresponding benzyl derivative (VIII G) wherein R^1 is a group of formula (i) and -Z- is -CH₂- via hydrogenation, in the presence of a suitable catalyst such as for example palladium on charcoal, in a suitable solvent such as for example ethanol.


Intermediates of formula (VIII G) wherein R^1 is a group of formula (i) and -Z- is -O- can be prepared by the method illustrated below in Error! Reference source not
 25 found. G.

117

**Scheme 8G**


Readily available pyridinols of formula (XVIIIG), wherein L_1 has the values mentioned above react with phenylboronic acids of formula (XIIIIG) in the presence of copper(II)acetate, powdered 4Å molecular sieves, and a suitable base such as triethylamine, in a suitable solvent such as for example dichloromethane to give intermediates of formula (VIIIIG) wherein R^1 is a group of formula (i) and $-Z-$ is $-O-$.

Compounds of formula (IG) wherein $-X-$ is $-O-$ may also be prepared by conventional chemistry techniques from the (2R) alcohol (IVG)_a using standard methods known in the art. For example as shown in **Scheme 9G** by reaction of said alcohol with a pyridine of the formula (XVIIIIG) or the ketone tautomer of this pyridine wherein R^1 has the values defined for formula (IG) above, in the presence of a suitable phosphine such as triphenyl phosphine and diethyl azodicarboxylate, using an appropriate solvent such as THF, dimethoxyethane, (DME), or chloroform ($CHCl_3$), as described by D.L. Comins and G. Jianhua, in *Tetrahedron Letters*, **1994**, 35 (18), pp2819-2822. This reaction is usually carried out with inversion of the stereocentre to (2S)

**Scheme 9G**

As previously mentioned, compounds of formula (IG) wherein $-X-$ is $-O-$ may alternatively be prepared by the reaction of the (2S) alcohol (IVG) with a pyridine of the

formula (VIII G), where L_1 is preferably chloro and R^1 has the values defined for formula (IG) above, using a suitable base such as potassium hydroxide, in a suitable solvent such as benzene or toluene, in the presence of a suitable phase transfer catalyst such as 18-Crown-6 as described by A.J.S. Duggan *et al*, in *Synthesis*, **1980**, 7, p573.



Scheme 10G

Compounds of formula (IG) wherein $-X-$ is $-O-$ may alternatively be prepared by the reaction of intermediate (VG) wherein L is Br with a pyridine of the formula (VIII G) wherein $-L_1$ is $-OAg$ and R^1 has the values defined for formula (IG) above, in a non-polar solvent such as benzene, as described by U. Schollkopf *et al*, in *Liebigs Ann. Chem.* **1972**, 765, pp153-170 and G.C. Hopkins *et al*, in *J. Org. Chem.* **1967**, 32, pp4040.

15 It will be appreciated that compounds of Formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) above possess one or more asymmetric carbon atoms, and that in the present invention specific individual stereoisomers are preferred. In the present specification, where a structural formula does not specify the stereochemistry at one or more chiral centres, it encompasses all possible stereoisomers and all possible mixtures of stereoisomers (including, but not limited to, racemic mixtures), which can result from stereoisomerism at each of the one or more chiral centers.

20

The following examples illustrate compounds of of Formula (IA) above and methods for their preparation.

Example 1A: N-(2-methylpropyl)-N-[(2-fluorophenyl)methyl]piperidin-4-amine fumarate

To a dry boiling tube (50 ml), under nitrogen, was added tert-butyl-4-(2-methyl-
5 propylamino)-piperidine-1-carboxylate (0.200g, 0.780 mmol), 2-fluorobenzaldehyde (0.087 ml, 0.102g, 0.819 mmol), and titanium isopropoxide (0.268 ml, 0.937 mmol) to give a yellow/orange solution. This was heated to 90°C for 2 hours. Solution cooled, and ethanol (5 ml) added. Sodium borohydride (0.030g, 0.780 mmol) was then added and allowed to stir for 2 days. Further sodium borohydride (0.300g, 7.80 mmol) was added,
10 and after 6 hours, this was diluted with methanol (10 ml) with stirring for 20 hours. This was concentrated in vacuo, dissolved in dichloromethane (5 ml), and acetic anhydride (0.371 ml, 39.00 mmol) added with stirring for 30 minutes. Solution was diluted with methanol (10 ml), and passed through an SCX-2 column to give an oil (0.150g, 0.412 mmol).

15 The resultant oil was dissolved in dichloromethane (5 ml), and trifluoroacetic acid (2 ml) added. Reaction was monitored by thin layer chromatography (100% ethyl acetate; reactant. r.f. 0.4, product r.f. 0.0). After 2 hours, reaction was concentrated in vacuo, azeotroped with dichloromethane (c.a. 25 ml), taken up in methanol (c.a. 5 ml), and passed through an SCX-2 column. The resultant colourless oil was purified using reverse
20 phase chromatography, concentrated in vacuo, taken up in 5 M hydrochloric acid (10 ml), and heated to 90°C for 3 hours. This solution was freeze dried to give an oil (0.049g, 0.185 mmol). Resultant oil was passed through an SCX-2 column, dissolved in aqueous acetonitrile (c.a. 20 ml), and fumaric acid (0.0214g, 0.1850 mmol) added. After 5 minutes, this was freeze dried to give a white solid (0.070g, 0.185 mmol) as the title
25 compound. δ_H (300 MHz, MeOD) 7.47 (1H, t, Ar), 7.25 (1H, m, Ar), 7.13 (1H, t, Ar), 7.02 (1H, t, Ar), 6.70 (2H, s, fumarate), 3.21 (2H, s, NCH₂Ar), 3.45 (2H, d, CH), 2.95 (2H, t, CH), 2.82 (1H, t, CH), 2.29 (2H, d, NCH₂), 2.00 (2H, d, CH), 1.80 (2H, t, m), 1.68 (1H, t, CH), 0.85 (6H, d, CHMe₂). LCMS 12 minute gradient, Rt = 1.99 mins, (M⁺+1) = 265.2

Example 2A: N-(3,3-dimethylbutyl)-N-[(2-biphenyl)methyl]piperidin-4-amine fumarate

To a 100 ml round bottomed flask, under nitrogen, was added the 1,1-dimethylethyl 4-[(2-bromophenylmethyl)(3,3-dimethylbutyl)amino]piperidine-1-carboxylate (0.675 g, 1.49 mmole, 1.0eq.), phenylboronic acid (0.363 g, 2.98 mmole, 2.0 eq.), dichlorobis(triphenylphosphine)palladium(II) (0.104 g, 0.15 mmole, 0.1 eq.), sodium carbonate (0.158 g, 2.98 mmole, 2.0 eq.) and a 1:1 mixture of tetrahydrofuran : water (50 ml). The mixture was heated at 90°C for two hours. The reaction mixture was allowed to cool then poured into diethyl ether (100 ml). This organic mixture was washed with a solution of sodium hydroxide (2M, aqueous, 80 ml) then concentrated *in vacuo* to give a dark yellow oil (1.18 g). This oil was purified by automated flash chromatography using an ISCO Combiflash system (SiO₂ (120 g); 0-10% methanol (+5% 7M NH₃/MeOH) in dichloromethane gradient elution over 40 minutes) to give a yellow oil (0.683 g). This oil was further purified by automated flash chromatography using an ISCO Combiflash system (SiO₂ (120 g); ethyl acetate gradient elution over 40 minutes) to give 1,1-dimethylethyl 4-[(2-biphenyl)methyl](3,3-dimethylbutyl)amino]piperidine-1-carboxylate as a yellow oil (0.549 g, 82%). To a solution of this oil (0.549 g, 1.22 mmole, 1.0 eq.) in dichloromethane (10 ml) was added trifluoromethanesulfonic acid (TFA) (1.36 ml, 18.27 mmole, 15 eq.). The solution was stirred for one hour at room temperature. Solvent and TFA were removed *in vacuo*. The resulting oil was taken up in methanol and loaded onto an SCX-2 (10 g) column. The column was washed with methanol (50 ml). Basic material was then eluted using 2N ammonia in methanol (50 ml). Removal of solvent from the ammonia/methanol mixture under vacuum, gave a colourless oil (0.27 g). This oil was purified on the Biotage Parallel Flex Purification System (UV-guided HPLC) followed by SCX-2 treatment (to obtain the free base) to give a colourless oil (0.132 g). To a solution of this oil in methanol was added a solution of fumaric acid (0.044 g, 0.38 mmole, 1 eq) in methanol. The mixture was left to stir for a couple of minutes, ethyl acetate and cyclohexane were then added. The resulting precipitate was collected by filtration to give the title compound as a white solid (0.121 g, 17%). δ_H (300 MHz, MeOD) 7.50-7.47 (1H, m, ArH), 7.35-7.18 (7H, m, ArH), 7.10-7.07 (1H, m, ArH), 6.61 (3H, s, fumarate CH), 3.58 (2H, s, CH₂Ar), 3.25-3.24 (2H, m, NCH₂), 2.74 (2H, dt,

NCH₂), 2.67-2.57 (1H, m, NCH), 2.34-2.29 (2H, m, NCH₂), 1.65-1.45 (4H, m, CCH₂), 1.13-1.08 (2H, m, CH₂tBu), 0.70 (9H, s, CH₃); LCMS 12 min, Rt = 4.3 min, (M⁺+1) = 351.

5 **Example 3A: N-(2-ethylbutyl)-N-[(2-biphenyl)methyl]piperidin-4-amine fumarate**

As method previously described for Example 2A, using 1,1-dimethylethyl 4-[(2-bromophenylmethyl)(2-ethylbutyl)amino]piperidine-1-carboxylate. Isolation of the fumarate salt from methanol, diethyl ether, cyclohexane yielded the title compound as a white solid (0.238 g, 34%). δ_H (300 MHz, MeOD) 7.59-7.57 (1H, m, ArH), 7.45-7.27 (7H, m, ArH), 7.19-7.16 (1H, m, ArH), 6.69 (1.5H, s, fumarate CH), 3.62 (2H, s, CH₂Ar), 3.34-3.32 (2H, m, NCH₂), 2.79 (2H, dt, NCH₂), 2.66-2.57 (1H, m, NCH), 2.21 (2H, d, NCH₂), 1.64-1.50 (4H, m, CCH₂), 1.38-1.17 (5H, m, CH(CH₂Me)₂), 0.78 (6H, t, CH₃); LCMS 12 min, Rt = 5.1 min, (M⁺+1) = 351.

15 **Example 4A: N-(cyclohexylmethyl)-N-[(2-biphenyl)methyl]piperidin-4-amine fumarate**

(i) To a solution of cyclohexylmethylaniline (0.461 g, 4.08 mmole, 1.02 eq.) in 1,2-dichloroethane (10 ml) was added 1-Boc-4-piperidone (0.797 g ml, 4.00 mmole, 1.0 eq.). To this was added a solution of sodium triacetoxymethylborohydride (0.865 g, 4.08 mmole, 1.02 eq.) in dimethylformamide (2 ml). This mixture was left to stir under nitrogen, at room temperature, over the weekend. To the reaction mixture was then added water (10 ml) and the mixture stirred vigorously for several minutes. The chlorinated organic layer was then run through a hydrophobic frit then diluted with methanol (10 ml) and loaded onto an SCX-2 (10 g) column. The column was washed with methanol (50 ml) then basic material eluted with 2N ammonia in methanol. The ammonia/methanol solution was concentrated *in vacuo* to give a pale yellow oil (1.2 g). This was purified by automated flash chromatography using an ISCO Combiflash system (SiO₂ (40 g); 0-10% methanol in ethyl acetate gradient elution over 40 minutes) to give 1,1-dimethylethyl 4-[(cyclohexylmethyl)amino]piperidine-1-carboxylate as a colourless oil (0.98 g, 83%). δ_H (300 MHz, CDCl₃) 4.03-4.00 (2H, m, NCH₂), 2.83-2.75 (2H, m, NCH₂), 2.60-2.49 (1H,

m, NCH), 2.45 (2H, d, NCH₂), 1.18-0.83 (15H, m, CCH₂), 1.45 (9H, s, OC(CH₃)₃); LCMS 6 min, Rt = 2.7 min, (M⁺+1) = 297.

(ii) To a solution of 1,1-dimethylethyl 4-[(cyclohexylmethyl)amino]piperidine-1-carboxylate (0.245 g, 0.840 mmole, 1.0 eq.), 2-phenylbenzyl bromide (0.185 ml, 1.01 mmole, 1.2 eq.) in dry acetonitrile (5 ml) was added anhydrous potassium carbonate (0.19 g, 1.35 mmole, 1.6 eq.). The mixture was stirred overnight at room temperature. The reaction mixture was concentrated under vacuum to give a white solid. The white solid was taken up in dichloromethane (10 ml) and this washed with water (10 ml). The dichloromethane layer was passed through a hydrophobic frit then diluted with methanol (10 ml). This solution was loaded onto an SCX-2 (10 g) column. The column was washed with methanol (50 ml) then basic material was eluted using 2N ammonia in methanol (50 ml). Concentration of the ammonia/methanol solution under vacuum yielded a colourless oil (0.344 g, 90%). To a solution of this oil (0.344 g, 0.74 mmole, 1.0 eq.) in dichloromethane (10 ml) was added trifluoroacetic acid (TFA) (0.83 ml, 11.2 mmole, 15 eq.). The solution was stirred overnight at room temperature. Solvent and TFA were removed *in vacuo*. The resulting oil was taken up in methanol and loaded onto an SCX-2 (10 g) column. The column was washed with methanol (50 ml). Basic material was then eluted using 2N ammonia in methanol (50 ml). Removal of solvent from the ammonia/methanol mixture under vacuum, gave a colourless oil (0.298 g, 99%). The oil was taken up in methanol. To this solution was added a solution of fumaric acid (0.095 g, 0.08 mmole, 1 eq) in methanol followed by diethyl ether and cyclohexane. The resulting precipitate was collected by filtration to give the title compound as a white solid (0.302 g, 76 %). δ_H (300 MHz, MeOD) 7.58 (1H, d, ArH), 7.45-7.29 (7H, m, ArH), 7.18 (1H, d, ArH), 6.70 (2H, s, fumarate CH), 3.64 (2H, s, CH₂Ar), 3.33-3.32 (2H, m, NCH₂), 2.79 (2H, dt, NCH₂), 2.65-2.54 (1H, m, NCH), 2.17 (2H, d, NCH₂), 1.74-1.47 (9H, m, CCH₂), 1.28-1.11 (4H, m, CH, CCH₂), 0.78-0.67 (2H, m, CH₂); LCMS 12 min, Rt = 5.0 min, (M⁺+1) = 363.

Example 5A: N-(cyclopropylmethyl)-N-[(2-biphenyl)methyl]piperidin-4-amine fumarate

As method previously described for Example 4A, using 1,1-dimethylethyl 4-[(cyclopropylmethyl)amino]piperidine-1-carboxylate and 2-phenylbenzyl bromide. Isolation of the fumarate salt from methanol and diethyl ether yielded the title compound as a white solid (0.485 g, 74%). δ_H (300 MHz, MeOD) 7.68 (1H, dd, ArH), 7.47-7.29 (7H, m, ArH), 7.21 (1H, d, ArH), 6.72 (2H, s, fumarate CH), 3.76 (2H, s, CH₂Ar), 3.38-3.34 (2H, m, NCH₂), 2.92-2.82 (3H, m, NCH, NCH₂), 2.32 (2H, d, NCH₂), 1.79-1.57 (4H, m, CCH₂), 0.77-0.66 (1H, m, CH), 0.46-0.40 (2H, m, CH₂), 0.03-0.02 (2H, m, CH₂); LCMS 12 min, Rt = 3.5 min, (M⁺+1) = 321.

10 **Example 6A: N-(3-methylbutyl)-N-[(2-phenoxyphenyl)methyl]piperidin-4-amine difumarate**

(i) To 10% Pd/C (1.0 g, 10%wt), under nitrogen, was added a solution of the 1-Boc-4-piperidone (10.0 g, 50.1 mmole, 1.0 eq.) and isoamylamine (4.46 g, 51.2 mmole, 1.02 eq.) in ethanol (60 ml). This was hydrogenated overnight, at 60 psi using a Parr
15 hydrogenator. The catalyst was removed by filtration through Celite. Solvent was removed under vacuum to give 1,1-dimethylethyl 4-[(3-methylbutyl)amino]piperidine-1-carboxylate as a colourless, slightly cloudy, oil (13.59 g, 100%). δ_H (300 MHz, CDCl₃) 4.05-4.02 (2H, m, NCH₂), 2.82-2.75 (2H, m, NCH₂), 2.66-2.54 (3H, m, NCH, NCH₂), 1.86-1.82 (2H, m, CCH₂), 1.62 (1H, septet, CHMe₂), 1.45 (9H, s, OC(CH₃)₃), 1.41-1.17
20 (4H, m, CCH₂), 0.90 (6H, d, C(CH₃)₂); LCMS 6 min, Rt = 2.7 min, (M⁺+1) = 271.

(ii) To a solution of 1,1-dimethylethyl 4-[(3-methylbutyl)amino]piperidine-1-carboxylate in 1,2-dichloroethane (10 ml) was added 2-phenoxybenzaldehyde. To this was added a solution of sodium triacetoxymethylborohydride (3.0 eq.) in dimethylformamide (2 ml). This mixture was left to stir for 3 days under nitrogen, at room temperature. To the
25 reaction mixture was added water (10 ml) and the mixture stirred vigorously for several minutes. The chlorinated organic layer was run through a hydrophobic frit to remove water, diluted with methanol (10 ml) and loaded onto an SCX-2 (10 g) column. The column was washed with methanol (50 ml) then basic material eluted with 2N ammonia in methanol. The ammonia/methanol solution was concentrated *in vacuo* to give 1,1-dimethylethyl 4-[(2-phenoxyphenyl)methyl](3-methylbutyl)amino]piperidine-1-
30 carboxylate as a colourless oil. To a solution of this oil in dichloromethane (10 ml) was

added trifluoroacetic acid (TFA) (15 eq). The solution was stirred overnight at room temperature. Solvent and TFA were removed *in vacuo*. The resulting oil was taken up in methanol and loaded onto an SCX-2 (10 g) column. The column was washed with methanol (50 ml). Basic material was then eluted using 2M ammonia in methanol (50 ml). Removal of solvent from the ammonia/methanol mixture under vacuum, gave a colourless oil. The oil was taken up in methanol. To this solution was added a solution of fumaric acid (1 eq) in methanol. The mixture was left to stir for a couple of minutes, then ethyl acetate and cyclohexane were added. The resulting precipitate was collected by filtration to give the title compound as a white solid (0.264 g, 30%). δ_H (300 MHz, MeOD) 7.46 (1H, dd, ArH), 7.26-7.16 (3H, m, ArH), 7.10-7.04 (1H, m, ArH), 7.00-6.95 (1H, m, ArH), 6.86-6.79 (3H, m, ArH), 6.61 (4H, s, fumarate CH), 3.68 (2H, s, CH₂Ar), 3.33-3.28 (2H, m, NCH₂), 3.04-2.96 (3H, m, NCH, NCH₂), 2.56-2.51 (2H, m, NCH₂), 1.91-1.87 (2H, m, CCH₂), 1.76-1.62 (2H, m, CCH₂), 1.52-1.41 (1H, m, CH), 1.30-1.23 (2H, m, CH₂), 0.74 (6H, d, CH₃); LCMS 12 min, Rt = 4.2 min, (M⁺+1) = 353.

15

Example 7A: N-(3-methylbutyl)-N-[(2-biphenyl)methyl]piperidin-4-amine difumarate

As method previously described for Example 4A, using 1,1-dimethylethyl 4-[(3-methylbutyl)amino]piperidine-1-carboxylate and 2-phenylbenzyl bromide. Isolation of the fumarate salt from methanol and diethyl ether yielded the title compound as a white solid (0.239 g, 24%). δ_H (300 MHz, MeOD) 7.49 (1H, dd, ArH), 7.35-7.18 (7H, m, ArH), 7.10 (1H, dd, ArH), 6.61 (4H, s, fumarate CH), 3.62 (2H, s, CH₂Ar), 3.25 (2H, m, NCH₂), 2.78-2.59 (3H, m, NCH, NCH₂), 2.36-2.31 (2H, m, NCH₂), 1.64-1.45 (4H, m, CCH₂), 1.42-1.31 (1H, m, CH), 1.13-1.05 (2H, m, CH₂), 0.69 (6H, d, CH₃); LCMS 12 min, Rt = 4.1 min, (M⁺+1) = 337.

25

The following examples illustrate compounds of Formula (IB) above and methods for their preparation.

30

Synthesis of Intermediates.


Preparation of (4-Benzyl-morpholin-2-yl)-phenyl-methanone.

A 1600 L GL reactor under N₂ was successively loaded with 2-chloroacrylonitrile
5 (33.2 kg, 379 moles) and toluene (114 L) at 21°C. Then, N-benzylethanolamine (57 kg,
377 moles) was added and the reaction mixture was post-agitated at room temperature for
about 17 h. Then, the mixture was diluted with toluene (336 L), cooled down to – 12.4
°C and potassium t-butoxide (42.3 kg, 377 moles) was added in portions (10) maintaining
– 13.7 °C ≤ T_{mass} ≤ – 2.8 °C. The mixture was post-agitated at about 0°C for 2.5 h,
10 quenched by adding ultra pure water (142.5 L) maintaining 2.1 °C ≤ T_{mass} ≤ 8.7 °C. The
aqueous layer (176 kg) was separated after 35 minutes of post-stirring allowing the
mixture to reach 15 °C and the toluene layer was washed with ultra pure water (142.5 L)
and the aqueous layer (162 kg) was separated. The organic layer was then concentrated
under reduced pressure (150 mbars) maintaining T_{mass} ≤ 60 °C in order to distill 162 kg
15 of toluene. The filtrates were then diluted with toluene (114 L) and treated with SiO₂
(Merck silica gel 60, 0.063-0.1 mm, 74.1 kg) under agitation at room temperature for 1.25
h. SiO₂ was filtered and rinsed with toluene (2x114 L). Then, the filtrates were
concentrated under reduced pressure (150 mbars) maintaining T_{mass} ≤ 60 °C in order to
distill 351.8 kg of toluene (KF : 0.01 % w/w H₂O).
20 The solution of 4-Benzyl-morpholine-2-carbonitrile (169.2 kg) was diluted with
toluene (157 L) and was cooled to 0°C and phenylmagnesiumchloride (25 wt. % solution
in THF, 213 kg, 389 moles, 1.36 molar equiv.) was slowly added (over 3.5 h) to the
reaction mixture, maintaining the temperature at – 3 °C ≤ T_{mass} ≤ 7 °C. The reaction
mixture was post-stirred for 2 hours at T_{mass} ≈ 0°C. Then, the quench was performed by
25 adding acetic acid (8.55 L, T_{mass} = 5 → 17.2 °C), post stirring 10 minutes and cooling to
5 °C before adding an acetic acid / water mixture (229 L, 33/67 v/v). During the quench,
addition was performed at such a rate that T_{mass} did not exceed 20°C (typical T_{mass} =
4.6 °C to 10.4 °C). The mixture was post-agitated overnight at RT and the aqueous layer
(285.8 kg) was extracted.

The toluene layer was cooled to 0°C and a 5 N NaOH aqueous solution (420.1 kg) was slowly added maintaining the temperature at $-2.4\text{ }^{\circ}\text{C} \leq T_{\text{mass}} \leq 11\text{ }^{\circ}\text{C}$. The reaction mixture was post-stirred for 1h and the aqueous layer (494.8 kg) was extracted. The toluene layer was concentrated under reduced pressure (50 mbars) maintaining $T_{\text{mass}} \leq 60\text{ }^{\circ}\text{C}$ in order to distill 356.2 kg of toluene and isopropanol (180.4 kg) was added. The toluene was stripped off under reduced pressure (100 mbars) maintaining $T_{\text{mass}} \leq 60\text{ }^{\circ}\text{C}$ in order to distill 186.4 kg of toluene and isopropanol (135 kg) was added again to the mixture. A last distillation of toluene was performed under reduced pressure (50 mbars) maintaining $T_{\text{mass}} \leq 60\text{ }^{\circ}\text{C}$ in order to distill 131 kg of toluene and isopropanol (49.4 kg) was finally added to the mixture and the solution was stirred at RT until crystallization (17 minutes).

Ultra pure water was added (125.4 L) and the mixture was stirred overnight at RT and cooled down to about 0 °C for 1 hour. The precipitate was filtered and rinsed with a cooled water/isopropanol 50/50 v/v solution (76.6 kg). The wet precipitate was dried under vacuum at $T_{\text{jack}} = 35^{\circ}\text{C}$ for 96 hours to obtain the title compound as an off-white powder with 59 % overall yield. The title compound can be resolved by the fractional crystallisation process described above.

Preparation of (4-Benzyl-morpholin-2-yl)-(3-fluoro-phenyl)-methanone.



20


To a solution of 4-Benzyl-morpholine-2-carbonitrile (10g, 50 mmol) in dry diethyl ether (100 ml) at $-10\text{ }^{\circ}\text{C}$ under an atmosphere of nitrogen was added (time of addition 30 minutes) a solution of 3-fluorophenylmagnesium bromide (0.5N solution in tetrahydrofuran, 120 ml, 60 mmol, 1.2 equivalents, available from Aldrich Chemical Company or Rieke Metals) and the reaction mixture was further stirred at $-10\text{ }^{\circ}\text{C}$ for 30 minutes. Then the reaction was allowed to warm to room temperature and stirred for one

25

hour. The reaction was then cooled to 0 °C and quenched by addition of hydrochloric acid (2N aqueous solution, 50 ml) and the resulting mixture was stirred for 30 minutes at 0 °C. Then the solution was concentrated *in vacuo* and the residue was taken-up by sodium hydroxide (2N aqueous solution, 60 ml). The aqueous solution was extracted with diethyl ether, the organics fractions were collected and dried (MgSO₄) and the solvent removed under reduced pressure to give the title compound as a brown oil (15g, 100%). FIA [M+H]⁺=300.1.

Preparation of 2-Chloromethyl-4-fluoro-1-methoxy-benzene.


a) (5-Fluoro-2-methoxy-phenyl)-methanol.



To a solution of 2-Methoxy-5-fluorobenzaldehyde (11.093g, 1 equiv.- available from Aldrich Chemical Company) in methanol at -10 °C under nitrogen atmosphere was added NaBH₄ (7.515g, 2.7 equiv.) portionwise. The solution was allowed to warm to room temperature and after 30 minutes the reaction solvent was removed under reduced pressure and replaced with dichloromethane. This solution was poured onto ice water and further extracted with dichloromethane. The organic fractions were collected and dried (MgSO₄) and the solvent removed under reduced pressure to give the title compound as an oil (9.794g, 87%). ¹H NMR (300MHz, CDCl₃): δ 2.58 (m, 1H), 3.81 (s, 3H), 4.63 (d, 2H, *J* = 6.3 Hz), 6.78 (dd, 1H, *J* = 8.9 and 4.3 Hz), 6.94 (td, 1H, *J* = 8.5 and 3.1Hz), 7.04 (dd, 1H, *J* = 8.7 and 3.1Hz).

b) 2-Chloromethyl-4-fluoro-1-methoxy-benzene.


128



Neat (5-Fluoro-2-methoxy-phenyl)-methanol (19.587g, 1 equiv.) was added to neat SOCl_2 (42.2 mL, 4.6 equiv.) at -78°C under a nitrogen atmosphere and the solution was then allowed to warm to room temperature and stirred until evolution of gas had
5 ceased. An equivalent volume of anhydrous toluene was added to the flask and the solution heated to 60°C . On cooling the reaction solution was poured onto ice water. The toluene layer was separated and dried (MgSO_4) and the solvent removed under reduced pressure. The crude material was sublimed ($60\text{--}80^\circ\text{C}/0.05\text{ mBarr}$) to give the title
10 compound as a white solid (13.40 g, 61%). ^1H NMR (300MHz, CDCl_3): δ 3.87 (s, 3H), 4.60 (s, 2H), 6.79-7.20 (m, 3H).

Preparation of 1-Chloromethyl-2-isopropoxy-benzene.

a) (2-Isopropoxy-phenyl)-methanol.




15

A mixture of 2-hydroxybenzyl alcohol (21.04g, 1 equiv., available from Aldrich Chemical Company), 2-isopropyl iodide (32.3 mL, 1.9 equiv., available from Aldrich Chemical Company) and K_2CO_3 (71.42g, 3 equiv.) in ethanol was refluxed for 3 hours. On cooling the reaction mixture was filtered and the solvent removed under reduced
20 pressure and replaced with dichloromethane, and then filtered and the solvent removed to give the title compound as an oil (27.751g, 99%). ^1H NMR (300MHz, CDCl_3): δ 1.37 (d, 6H, $J = 6.0\text{Hz}$), 3.55 (bs, 1H), 4.50-4.70 (m, 3H), 6.78-6.90 (m, 2H), 7.15-7.25 (m, 2H).

b) 1-Chloromethyl-2-isopropoxy-benzene.

129




The title compound was prepared using the general procedure outlined above for the preparation of 2-chloromethyl-4-fluoro-1-methoxy-benzene followed by the following treatment:

- 5 The crude reaction material was chromatographed on silica gel and eluted 1:9 ethyl acetate/heptane prior to distillation (40-60 °C/0.05 mBar). ¹H NMR (300MHz, CDCl₃): δ 1.37 (d, 6H, J = 6.0Hz), 4.50-4.70 (m, 3H), 6.80-7.00 (m, 2H), 7.23-7.30 (m, 2H).

10 **Synthesis of Compounds of Formula (IB).**

Example 1B: (S, R)-2-(2-Methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.


- 15 a) **1-(4-Benzyl-morpholin-2-yl)-2-(2-methoxy-phenyl)-1-phenyl-ethanol.**



- 20 Solid magnesium turnings (9.5 g, 28 equiv.) under nitrogen atmosphere at room temperature were stirred vigorously with a magnetic stirring bar overnight. The magnesium was then covered with dry diethyl ether and to the suspension was added 1,2-dibromoethane (50 µL). A cold bath was then applied followed by dropwise addition of 1-chloromethyl-2-methoxy-benzene (18.18 g, 5 equiv. available from Aldrich Chemical Company) in diethyl ether (71 mL) which maintained the temperature at up to 15 °C. The

resulting black suspension was stirred at room temperature for 30 minutes and cooled down at -20°C . A solution of (4-Benzyl-morpholin-2-yl)-phenyl-methanone (4g, 1 equiv.) in diethyl ether (50 mL) was then added dropwise *via canula*. The reaction mixture was left to warm to room temperature over two hours and then quenched by
5 addition of aqueous saturated solution of NaHCO_3 (50 mL). The aqueous solution was extracted with diethyl ether, the organic phase dried with MgSO_4 , evaporated *in vacuo* to give 7 g of a yellow amorphous solid. The compound was taken without further purification in the next step. FIA $[\text{M}+\text{H}]^+=404$.


10 b) 2-(2-Methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.



To a solution of 1-(4-Benzyl-morpholin-2-yl)-2-(2-methoxy-phenyl)-1-phenyl-ethanol (1 g, 1 equiv.) in ethyl acetate (100 mL) at room temperature under nitrogen atmosphere was added ammonium formate (3.9 g, 25 equiv.) followed by addition of
15 palladium on charcoal (10 %, 1g.). The reaction mixture was heated to reflux for 1 hour, cooled to room temperature and then filtered through Celite. All volatiles were evaporated under *vacuum*, and the resulting solid was purified via preparative HPLC. The isolated white solid was taken up in ethanol. Hydrogen chloride was added (large excess of 2M solution in diethyl ether) and the mixture was stirred until it became a clear
20 solution. Then all the volatiles were evaporated in *vacuo*, to give 650 mg of the title compound as white solid (75 %). ^1H NMR (300MHz, DMSO D_6) δ : 2.43-2.51 (m, 2H), 2.77-2.92 (m, 2H), 3.15-3.23 (m, 3H), 3.41 (s, 3H), 4.10-4.19 (m, 2H), 6.66-6.72 (m, 2H), 6.98-7.07 (m, 2H), 7.13-7.20 (m, 5H), 9.32 (bs, 2H). LCMS (12 minute method) $[\text{M}+\text{H}]^+=314$ @ Rt 3.96 min. single major peak.


Example 2B: (S, R) 2-(2-Ethoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.

a) 1-(4-Benzyl-morpholin-2-yl)-2-(2-ethoxy-phenyl)-1-phenyl-ethanol.



The procedure for the synthesis of example **1Ba**, 1-(4-Benzyl-morpholin-2-yl)-2-(2-methoxy-phenyl)-1-phenyl-ethanol, was followed using commercially available 2-ethoxybenzylmagnesium bromide (available from Rieke-Metals) as starting material and making non-critical variations, to yield the title compound. FIA $[M+H]^+=418$.


b) 2-(2-Ethoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.



The procedure for the synthesis of example **1Bb**, 2-(2-Methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride was followed making non-critical variations, to yield the title compound. ^1H NMR (300MHz, DMSO D_6) δ : 1.11 (t, 3H, $J=6.97\text{Hz}$), 2.43-2.56 (m, 1H), 2.81-2.96 (m, 2H), 3.17-3.27 (m, 3H), 3.55-3.67 (m, 2H), 3.84-3.92 (m, 1H), 4.05-4.20 (m, 2H), 6.68-6.74 (m, 2H), 7.01-7.18 (m, 8H), 8.92 (bs, 2H) ppm. LCMS (12 minute method) $[M+H]^+=328$ @ R_t 4.57 min. single major peak.

Example 3B: S, R) 2-(2-Isopropoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.

a) 1-(4-Benzyl-morpholin-2-yl)-2-(2-isopropoxy-phenyl)-1-phenyl-ethanol.



5

Solid magnesium turnings (4.6 g, 48 equiv.) under nitrogen atmosphere at room temperature were stirred vigorously with a magnetic stirring bar overnight. The magnesium was then covered with dry tetrahydrofuran. A cold bath was then applied followed by dropwise addition of 1-chloromethyl-2-isopropoxy-benzene (3.0 g, 4 equiv. prepared as described above) in tetrahydrofuran (40 mL). During slow addition of the electrophile no exotherm was observed so on completion of addition 3 crystals of Iodine were added to promote initiation of the reaction. After this addition the reaction temperature was allowed to spike to 50 °C then cooled rapidly to 8 °C before being left to warm to room temperature for one hour. The resulting black suspension was cooled down to -10 °C and a solution of (4-Benzyl-morpholin-2-yl)-phenyl-methanone (1.2 g, 1 equiv.) in tetrahydrofuran (10 mL) was then added dropwise. The reaction mixture was left to warm to room temperature over thirty minutes and then quenched by addition of aqueous saturated solution of NaHCO₃ (50 mL) prior to filtration through Celite. The aqueous solution was extracted with diethyl ether, the organic phase dried with MgSO₄, evaporated *in vacuo* to give 3 g of a yellow amorphous solid. The compound was taken without further purification in the next step. LCMS (6 minutes method) [M+H]⁺=432 @ Rt 3.25 min. major peak.


10

15

20

b) 2-(2-Isopropoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.


133



The procedure for the synthesis of example **1Bb**, 2-(2-Methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride was followed making non-critical variations, to yield the title compound. ^1H NMR (300MHz, MeOH D₃) δ : 1.12-1.16 (m, 6H), 2.51-2.55 (m, 1H), 2.89-3.14 (m, 4H), 3.56-3.60 (m, 1H), 3.82-3.92 (m, 1H), 3.99-4.03 (m, 1H), 4.17-4.22 (m, 1H), 4.36-4.44 (m, 1H), 6.50-6.55 (m, 1H), 6.66-6.73 (m, 2H), 6.92-6.98 (m, 1H), 7.07-7.20 (m, 5H) ppm. LCMS (12 minutes method) $[\text{M}+\text{H}]^+ = 342$ @ Rt 4.90 min. major peak.

Example 4B: (S, R) 1-(3-Fluoro-phenyl)-2-(2-methoxy-phenyl)-1-morpholin-2-yl-ethanol hydrochloride

a) **1-(4-Benzyl-morpholin-2-yl)-1-(3-fluoro-phenyl)-2-(2-methoxy-phenyl)-ethanol.**




15

A magnetically stirred 0.25M tetrahydrofuran solution of commercially available 2-methoxybenzylmagnesium bromide (available from Rieke-Metals) (80ml, 3equiv.) under nitrogen atmosphere was cooled to -10°C and to this was added neat (4-Benzyl-morpholin-2-yl)-1-(3-fluoro-phenyl)-methanone (2.1g, 1equiv.). The solution was allowed to warm to room temperature and reaction progress followed using mass

spectrometry. After 1.5 hours 2-methoxybenzylmagnesium bromide solution (14ml, 0.5equiv.) was again added to the reaction and after a further 0.5 hours an aqueous saturated solution of NaHCO₃ (50 mL) was added to halt the reaction. The aqueous solution was extracted with diethyl ether, the organic phase dried with MgSO₄,

5 evaporated *in vacuo* to give 2.8 g of a yellow amorphous solid. The compound was taken without further purification in the next step. LCMS (6 minutes method) [M+H]⁺=422 @ Rt 3.03 and 2.86 min. major peaks.


b) (S, R)-1-(3-Fluoro-phenyl)-2-(2-methoxy-phenyl)-1-morpholin-2-yl-ethanol
10 hydrochloride.



To a solution of 1-(4-Benzyl-morpholin-2-yl)-1-(3-fluoro-phenyl)-2-(2-methoxy-phenyl)-ethanol (2.8 g, 1 equiv.) in ethyl acetate (100 mL) at room temperature under nitrogen atmosphere was added ammonium formate (4.3 g, 10 equiv.) followed by
15 addition of palladium on charcoal (10 %, 2.7g.). The reaction mixture was heated to reflux for 1 hour, cooled to room temperature and then filtered through Celite. All volatiles were evaporated under *vacuum*, and the resulting solid was purified via preparative HPLC to give the desired diastereoisomers. The active enantiomer was obtained after a further preparative chiral HPLC separation. The active enantiomer, a
20 white solid, was next taken up in ethanol and hydrogen chloride was added (large excess of 2M solution in diethyl ether) and the mixture was stirred until it became a clear solution. Then all the volatiles were evaporated in *vacuo*, to give 447mg of the title compound as white solid. ¹H NMR (300MHz, DMSO D₆) δ: 2.49-2.53 (m, 1H), 2.80-2.93 (m, 2H), 3.12-3.33 (m, 4H), 3.41 (s, 3H), 3.85-3.92 (m, 1H), 4.07-4.20 (m, 2H),
25 6.70-6.75 (m, 2H), 6.92-7.10 (m, 5H), 7.20-7.27 (m, 1H), 9.08 (bs, 2H). LCMS (12 minutes method) [M+H]⁺=332. Rt 4.11 min.

Example 5B: (S, R) 1-Morpholin-2-yl-1-phenyl-2-(2-trifluoromethoxy-phenyl)-ethanol hydrochloride

5 **a) 1-(4-Benzyl-morpholin-2-yl)-1-phenyl-2-(2-trifluoromethoxy-phenyl)-ethanol.**




Magnesium turnings (24.2 g, 0.935 mole, 2 eq.) and diethyl ether (300 ml) were loaded in a reactor under N₂. A solution of 2-trifluoromethoxybenzyl bromide (165 g, 0.647 mole, 1.3 eq.) in diethyl ether (300 ml) was loaded in an addition funnel. Iodine crystals and a small amount of the 2-trifluoromethoxybenzyl bromide solution were added and the reaction mixture was stirred to initiate the reaction. The remainder of the 2-trifluoromethoxybenzyl bromide solution was then added drop-wise maintaining the temperature of the reaction mixture below 35°C. The mixture was stirred for another 5 minutes at 23°C after completion of the addition. A solution of (4-Benzyl-morpholin-2-yl)-phenyl-methanone (140 g, 0.498 mole) in diethyl ether (2.1 L) was added drop-wise, maintaining the temperature of the reaction mixture below 25°C. The solution obtained was stirred for 1 hour at 20°C. The reaction mixture was quenched through the addition of a saturated aqueous NaHCO₃ solution (700 ml) and water (700 ml). The solids were filtered and washed with diethyl ether (200 ml). The filtrates were loaded into a separation funnel and the layers were separated. The aqueous layer was extracted with diethyl ether (1 L). The organic layers were combined and the filtrates were concentrated under vacuum to about 2 liters. The solution was dried over MgSO₄, filtered and the filter cake was washed with diethyl ether (200 ml). The filtrate was concentrated under vacuum to orange oil. The residue was twice dissolved in toluene (500 ml) and concentrated to a solid product. The yield of crude title compound was 235 g (103%). ¹H-NMR (CDCl₃):

6.80-7.07 ppm, 11 H, mp; 7.04-7.01 ppm, 1H, mp; 7.01-6.86 ppm, 1H, dt; 6.84-6.80 ppm, 1H, d; 3.98-4.03 ppm, 1H, dt; 3.86-3.89 ppm, 1H, dd; 3.70-3.60 ppm, 1H, dt; 3.52-3.58 ppm, 1H, d; 3.37-3.42 ppm, 1H, d; 3.13-3.37 ppm, 1H, d; 3.05-3.08 ppm, 1H, d; 2.44-2.45 ppm, 1H, d; 2.30-2.00 ppm, 3H, mp.

5

b) (S, R) 1-Morpholin-2-yl-1-phenyl-2-(2-trifluoromethoxy-phenyl)-ethanol hydrochloride.




A stainless steel Buchi hydrogenation reactor was loaded with 1-(4-Benzyl-morpholin-2-yl)-1-phenyl-2-(2-trifluoromethoxy-phenyl)-ethanol (230 g, 0.503 mole), methanol (1 L), a suspension of Pd/C (10%, 46 g, 20% loading) in methanol (500 ml), and methanol (500 ml) from equipment rinses. A solution of HCl in ethanol (1.6N, 460 ml, 0.736 mole, 1.5 eq.) was added and the reactor was pressurized with H₂ (3 Bar). The reaction mixture was heated to 40°C and stirred for 3 hours. The reaction mixture was cooled to 20°C and flushed with N₂. The catalyst was filtered off and washed with methanol (0.5 L). The filtrates were concentrated under vacuum to a yellow solid. The yield of crude title compound was 198 g (97.5%). A reactor was loaded with crude title compound (190 g, 0.47 mole) and toluene (6.65 L) under N₂. The suspension was heated under reflux and toluene (150 ml) was added until all solid dissolved. The solution was stirred for 15 minutes more under reflux and then cooled slowly to 20°C. The suspension was stirred for 1 hour at 20°C. The solid was filtered, washed with toluene (680 ml), and dried at 40°C under vacuum. The yield of pure anhydrous title compound was 158.5 g (83.4%).

Alternatively, the following method can be used. In a glass-lined nitrogen purged hydrogenator are charged 1-(4-Benzyl-morpholin-2-yl)-1-phenyl-2-(2-trifluoromethoxy-

phenyl)-ethanol hydrochloride (150g, 303.7 mmol), demineralized water (352 mL), *i*-PrOH (375 mL) and 5% Pd/C (30 g, 50% water, Johnson & Matthey type 440). The heterogeneous reaction mixture was then purged 5 times with 25 psi nitrogen then purged 5 times with 50 psi hydrogen, and the hydrogenation was performed at RT. The initial
5 T_{mass} was 22°C and the maximum T_{mass} during the hydrogenation was 23°C. The reactor was stirred vigorously. In-process analysis after 2 hours indicated complete hydrogenolysis. The hydrogenation was stopped after 3 hours. The nitrogen purged reaction mixture was then filtered at RT through an hyflo filter (56 g), impregnated beforehand with 75 mL of a 50/50 v/v isopropanol/water mixture and washed with 300
10 mL of a 50/50 v/v isopropanol/water mixture. The filtrates were stored overnight at RT. The filtrates were concentrated at 40-50°C under reduced pressure (typical 622 g distilled). The reaction mixture was cooled to RT and post-agitated. After 3 hours, 1 mL of the solution was taken and cooled to 0°C to initiate crystallization. These seeds were added to the reaction mixture and precipitation was observed within a few minutes. The
15 mixture was post-agitated at RT for 2 hours. The crystals were filtered and rinsed with H₂O (30 mL). Then, the precipitate was dried under reduced pressure (400 mmHg) with a nitrogen flow (0.1 bar) for 4 hours affording the title compound as the hydrate polymorph (103.5 g, 81% yield).


20 **Example 6B: (S, R) 2-Biphenyl-2-yl-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride**

a) **1-(4-Benzyl-morpholin-2-yl)-2-biphenyl-2-yl-1-phenyl-ethanol.**



1-(4-Benzyl-morpholin-2-yl)-2-(2-bromo-phenyl)-1-phenyl-ethanol (0.50 g, 1.0 equiv. prepared according to Example 15Ba below) and phenylboronic acid (0.402 g, 3.0 equiv., available from Aldrich Chemical Company) were suspended in a mixture ethanol/water (2/1, 7.5 mL) and Pd(Ph₃)₄ (0.022 g, 0.04 equiv.), then K₂CO₃ (0.654 g, 4.30 equiv.) were added. The mixture was heated to 80°C under nitrogen atmosphere. After 16 hours, the reaction was cooled down to room temperature and filtered through Celite, then extracted with ethyl acetate. The organic layers were combined, dried with MgSO₄, filtered and concentrated in vacuo yielding a yellow oil, which was purified by column chromatography on silica gel (10% EtOAc:Hexane) to give 0.491g (98%) of the title compound as a white solid.

b) (S, R) 2-Biphenyl-2-yl-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.




The procedure for the synthesis of example 1Bb, 2-(2-methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride, was followed making non-critical variations, to yield the title compound. ¹H NMR (300MHz, DMSO D₆) δ: 2.16-2.20 (m, 1H), 2.54-2.62 (m, 1H), 2.67-2.76 (m, 1H), 2.85-2.89 (m, 1H), 3.24 (s, 2H), 3.61-3.69 (m, 2H), 3.93-3.98 (m, 1H), 5.14 (bs, 1H), 6.80-6.92 (m, 5H), 7.04-7.17 (m, 5H), 7.27-7.30 (m, 3H), 7.36-7.39 (m, 1H). LCMS (12 minutes method) [M+H]⁺=360 @ Rt 5.15 min. single major peak.

Example 7B: (S, R) 2-(2-Chloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride


a) 1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-phenyl)-1-phenyl-ethanol.

139



The procedure for the synthesis of example **1Ba**, 1-(4-Benzyl-morpholin-2-yl)-2-(2-methoxy-phenyl)-1-phenyl-ethanol, was followed using 2-chlorobenzyl chloride (available from Aldrich Chemical Company) as starting material and making non-critical variations, to yield the title compound. FIA $[M+H]^+$ =408 and 410.

b) (S, R) 2-(2-Chloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride




The procedure for the synthesis of example **5Bb**, (S, R) 1-Morpholin-2-yl-1-phenyl-2-(2-trifluoromethoxy-phenyl)-ethanol hydrochloride, was followed making non-critical variations, to yield the title compound. ^1H NMR (300MHz, DMSO D_6) δ : 2.45-2.54 (m, 1H), 2.84-2.93 (m, 2H), 3.17-3.22 (m, 1H), 3.33-3.38 (m, 3H), 3.89-3.97 (m, 1H), 4.14-4.18 (m, 2H), 7.06-7.11 (m, 2H), 7.15-7.26 (m, 7H), 9.24 (bs, 2H) ppm. LCMS (12 minutes method) $[M+H]^+$ =318-320 @ Rt 4.36 min. single peak.

Example 8B: (S, R) 2-(5-Fluoro-2-methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride

a) 1-(4-Benzyl-morpholin-2-yl)-2-(5-fluoro-2-methoxy-phenyl)-1-phenyl-ethanol.

140




Magnesium turnings (21.6 g, 0.888 mole, 2 eq.) and diethyl ether (300 ml) were loaded in a reactor under N₂. A solution of 5-fluoro-2-methoxybenzyl chloride (116 g, 0.664 mole, 1.5 eq.) in diethyl ether (200 ml) was loaded in an addition funnel. Iodine crystals and a small amount of the 5-fluoro-2-methoxybenzyl chloride solution were added and the reaction mixture was stirred to initiate the reaction. The remainder of the 5-fluoro-2-methoxybenzyl chloride solution was then added drop-wise maintaining the temperature of the reaction mixture below 28 °C. The mixture was stirred for another 5 minutes at 19 °C after completion of the addition and a white suspension was formed. A solution of (4-Benzyl-morpholin-2-yl)-phenyl-methanone (125 g, 0.444 mole) in diethyl ether (1.8 L) was added drop-wise, maintaining the temperature of the reaction mixture below 25 °C. The suspension obtained was stirred for 2 hours. The reaction mixture was quenched through the addition of a saturated aqueous NaHCO₃ solution (625 ml) and water (500 ml), maintaining the temperature below 20 °C. The mixture was stirred for 30 minutes and the solids were filtered, washed with water (125 ml) and diethyl ether (200 ml). The filtrates were loaded into a separation funnel and the layers were separated. The aqueous layer was extracted with diethyl ether (1 L). The organic layers were combined and dried over MgSO₄, filtered and the filter cake was washed with diethyl ether (100 ml). The filtrates were concentrated under vacuum. The yield of title compound was 201 g as a yellow solid (107%). Title compound (200 g, 0.474 mole) was then suspended in isopropanol (400 ml) under N₂. The suspension was heated under reflux until all solids were dissolved. The solution is allowed to cool to 20 °C over 4 hours under stirring. The solid is filtered, washed with isopropanol (100 ml) and dried at 40°C under vacuum. The yield of pure title compound is 158 g (79%). ¹H-NMR (CDCl₃): 6.99-7.26 ppm, 10H, mp; 6.60-6.71 ppm, 1H, dt; 6.49-6.60 ppm, 1H, dd; 6.31-6.44 ppm, 1H, dd; 3.92-4.01 ppm, 1H, dt; 3.80-3.90 ppm, 1H, dd; 3.64-3.73 ppm, 1H, dd; 3.59-3.64 ppm, 1H, d; 3.52-

141

3.59 ppm, 3+1 H, 2s; 3.37-3.45 ppm, 1H, d; 3.07-3.17 ppm, 1H, d; 2.84-2.92 ppm, 1H, d; 2.43-2.53 ppm, 1H, d; 2.20-2.28 ppm, 1H, d; 1.98-2.11 ppm, 2H, mp.

b) (S, R) 2-(5-Fluoro-2-methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol
5 hydrochloride




A glass hydrogenation flask was loaded with methanol (1.55 L), Pd/C (10%, 31 g, 20% loading), 1-(4-benzyl-morpholin-2-yl)-2-(5-fluoro-2-methoxy-phenyl)-1-phenyl-ethanol (155 g, 0.368 mole) and a solution of HCl in ethanol (2.5N, 233 ml, 0.582 mole, 10 1.6 eq.). The reactor was mounted on a Parr instrument and pressurized with H₂ (49 Psi). The reaction mixture was shaken overnight between 20°C and 15°C. The catalyst was filtered off and washed with methanol (0.5 L). The filtrates were concentrated under vacuum. The yield of crude title compound was 109.5 g (81%). The catalyst was washed again with methanol (2 x 500 ml). The filtrates were combined and concentrated under 15 vacuum. The yield of the second crop of crude title compound was 21.7 g (16%). A reactor was loaded with crude title compound (131 g, 0.356 mole) and isopropanol (1,3 L) under N₂. The suspension was heated under reflux for 4 hours. The mixture was cooled to 20°C and the solid was filtered, washed with isopropanol (130 ml), and dried at 50°C under vacuum. The yield of pure title compound was 115.9 g (88.5% yield).

20

Example 9B: (S, R) 1-Morpholin-2-yl-1-phenyl-2-(2-trifluoromethylsulfanyl-phenyl)-ethanol acetate


a) 1-(4-Benzyl-morpholin-2-yl)-1-phenyl-2-(2-trifluoromethylsulfanyl-phenyl)-
25 ethanol.

142



The procedure for the synthesis of example **1Ba**, 1-(4-benzyl-morpholin-2-yl)-2-(2-methoxy-phenyl)-1-phenyl-ethanol, was followed using 1-bromomethyl-2-trifluoromethylsulfanyl-benzene (available from Fluorochem Ltd.) as starting material and making non-critical variations, to yield the title compound. ¹H NMR (300MHz, CDCl₃) δ: 2.05-2.33 (m, 3H), 2.49-2.65 (m, 1H), 3.10-3.35 (m, 2H), 3.43-3.55 (m, 1H), 3.67-3.89 (m, 2H), 3.91-4.08 (m, 2H), 4.09-4.22 (m, 1H), 6.91-7.05 (m, 1H), 7.10-7.42 (m, 12H), 7.50-7.63 (m, 1H) ppm.

b) (S, R) 1-Morpholin-2-yl-1-phenyl-2-(2-trifluoromethylsulfanyl-phenyl)-ethanol acetate




To a solution of 1-(4-benzyl-morpholin-2-yl)-1-phenyl-2-(2-trifluoromethylsulfanyl-phenyl)-ethanol (218 mg g, 1 equiv.) and solid supported Hunig's base (available from Argonaut, 1g, 5 equiv.) in dry tetrahydrofuran (4 mL) at 0 °C under nitrogen atmosphere was added ACE-Cl (502 μL, 10 equiv.). The reaction mixture was left to warm to room temperature for 48 hours. All volatiles were evaporated under vacuum, and the resulting solid was taken-up with methanol (50 mL) and stirred at room temperature overnight. The solution was filtered through acid ion exchange column and the required fractions evaporated to dryness. The resulting solid was purified *via* preparative HPLC to give 62 mg of the title compound as a colourless oil. ¹H NMR (300MHz, CDCl₃) δ: 2.01 (s, 3H), 2.43-2.47 (m, 1H), 2.63-2.70 (m, 1H), 2.81-2.94 (m,

2H), 3.24 (d, 1H, $J=13.57\text{Hz}$), 3.85-3.96 (m, 2H), 4.01-4.05 (m, 1H), 4.09-4.13 (m, 1H), 4.45 (bs, 4H), 6.90-6.93 (m, 1H), 7.13-7.26 (m, 7H), 7.55-7.58 (m, 1H) ppm. LCMS (12 minute method) $[M+H]^+=384$ @ Rt 5.13 min. single peak.

5 **Example 10B: (S, R) 1-Morpholin-2-yl-1-phenyl-2-(2-trifluoromethyl-phenyl)-ethanol**


a) **4-Benzyl-2-(2-phenyl-oxiranyl)-morpholine.**



10 To a mixture of trimethylsulfoxonium iodide (783 mg, 1equiv.) and sodium hydride (142 mg, 1 equiv.) in dimethylformamide (17 mL) at 0 °C under nitrogen atmosphere was added dimethylsulfoxide (251 μL , 1 equiv.) and the resulting suspension was stirred for 30 minutes. A solution of (4-Benzyl-morpholin-2-yl)-phenyl-methanone (1 g, 1equiv.) in dimethylformamide (10 mL) was then added dropwise. Stirring was
15 continued for 30 minutes and the reaction was stopped by addition of water (50 mL). The aqueous solution was extracted with diethyl ether, the organic phase dried with MgSO_4 , and evaporated *in vacuo*. The crude material was purified using a column chromatography on silica gel eluting with a mixture of ethyl acetate/heptane (20/80) to give 825 mg of the title compound as a colourless oil (78 %), mixture of two
20 diastereoisomers. LCMS (6 minute method) $[M+H]^+=296$ @ Rt 1.88 min. single peak.


b) **1-(4-Benzyl-morpholin-2-yl)-1-phenyl-2-(2-trifluoromethyl-phenyl)-ethanol.**

144



To a suspension of magnesium turnings in tetrahydrofuran (2mL) at room temperature under nitrogen atmosphere was added a solution of 1-bromo-2-trifluoromethyl-benzene (7.6g, 5equiv., available from Acros) in tetrahydrofuran (32 mL) and the mixture was stirred for an hour. The solution was cooled to -78°C and copper iodide (646 mg) was added followed by dropwise addition of a solution of 4-Benzyl-2-(2-phenyl-oxiranyl)-morpholine (2g, 1 equiv.) in tetrahydrofuran (10 mL). The resulting mixture was warmed to room temperature over 2 hours and then treated with water (10 mL). The solution was extracted with diethyl ether, the organic phase dried with MgSO_4 , and evaporated *in vacuo*. The crude material was purified using a column chromatography on silica gel eluting with a mixture of ethyl acetate/heptane (10/90) to give 352 mg of the title compound as a colourless oil (12 %). LCMS (6 minutes method) $[\text{M}+\text{H}]^+=442$ @ Rt 3.05 min. major peak.

c) (S, R) 1-Morpholin-2-yl-1-phenyl-2-(2-trifluoromethyl-phenyl)-ethanol



To a solution of 1-(4-Benzyl-morpholin-2-yl)-1-phenyl-2-(2-trifluoromethyl-phenyl)-ethanol (352 mg, 1 equiv.) in ethanol (15 mL) at room temperature under nitrogen atmosphere was added ammonium formate (507 mg g, 10 equiv.) followed by addition of palladium on charcoal (10 %, 355 mg.). The reaction mixture was heated to reflux for 1 hour, cooled to room temperature and then filtered through Celite. All volatiles were evaporated under vacuum to give 265 mg of the title compound as white solid (94 %). The enantiomeric mixture was resolved using chiral HPLC, to give the title


145

compound as a single enantiomer. ^1H NMR (300MHz, CDCl_3) δ : 2.25-2.30 (m, 1H), 2.56-2.64 (m, 1H), 2.75-2.87 (m, 2H), 3.18 (d, 1H, $J=14.88\text{Hz}$), 3.71-3.81 (m, 2H), 3.89 (d, 1H, $J=14.88\text{Hz}$), 4.02-4.05 (m, 1H), 6.83-6.86 (m, 1H), 7.09-7.34 (m, 7H), 7.53-7.55 (m, 1H) ppm. LCMS (12 minute method) $[\text{M}+\text{H}]^+=352$ @ R_t 4.73 min. single peak.

5

Example 11B: (S, R) 2-(2-Chloro-phenyl)-1-(3-fluoro-phenyl)-1-morpholin-2-yl-ethanol hydrochloride

a) 1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-phenyl)-1-(3-fluoro-phenyl)-ethanol.




10

The procedure for the synthesis of **4Ba**, 1-(4-Benzyl-morpholin-2-yl)-1-(3-fluoro-phenyl)-2-(2-methoxy-phenyl)-ethanol was followed using 2-chlorobenzyl chloride (available from Aldrich Chemical Company) as starting material, and making non-critical variations, to yield the title compound which was taken without further purification in the next step. LCMS (6 minutes method) $[\text{M}+\text{H}]^+=426$ @ R_t 2.85 min. major peak.

15

b) (S, R) 2-(2-Chloro-phenyl)-1-(3-fluoro-phenyl)-1-morpholin-2-yl-ethanol hydrochloride




20

To a solution of 1-(4-Benzyl-morpholine-2-yl)-2-(2-chloro-phenyl)-1-(3-fluoro-phenyl)-ethanol. (3.2g, 1 equiv.) in dry 1,2-dichloroethane (40 mL) under nitrogen atmosphere was added ACE-Cl (20.33 g, 5 equiv.). The reaction mixture was stirred at

room temperature overnight then refluxed until completion. All volatiles were evaporated under vacuum, and the resulting residue redissolved in acetonitrile. This solution was filtered through an ion exchange column and the filtrate taken-up with methanol (50 mL) and refluxed for 3h. The solution was again filtered through acid ion exchange column and the required fractions evaporated to dryness. The resulting solid was next purified via preparative HPLC followed by chiral HPLC. The purified active enantiomer was taken up in ethanol and hydrogen chloride was added (large excess of 2M solution in diethyl ether) and the mixture stirred. Then all the volatiles were evaporated in vacuo, to give 519mg of the title compound as a white solid (18 %). ¹H NMR (300MHz, DMSO D6) δ : 2.43-2.54 (m, 1H), 2.81-2.95 (m, 2H), 3.16-3.23 (m, 1H), 3.30-3.44 (m, 2H), 3.54 (bs, 1H), 3.92-4.00 (m, 1H), 4.15-4.23 (m, 2H), 6.96-7.29 (m, 8H), 9.32-9.45 (m, 2H). LCMS (12minute method) $[M+H]^+=336$.

Example 12B: (S, R) 1-Morpholin-2-yl-1-phenyl-2-*o*-tolyl-ethanol hydrochloride


a) 1-(4-Benzyl-morpholin-2-yl)-1-phenyl-2-*o*-tolyl-ethanol.



The procedure for the synthesis of example **1Ba**, 1-(4-benzyl-morpholin-2-yl)-2-(2-methoxy-phenyl)-1-phenyl-ethanol, was followed using commercially available 2-methylbenzylmagnesium bromide (available from Rieke-Metals) as starting material and making non-critical variations, to yield the title compound. FIA $[M+H]^+=388$.

b) (S, R) 1-Morpholin-2-yl-1-phenyl-2-*o*-tolyl-ethanol hydrochloride

147




The procedure for the synthesis of example **1Bb**, 2-(2-methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride was followed making non-critical variations, to yield the title compound. ^1H NMR (300MHz, DMSO D6) δ : 1.62 (s, 3H),
5 2.40-2.58 (m, 1H), 2.78-3.01 (m, 2H), 3.03-3.09 (m, 1H), 3.15-3.31 (m, 2H), 3.90-4.05 (m, 1H), 4.15-4.25 (m, 2H), 6.89-7.28 (m, 9H), 9.21-9.55 (m, 2H). LCMS (12 minute method) $[\text{M}+\text{H}]^+ = 298$ single peak.

Example 13B: (S, R) 1-Morpholin-2-yl-1,2-diphenyl-ethanol hydrochloride.

10


a) 1-(4-Benzyl-morpholin-2-yl)-1,2-diphenyl-ethanol.



The procedure for the synthesis of example **1Ba**, 1-(4-benzyl-morpholin-2-yl)-2-(2-methoxy-phenyl)-1-phenyl-ethanol, was followed using commercially available
15 benzylmagnesium bromide (available from TCI America) as starting material and making non-critical variations, to yield the title compound. LCMS $[\text{M}+\text{H}]^+ = 374.1$ major single peak @ 3.82 min.

b) (S, R) 1-Morpholin-2-yl-1,2-diphenyl-ethanol hydrochloride


148



The procedure for the synthesis of example **1Bb**, 2-(2-methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride was followed making non-critical variations, to yield the title compound. ^1H NMR (300MHz, CDCl_3) δ : 2.36-2.41 (m, 1H),
 5 2.64-2.71 (m, 1H), 2.78-2.91 (m, 3H), 3.16-3.32 (m, 2H), 3.73-3.82 (m, 2H), 4.08-4.11 (m, 1H), 6.80-6.83 (m, 2H), 7.07-7.12 (m, 3H), 7.16-7.27 (m, 6H). LCMS $[\text{M}+\text{H}]^+=284.1$ single peak @ 3.82 minutes.

Example 14B: (S, R) 2-(2-Fluoro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride


a) 1-(4-Benzyl-morpholin-2-yl)-2-(2-fluoro-phenyl)-1-phenyl-ethanol.



The procedure for the synthesis of example **1Ba**, 1-(4-benzyl-morpholin-2-yl)-2-(2-methoxy-phenyl)-1-phenyl-ethanol, was followed using commercially available 2-fluoro-benzylmagnesium chloride (available from Rieke Metals) as starting material and making non-critical variations, to yield the title compound. FIA $[\text{M}+\text{H}]^+=392.1$.

b) (S, R) 2-(2-Fluoro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride


149



The procedure for the synthesis of example **1Bb**, 2-(2-methoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride was followed making non-critical variations, to yield the title compound. ¹H NMR (300MHz, DMSO D6) δ: 2.40-2.56 (m, 1H), 2.78-2.97 (m, 2H), 3.17-3.29 (m, 3H), 3.89-3.96 (m, 1H), 4.14-4.19 (m, 2H), 5.47 (bs, 1H), 6.82-6.94 (m, 2H), 7.01-7.25 (m, 7H), 9.28-9.38 (m, 2H). LCMS [M+H]⁺=302.1 single major peak @ 3.82 minutes.

Example 15B: (S, R) 2-(2-bromo-phenyl)-1-phenyl-1-morpholin-2-yl-ethanol.


a) 1-(4-Benzyl-morpholin-2-yl)-2-(2-bromo-phenyl)-1-phenyl-ethanol.



The procedure for the synthesis of example **1Ba**, 1-(4-Benzyl-morpholin-2-yl)-2-(2-methoxy-phenyl)-1-phenyl-ethanol, was followed using commercially available 2-bromobenzylmagnesium bromide (available from Rieke-Metals) as starting material and making non-critical variations, to yield the title compound. FIA [M+H]⁺= 452/454.

b) (S, R) 1-Morpholin-2-yl-2-(2-bromo-phenyl)-1-phenyl-ethanol.


150



The procedure for the synthesis of example **5Bb**, (S, R) 1-Morpholin-2-yl-1-phenyl-2-(2-trifluoromethoxy-phenyl)-ethanol, was followed making non-critical variations, to yield the title compound. ¹H NMR (300MHz, CDCl₃) δ: 2.64-2.68 (m, 1H), 3.02-3.21 (m, 2H), 3.27-3.33 (m, 3H), 3.45-3.50 (m, 1H), 3.63-3.68 (m, 1H), 3.99-4.09 (m, 1H), 4.20-4.24 (m, 1H), 4.29-4.34 (m, 1H), 4.87 (s, 1H), 6.98-7.21 (m, 2H), 7.24-7.59 (m, 7H) ppm. LCMS (6 minutes method) [M+H]⁺= 362.3 @ Rt 2.85 min. single peak.

Example 16B: (S, R) 2-(2'-chloro[1-1'biphenyl]-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride


a) **2-(2'-chloro[1-1'biphenyl]-2-yl)-1-phenyl-1-[4-(phenylmethyl)morpholin-2-yl]ethanol.**



The procedure for the synthesis of example **6Ba**, was followed using 2-chloro phenyl boronic acid (available from Aldrich Chemical Company) as starting material and making non-critical variations, to yield the title compound. FIA [M+H]⁺= 485

b) **(S, R) 2-(2'-chloro[1-1'biphenyl]-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride**


151



The procedure for the synthesis of example **6Bb**, was followed making non-critical variations, to yield the title compound. ^1H NMR (300MHz, CDCl_3) δ : 2.10-2.21 (m, 1H), 2.57-2.65 (m, 1H), 2.62-2.75 (m, 1H), 2.83-2.87 (m, 1H), 3.20 (s, 2H), 3.63-3.70 (m, 2H), 3.95-3.97 (m, 1H), 5.12 (bs, 1H), 6.80-6.92 (m, 5H), 7.04-7.17 (m, 5H), 7.27-7.37 (m, 3H). LCMS (12 minutes method) $[\text{M}+\text{H}]^+=393$ @ Rt 4.75 min. single major peak.

Example 17B: 4-Fluoro-2-(2-morpholin-2-yl-2-phenylpropyl)phenol hydrochloride

a) 4-Fluoro-2-(2-morpholin-2-yl-2-phenylpropyl)phenol hydrochloride



Sodium thiomethoxide (13 eq, 186 mg) was added at once to a solution of 2-{2-[5-fluoro-2-(methoxy)phenyl]-1-methyl-1-phenylethyl}morpholine hydrochloride (75.2 mg, 0.204 mmol, synthesized as described in Example 8 above) in anhydrous DMF (3 ml) in a microwave vessel. Upon addition, the reaction vessel was sealed and heated up in a CEM-Discovery microwave at 150 Watts, reaching 110 °C in 5 minutes and maintaining this temperature 6 minutes. The reaction vessel was cooled to room temperature and the reaction mixture taken into methanol (5 ml) and purified by SCX-2 chromatography to obtain the free base as clear oil (50 mg). The hydrochloride salt was obtained following general procedures as a white solid (52 mg, 72 % after salt formation.). MW 353.83; $\text{C}_{18}\text{H}_{22}\text{NO}_3\text{FCl}$; ^1H NMR (CD_3OD): 7.29-7.26 (2H, m), 7.20-7.08 (2H, m), 6.53-6.50 (2H, m), 6.30-6.26 (1H, m), 4.18 (1H, dd, 12.6 Hz, 2.6 Hz), 4.02 (1H, dd, 10.9 Hz, 2.3 Hz),

3.86 (1H, td, 12.6 Hz, 2.6 Hz), 3.60 (1H, $\frac{1}{2}$ AB), 3.16 (1H, d, 12.6 Hz), 3.08-2.90 (3H, m), 2.58 (1H, m); ^{19}F NMR (CD_3OD) -128.4 ; LCMS: (12 min method) m/z 318.1 $[\text{M}-\text{HCl}+\text{H}]^+$ @ Rt 3.954 min.


5 **Example 18B: 2-(2-Fluoro-6-chloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.**

a) **1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-6-fluoro -phenyl)-1-phenyl-ethanol.**



10 To a stirred solution of 2-chloro-6-fluorobenzyl magnesium chloride (12.8mL, 3.20 mmol, 3 equiv., available from Rieke Metals) in anhydrous tetrahydrofuran (15 ml) at 0 °C under nitrogen was added a solution of (4-Benzyl-morpholin-2-yl)-phenyl-methanone (300mg, 1.07mmol, 1 equiv.) in tetrahydrofuran (5ml) dropwise over 15 minutes. The reaction was then stirred at 0 °C for one hour. The reaction mixture was
15 allowed to warm to room temperature over two hours and stirred for a further 18h. The solvent was then evaporated "*in vacuo*" and the residue redissolved in dichloromethane (30mL). The organic solution was washed with aqueous saturated solution of NaHCO_3 (50 mL). The aqueous solution was extracted with dichloromethane using a hydrophobic phase separator. The dichloromethane was evaporated "*in vacuo*" and redissolved in
20 methanol (2 mL). The sample was bound to SCX-2 (5g) and washed with methanol (30mL). The sample was eluted using 2M ammonia in methanol (30mL). The solvent was then evaporated using a reacti-therm blow down station to give 450 mg of a yellow amorphous solid. This material was used in step b) without further purification. LCMS (6 minutes method) $[\text{M}+\text{H}]^+ = 426$ @ Rt 3.27 min. major peak.

b) **2-(2-Fluoro-6-chloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.**




To a solution of 1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-6-fluoro -phenyl)-1-phenyl-ethanol (450mg, 1 equiv.) in ethyl acetate (15mL) at room temperature under nitrogen atmosphere was added ammonium formate (1.69 g, 25 equiv.) followed by addition of palladium on charcoal (10 %, 450g.). The reaction mixture was heated to reflux for 1.5 hours, cooled to room temperature and then filtered through Celite. All volatiles were evaporated under *vacuum*, and the resulting solid was purified via preparative HPLC. The isolated white solid was taken up in ethanol. Hydrogen chloride was added (large excess of 2M solution in diethyl ether) and the mixture was stirred until it became a clear solution. Then all the volatiles were evaporated "in vacuo", to give 147 mg of the title compound as white solid. ¹H NMR (300MHz, CD₃OD D₄) δ: 2.51-2.61 (d, 1H), 2.79-2.91 (t, 1H), 2.96-3.09 (m, 1H), 3.09-3.16 (m, 1H), 3.32-3.54 (q, 2H), 3.82-3.97 (t, 1H), 4.09-4.24 (t, 2H), 6.73-6.84 (t, 1H), 6.93-7.08 (m, 2H), 7.08-7.21 (m, 5H). LCMS (12 minutes method) [M+H]⁺ = 336 @ Rt 4.44 min. single major peak.

Example 19B: 2-(2,5-Dimethoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride

20


a) **1-(4-Benzyl-morpholin-2-yl)-2-(2,5-dimethoxy-phenyl)-1-phenyl-ethanol.**

154



The procedure for the synthesis of example **18Ba**, 1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-6-fluoro-phenyl)-1-phenyl-ethanol, using 2,5-dimethoxybenzyl magnesium chloride as starting material (available from Rieke Metals) was followed making non-critical variations, to yield the title compound. This material was used in step b) without further purification. LCMS (6 minutes method) $[M+H]^+ = 434$ @ Rt 3.10min. major peak.

b) 2-(2,5-Dimethoxy-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.



10

The procedure for the synthesis of example **18Bb**, 2-(2-Fluoro-6-chloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride, was followed making non-critical variations, to yield the title compound. ^1H NMR (300MHz, CD_3OD D4) δ : 2.53-2.62 (d, 1H), 2.86-3.10 (m, 3H), 3.13-3.27 (m, 2H), 3.36-3.51 (m, 6H), 3.81-3.93 (t, 1H), 4.02-4.08 (d, 1H), 4.15-4.25 (d, 1H), 6.28-6.33 (s, 1H), 6.49-6.64 (m, 2H), 7.06-7.22 (m, 5H). LCMS (12 minutes method) $[M+H]^+ = 344$ @ Rt 4.15 min. single major peak.


15

Example 20B: 2-(2,4-Difluoro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride

20

a) 1-(4-Benzyl-morpholin-2-yl)-2-(2,4-difluoro-phenyl)-1-phenyl-ethanol.

155



The procedure for the synthesis of example **18Ba**, 1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-6-fluoro-phenyl)-1-phenyl-ethanol, using 2,4-difluorobenzyl magnesium bromide as starting material (available from Rieke Metals) was followed making non-critical variations, to yield the title compound. This material was used in step b) without further purification. LCMS (6 minutes method) $[M+H]^+ = 410$ @ Rt 3.19 min. major peak.

b) **2-(2,4-Difluoro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.**




The procedure for the synthesis of example **18Bb**, 2-(2-Fluoro-6-chloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride, was followed making non-critical variations to yield the title compound. ^1H NMR (300MHz, CD_3OD D4) δ : 2.48-2.59 (d, 1H), 2.87-3.09 (m, 2H), 3.11-3.17 (m, 2H), 3.26-3.38 (m, 1H), 3.81-3.95 (t, 1H), 4.02-4.11 (d, 1H), 4.13-4.25 (d, 1H), 6.48-6.60 (m, 2H), 7.70-6.98 (m, 1H) 7.08-7.28 (m, 5H). LCMS (12 minutes method) $[M+H]^+ = 320$ @ Rt 4.20 min. major peak.

Example 21B: Preparation of 2-(2,6-Dichloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride


a) **1-(4-Benzyl-morpholin-2-yl)-2-(2,6-dichloro-phenyl)-1-phenyl-ethanol.**

156



The procedure for the synthesis of example **18Ba**, 1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-6-fluoro -phenyl)-1-phenyl-ethanol, using 2,6-dichlorobenzyl magnesium chloride as starting material (available from Rieke Metals) was followed making non-critical variations, to yield the title compound. This material was used in step b) without further purification. LCMS (6 minutes method) $[M+H]^+ = 442$ @ Rt 3.49 min. major peak.

b) 2-(2,6-Dichloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.



10


To a solution of 1-(4-Benzyl-morpholin-2-yl)-2-(2,6-dichloro-phenyl)-1-phenyl-ethanol (450mg, 1 equiv.) in ethyl acetate (15mL) at room temperature under nitrogen atmosphere was added ammonium formate (1.69 g, 25 equiv.) followed by addition of palladium on charcoal (10 %, 45mg.). The reaction mixture was heated to reflux for 3 hour, cooled to room temperature and then filtered through Celite. All volatiles were evaporated under *vacuum*, and the resulting solid was purified via preparative HPLC. The isolated white solid was taken up in ethanol. Hydrogen chloride was added (large excess of 2M solution in diethyl ether) and the mixture was stirred until it became a clear solution. Then all the volatiles were evaporated "in vacuo", to give 60 mg of the title compound as white solid. ^1H NMR (300MHz, CD_3OD D4) δ : 2.52-2.61 (d, 1H), 2.79-2.96 (t, 1H), 2.98-3.13 (t, 1H), 3.15-3.19 (s, 1H), 3.56-3.71 (q, 2H), 3.88-4.02 (t, 1H), 4.10-

20

4.21 (d, 1H), 4.29-4.39 (d, 1H), 6.97-7.08 (m, 1H), 7.10-7.21 (m, 7H). LCMS (12 minutes method) $[M+H]^+ = 352$ @ Rt 4.63 min. single major peak.


Example 22B: Preparation of 2-(2,5-Dichloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride

a) 1-(4-Benzyl-morpholin-2-yl)-2-(2,5-dichloro -phenyl)-1-phenyl-ethanol.



The procedure for the synthesis of example **18Ba**, 1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-6-fluoro -phenyl)-1-phenyl-ethanol, using 2,5-dichlorobenzyl magnesium chloride as starting material (available from Rieke Metals) was followed making non-critical variations, to yield the title compound. This material was used in step b) without further purification. LCMS (6 minutes method) $[M+H]^+ = 442$ @ Rt 3.48 min. major peak.

b) 2-(2,5-Dichloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.




The procedure for the synthesis of example **21Bb**, 1-(4-Benzyl-morpholin-2-yl)-2-(2,6-dichloro-phenyl)-1-phenyl-ethanol, was followed making non-critical variations to the title compound. ^1H NMR (300MHz, CD_3OD D4) δ : 2.49-2.61 (d, 1H), 2.88-3.11(m, 2H), 3.12-3.24 (m, 1H), 3.24-3.35 (m, 1H), 3.41-3.53 (d, 1H), 3.82-3.96 (m, 1H), 4.04-

4.25 (m, 2H), 6.90-7.00 (m, 1H), 7.02-7.29 (m, 7H). LCMS (12 minutes method)
[M+H]⁺ = 352 @ Rt 4.86 min. major peak


Example 23B: Preparation of 2-(2,5-Difluoro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride

a) 1-(4-Benzyl-morpholin-2-yl)-2-(2,5-difluoro -phenyl)-1-phenyl-ethanol.



The procedure for the synthesis of example 18Ba, 1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-6-fluoro -phenyl)-1-phenyl-ethanol, using 2,5-difluorobenzyl magnesium bromide as starting material (available from Rieke Metals) was followed making non-critical variations, to yield the title compound. This material was used in step b) without further purification. LCMS (6 minutes method) [M+H]⁺ = 410 @ Rt 3.11 min. major peak.

b) 2-(2,5-Difluoro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.




The procedure for the synthesis of example 18Bb, 2-(2-Fluoro-6-chloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride, was followed making non-critical variations, to yield the title compound. ¹H NMR (300MHz, CD₃OD D₄) δ: 2.48-2.59 (d, 1H), 2.87-3.09 (m, 2H), 3.11-3.17 (m, 1H), 3.26-3.38 (m, 2H), 3.81-3.95 (t, 1H), 4.02-

4.11 (d, 1H), 4.13-4.25 (d, 1H), 6.62-6.77 (m, 3H), 7.08-7.28 (m, 5H). LCMS (12 minutes method) $[M+H]^+ = 320$ @ Rt 4.20 min. single major peak.


Example 24B: Preparation of 2-(2-Fluoro-5-phenyl-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride

a) **1-(4-Benzyl-morpholin-2-yl)-2-(2-biphenyl-5-flouro-phenyl)-1-phenyl-ethanol.**



The procedure for the synthesis of example **18Ba**, 1-(4-Benzyl-morpholin-2-yl)-2-(2-chloro-6-fluoro -phenyl)-1-phenyl-ethanol, using 2-phenyl-5-fluorobenzyl magnesium bromide as starting material was followed making non-critical variations, to yield the title compound. This material was used in step b) without further purification. LCMS (6 minutes method) $[M+H]^+ = 468$ @ Rt 3.62 min. major peak.

b) **2-(2-Fluoro-5-phenyl-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride.**




The procedure for the synthesis of example **18Bb**, 2-(2-Fluoro-6-chloro-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol hydrochloride, was followed making non-critical variations to the title compound. ^1H NMR (300MHz, CD_3OD D4) δ : 2.35-2.48 (d, 1H), 2.77-2.91 (t, 1H), 2.91-3.04 (m, 1H), 3.04-3.16 (m, 1H), 3.22-3.28 (m, 1H), 3.30-3.42 (m,

1H), 3.66-3.87 (m, 2H), 4.01-4.14 (d, 1H), 6.70-6.89 (m, 5H), 6.98-7.11 (m, 4H), 7.14-7.25 (m, 4H). LCMS (12 minutes method) $[M+H]^+ = 378$ @ Rt 5.22 min. major peak.

Solid Phase Synthesis of Compounds of Formulae (IB)

- 5 Compounds of the invention wherein Ar₁ is substituted with an aromatic group (i.e., pyridyl, thiophenyl, and optionally substituted phenyl) can be prepared by solid phase synthesis using the route shown below (the black dot represents polystyrene resin).



- 10 The sequence is preferably performed on a polystyrene resin, without characterization of the resin-bound intermediates.

- i) Aliquots (52 mg, 0.05 mmoles) of p-nitrophenyl carbonate resin (Novabiochem) were dispensed into 4.5 ml MiniBlock reaction tubes (Mettler-Toledo). To each resin was added DMF (0.5 ml) followed by a 0.2M solution of 2-(2-bromo-phenyl)-1-morpholin-2-yl-1-phenyl-ethanol in DMF (0.5 ml, 0.1 mmoles). The tubes were sealed and agitated by orbital shaking for 24 hrs. The resins were then filtered and
- 15

washed with DMF (3 x 1.0 ml), a solution of diisopropylethylamine (0.25 ml) in DMF (1.0 ml) and finally DMF (4 x 1.0 ml).

ii) To each resin was added a 2M solution of an optionally substituted aryl boronic acid in DMF (0.5 ml, 1.0 mmoles), a 0.5M solution of triphenylphosphine in DMF (0.2 ml, 0.1 mmoles), a 0.25M solution of Pd(II) acetate in DMF (0.2 ml, 0.05 mmoles) and a 1.25M solution of caesium carbonate in water (0.1 ml, 0.125 mmoles). The tubes were sealed, agitated by orbital shaking and heated at 80° for 20 hrs. The reactions were then cooled to ambient temperature and the resins washed with DMF (2 x 1.0 ml), MeOH (3 x 1.0 ml) and DCM (4 x 1.0 ml).

iii) To each resin was added a TFA/H₂O mixture (95:5 v/v, 1 ml). The tubes were sealed and agitated by orbital shaking for 6 hrs. The reactions were filtered and washed with DCM (2 x 2 ml). Appropriate filtrates and washings were combined and volatile components removed by vacuum evaporation. Each residue was dissolved in MeOH (1 ml) and the solutions applied to MeOH-washed SCX-2 cartridges (0.5 g/3.0 ml) (Jones Chromatography). After draining under gravity the cartridges were washed with MeOH (2.5 ml) and the products then eluted using a 2M solution of ammonia in MeOH (2.5 ml). Removal of volatile components by vacuum evaporation gave the desired products which were purified by preparative LCMS.

By this means were prepared:

Example 25B

2-(4'-methyl-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient) 3.11 min, [M+H]⁺ 374.2

Example 26B

2-(4'-chloro-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient) 3.36 min, [M+H]⁺ 394.2

Example 27B

2-(4'-methoxy-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient)
3.37 min, $[M+H]^+$ 390.2

Example 28B

5 2-(3'-fluoro-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient) 3.39
min, $[M+H]^+$ 378.4

Example 29B

2-(3'-chloro-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient) 3.53
10 min, $[M+H]^+$ 394.4

Example 30B

2-(3'-methoxy-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient)
3.31 min, $[M+H]^+$ 390.4
15

Example 31B

2-(3'-methyl-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient) 3.45
min, $[M+H]^+$ 374.4

20 **Example 32B**

2-(3',5'-dichloro-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient)
3.71 min, $[M+H]^+$ 428.3

Example 33B

25 2-(2',4'-dimethyl-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient)
3.59 min, $[M+H]^+$ 388.4

Example 34B

2-(2',4'-dimethoxy-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min
30 gradient) 3.33 min, $[M+H]^+$ 420.4

Example 35B

1-morpholin-2-yl-1-phenyl-2-(2-pyridin-3-yl-phenyl)-ethanol, RT (6 min gradient) 2.17 min, $[M+H]^+$ 361.4

5 **Example 36B**

1-morpholin-2-yl-1-phenyl-2-(2-thiophen-3-yl-phenyl)-ethanol, 3.25 min, $[M+H]^+$ 366.4

Example 37B

2-(3',4'-dichloro-biphenyl-2-yl)-1-morpholin-2-yl-1-phenyl-ethanol, RT (6 min gradient)
10 3.56 min, $[M+H]^+$ 428.1

The following examples illustrate compounds of of Formulae (IC) above and methods for their preparation.

15 **General Synthetic Procedures for the preparation of Examples 1C-17C**

The numbers included in the following Sections refer to the compounds illustrated in Schemes 2C to 6C herein.

20 **General Procedure 1C: Preparation of racemic N-substituted aryl thiols**

To a solution of 5Ca,5Cb (0.02 g, 0.52 mmol) and the requisite aryl thiol (1.1 eq) in anhydrous dimethylformamide (1 ml) at room temperature under nitrogen was added cesium carbonate (1.1 eq, 0.19 g, 0.57 mmol). The reaction mixture was heated to 95°C for 2 hours. The reaction mixture was allowed to cool to room temperature, diluted with
25 ethyl acetate, then washed sequentially with water, brine, dried over magnesium sulphate and finally concentrated *in vacuo*.

General Procedure 2Ca: Deprotection of N-substituted aryl thiols

To a solution of the requisite N-benzyl aryl thiol in anhydrous dichloromethane
30 (5ml) was added solid supported Hünig's base (Argonaut, 3.56 mmol/g, 2 eq) and α -

chloroethyl chloroformate (3 to 10 eq) at room temperature under nitrogen. The reaction mixture was heated to 40°C and followed by LCMS analysis. After completion the reaction mixture was filtered, and the resin washed with dichloromethane. The combined organic phases were concentrated *in vacuo*. Methanol (HPLC grade, 25 ml) was added and the solution heated to 60°C for 1.5 to 4 hours. After complete consumption of starting material the methanol solution was evaporated to give a solid which was further purified as detailed for individual compounds.

General Procedure 2Cb: Deprotection of N-substituted aryl thiols

To a solution of the requisite *N*-benzyl aryl thiol (1 eq) in ethyl acetate at room temperature was added phenylchloroformate (3 eq). The mixture was warmed under reflux for 2 hours. The mixture was then cooled to room temperature and 30% NaOH with water was added over 1 hour. The biphasic system was stirred for 1.5 hours at room temperature and the organic layer was separated. The organic layer was washed with water, dried over MgSO₄, filtered and rinsed with ethyl acetate.

To the mixture of carbamate and benzylchloride in ethyl acetate was added 5.6M dimethylamine in ethanol. The solution was warmed under reflux (70-72°C) for 2 hours. After cooling at room temperature, water and 12N HCl were added and the mixture was stirred for 10 minutes. The layers were separated and the organic phase was washed twice with water. Then the organic layer was concentrated (T=50°C) until crystallization. MeOH was added and approx. 40% of solvent was then removed under reduce pressure, this operation was repeated. The heterogeneous mixture was stirred for 0.5 hours at room temperature and filtered. The precipitate was washed twice with MeOH and dried under reduce pressure at 40°C to yield the carbamate.

To a biphasic mixture of 30% NaOH and isopropanol warmed to 65°C, was added the carbamate. The heterogeneous mixture was warmed under reflux for 4 hours and then cooled to room temperature and post-agitated overnight. The organic layer was concentrated under reduce pressure and the yellow solid obtained was added to a mixture of AcOEt and 1N NaOH. After separation of the layers, the organic one was washed with 1N NaOH. The aqueous layers were combined and extracted with AcOEt. The combined

organic layers were dried over MgSO_4 , filtered and concentrated under reduce pressure to dryness to obtain the free amine.

General Procedure 3C: Conversion of amines into hydrochloride salts

- 5 To a solution of the requisite amine in dry diethyl ether (1 ml) was added hydrochloric acid (500 μl of a 1M solution in diethyl ether). A white precipitate immediately formed. The suspension was then sonicated for 5 minutes. Ether was blown off with a stream of nitrogen and the samples were dried under high vacuum for several hours to give the hydrochloride salts in near quantitative yield as white solids.

10

General Procedure 4C: Aldoladdition with substituted benzaldehydes

Preparation of 38Ca,38Cb; 39Ca,39Cb; 40Ca,40Cb

- N*-Benzylmorpholinone (1.0 eq) and the requisite aldehyde (1.1 eq) were dissolved in anhydrous tetrahydrofuran (25 ml) under nitrogen and the reaction cooled to -78°C .
15 Then, lithium diisopropylamide (1.1 eq of a 2M solution in heptane/tetrahydrofuran/ethylbenzene) was added over approximately 20 minutes, whilst maintaining the reaction temperature below -78°C . The resulting yellow solution was stirred at -78°C for 1 hour and then allowed to warm to room temperature. The reaction was quenched with saturated ammonium chloride solution (25 ml) and extracted into
20 ethyl acetate. The combined organic layers were dried with magnesium sulphate, filtered and concentrated *in vacuo*, to give a yellow oil which was purified by column chromatography on silica gel (eluent: ethyl acetate/hexane 70/100 [v/v]).

General Procedure 5C: Reduction of substituted aldol adducts


25 **Preparation of 41Ca,41Cb; 42Ca,42Cb; 43Ca,43Cb**

To a solution of the requisite amide 38Ca,38Cb, 39Ca,39Cb or 40Ca,40Cb (1.1 mmol) in anhydrous tetrahydrofuran under nitrogen at room temperature was slowly added borane (4 eq of a 1M solution in tetrahydrofuran). The solution was stirred at 60°C for 2 hours. The reaction was cooled to room temperature; dry methanol (excess) was

slowly added, followed by aqueous hydrochloric acid solution (1M, excess). The reaction mixture was heated to 60°C for 1 hour and quenched with aqueous potassium carbonate solution (1M, excess) and extracted with diethyl ether. The combined organic layers were washed with brine, dried with magnesium sulphate, filtered and concentrated *in vacuo* yielding a yellow oil which was purified by column chromatography on silica gel (eluent: ethyl acetate/hexane 10/100 [v/v]).

Preparation of intermediates for the synthesis of Examples 1C-17C

4-Benzylmorpholin-3-one (2C)




10

A solution of *N*-benzyl-*N*-(2-hydroxyethyl) chloroacetamide (627.7 g, 2.76 mol) in *tert*-butanol (0.9 l) was stirred under nitrogen while warming to 25-30°C. Potassium *tert*-butoxide (2.897 l of a 1M solution in *tert*-butanol, 2.90 mol, 1.05 eq) was added over 2 hours. The reaction mixture was then stirred at room temperature for 90 minutes. Ice-cold water (6 l) was added and the resultant cloudy solution extracted with ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulphate and evaporated *in vacuo* to give a light brown oil (441 g, 84%), which was used in the next stage without further purification; MW 191.23; C₁₁H₁₃NO₂; ¹H NMR (CDCl₃): 7.29-7.40 (5H, m), 4.67 (2H, s), 4.28 (2H, s), 3.87 (2H, t, 5 Hz), 3.31 (2H, t, 5 Hz); LCMS: (12 min method) m/z 192 [M+H]⁺ @ Rt 1.00 min.

20


(2S)-(4-Benzyl-morpholin-2-yl)-phenyl-methanone (3Ca) and (2R)-(4-Benzyl-morpholin-2-yl)-phenyl-methanone (3Cb)

167



Described above under the "Synthesis of Intermediates" section for compounds of Formula (IB).


5 **(S)-Phenyl[(2S)-4-(benzylmethyl)morpholin-2-yl]methanol (4Ca)**



To a stirred solution of [(-)-*B*-chlorodisopinocampheylborane] (45 g, 140 mmol) in dry tetrahydrofuran (300 ml) under nitrogen was added **3Ca** (7.97 g, 28.4 mmol) in one
 10 portion. The reaction mixture was stirred at room temperature for 18 hours. The mixture was evaporated *in vacuo* and extracted from 2M aqueous sodium hydroxide solution into ethyl acetate. The combined organic extracts were washed with brine, dried, filtered and evaporated. The crude product was taken up in chloroform/methanol (1:1 [v/v]) and absorbed onto 150g SCX-2 ion exchange resin. After elution of borane residues with
 15 methanol the product was eluted with 2M ammonia in methanol. Removal of solvent *in vacuo* yielded the product as yellow oil. This was further purified by flash chromatography (eluent: ethyl acetate/isohexane 80/20 [v/v]). After removal of solvents, the product crystallised on standing (6.73g, 84%); MW 283.37; C₁₈H₂₁NO₂; ¹H NMR (CDCl₃): 7.32-7.45 (10H, m), 4.67 (1H, d, 7 Hz), 4.03 (1H, dt, 11 Hz and 2 Hz), 3.86-
 20 3.73 (2H, m), 3.64 (1H, d, 13 Hz), 3.39 (1H, d, 13 Hz), 3.30 (1H, br, s), 2.68 (1H, d, 12 Hz), 2.56 (1H, d, 10 Hz), 2.28-2.15 (2H, m); LCMS: m/z 284 [M+H]⁺ @ Rt 0.95 min.


(2S)-2-[(R)-bromo(phenyl)methyl]-4-(benzylmethyl)morpholine (5Ca)

168



To a solution of **4Ca** (4.71 g, 16.6 mmol) in anhydrous chloroform (200 ml) under nitrogen was added triphenylphosphine dibromide (14.04 g, 33.26 mmol). The reaction mixture was heated at 60°C overnight. The mixture was allowed to cool to room temperature then washed with saturated aqueous sodium carbonate solution, dried over sodium sulphate and concentrated *in vacuo*. The resulting residue was purified by flash chromatography on silica (eluent: ethyl acetate/isohexane gradient 10/90 to 30/70 [v/v]) to give **5Ca** as a white solid (4.63 g, 81%); MW 346.27; C₁₈H₂₀BrNO; ¹H NMR (CDCl₃): 7.14-7.39 (10H, m), 4.83 (1H, d, 7 Hz), 4.01 (1H, br, t, 8 Hz), 3.73 (1H, br, d, 11 Hz), 3.60-3.48 (2H, m), 3.39 (1H, d, 12 Hz), 3.20 (1H, d, 11 Hz), 2.50 (1H, d, 10 Hz), 2.07 (2H, t, 10 Hz); LCMS: (6 min method) m/z 346 [M]⁺ @ Rt 2.51 min.

(2*S*)-2-[(*S*)-Hydroxy(phenyl)methyl]-4-(phenylmethyl)morpholin-3-one (**6Ca**) and (2*S*)-2-[(*R*)-Hydroxy(phenyl)methyl]-4-(phenylmethyl)morpholin-3-one (**6Cb**) and (2*R*)-2-[(*S*)-Hydroxy(phenyl)methyl]-4-(phenylmethyl)morpholin-3-one (**6Cc**) and (2*R*)-2-[(*R*)-Hydroxy(phenyl)methyl]-4-(phenylmethyl)morpholin-3-one (**6Cd**)




To a stirred solution of **2C** (5.02 g, 26 mmol) in anhydrous tetrahydrofuran (25 ml) under nitrogen at -78°C was added lithium diisopropylamide (1.5 eq, 39 mmol, 19.5 ml of a 2M solution in heptane/tetrahydrofuran/ethylbenzene) over approximately 20 minutes, whilst maintaining the reaction temperature below -75°C. The resulting brown solution was stirred for a further 30 minutes at -78°C, before being added over

approximately 30 minutes to a solution of benzaldehyde (1.2 eq, 3.29 g, 31 mmol) in anhydrous tetrahydrofuran (15 ml) under nitrogen at -78°C, whilst again maintaining the reaction temperature below -75°C. The resulting yellow solution was stirred at -78°C for 1 hour, before being allowed to warm to room temperature slowly over 1 hour. The reaction mixture was cautiously quenched by addition of saturated ammonium chloride solution (50 ml) and the tetrahydrofuran was evaporated *in vacuo*. The resulting cloudy aqueous solution was extracted with dichloromethane, and the organic extracts were combined, washed with brine, dried over sodium sulphate and the dichloromethane evaporated *in vacuo* to give a thick brown oil (9.2 g), which partially crystallised on standing. After purification by flash column chromatography (eluent: ethyl acetate/dichloromethane 10/90 to 20/80 gradient [v/v]) **6Ca,6Cb** was obtained as light red crystals (2.46 g, 32%); MW 297.36; C₁₈H₁₉NO₃; ¹H NMR (CDCl₃): 7.36-7.41 (2H, m), 7.16-7.31 (6H, m), 6.86-6.91 (2H, m), 5.14 (1H, d, J 3 Hz), 4.71 (1H, d, 14 Hz), 4.48 (1H, d, J 3 Hz), 4.25 (1H, d, 14 Hz), 4.20 (1H, br, s), 3.89 (1H, ddd, 12 Hz, 3 Hz, 2 Hz), 3.67 (1H, dt, 11 Hz, 3 Hz), 3.16 (1H, dt, 12 Hz and 4 Hz), 2.86 (1H, br, d, 12 Hz); LCMS: m/z 298 [M+H]⁺ @ Rt 1.24 min. **6Cc, 6Cd** was isolated as a brown solid (1.42 g) contaminated with **2C**. Trituration with ethyl acetate afforded pure **6Cc,6Cd** as a white solid (0.484 g, 6%); MW 297.36; C₁₈H₁₉NO₃; ¹H NMR (CDCl₃): 7.55-7.61 (2H, m), 7.36-7.50 (6H, m), 7.25-7.31 (2H, m), 5.21 (1H, d, 2 Hz), 5.09 (1H, d, J 7 Hz and 2 Hz), 4.73 (2H, s), 4.37 (1H, d, J 8 Hz), 4.01 (1H, ddd, 12 Hz, 3 Hz, 2 Hz), 3.77 (1H, dt, 11 Hz, 4 Hz), 3.50 (1H, dt, 12 Hz, 4 Hz), 3.16 (1H, br, d, 12 Hz); LCMS: m/z 298 [M+H]⁺ @ Rt 1.24 min.

(*S*)-Phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methanol (**4Ca**)

and

(*R*)-Phenyl[(2*R*)-4-(phenylmethyl)morpholin-2-yl]methanol (**4Cb**)




To a solution of **6Ca,6Cb** (0.033 g, 1.1 mmol) in anhydrous THF (5 ml) under nitrogen at room temperature was slowly added borane (4 eq, 4.4 ml of a 1M solution in tetrahydrofuran, 4.4 mmol). The solution was stirred at 60°C for 2 hours. After cooling down to room temperature, dry methanol (2 ml) was slowly added to quench excess borane reagent. After addition of aqueous hydrochloric acid solution (2 ml of a 1M solution) the reaction mixture was heated to 60°C for 1 hour. The organic solvents were evaporated *in vacuo* and the concentrated solution was poured onto aqueous potassium carbonate solution (10 ml of a 1M solution) and extracted with diethyl ether (2 x 20 ml). The combined organic layers were washed with brine, water, dried over magnesium sulphate and concentrated *in vacuo*. Purification by flash column chromatography (eluent: hexane/ethyl acetate/triethylamine 90/9/1 [v/v/v]) gave a viscous oil (0.19 g, 60%); MW 283.37; C₁₈H₂₁NO₂; ¹H NMR (CDCl₃): 7.45-7.32 (10H, m), 4.67 (1H, d, 7 Hz), 4.03 (1H, dt, 11 Hz, 2.7 Hz), 3.86-3.73 (2H, m), 3.64 (1H, d, 13 Hz), 3.39 (1H, d, 13 Hz), 3.30 (1H, br, s), 2.68 (1H, d, 13 Hz), 2.56 (1H, d, 11 Hz), 2.28-2.15 (2H, m); LCMS: m/z 284 [M+H]⁺ @ Rt 0.95 min.

(R)-[(2S)-4-Benzylmorpholinyl](phenyl)methanol (4Cc)

and

(S)-[(2R)-4-Benzylmorpholinyl](phenyl)methanol (4Cd)



20


Using the procedure described for the preparation of **4Ca,4Cb** starting from **6Cc,6Cd** (0.14 g, 0.45 mmol) **4Cc,4Cd** was obtained as a viscous oil (0.098 g, 68%); MW 283.37; C₁₈H₂₁NO₂; ¹H NMR (CDCl₃): 7.17-7.28 (10H, m), 4.80 (1H, d, 4 Hz), 3.88 (1H, dt, 11 Hz, 3 Hz), 3.72 (1H, m), 3.61-3.68 (1H, m), 3.50 (1H, d, 13 Hz), 3.25 (1H, d, 13 Hz), 2.52 (2H, br, t, 12 Hz), 2.17 (1H, t, 11 Hz), 2.08 (1H, td, 11 Hz, 3 Hz); LCMS: m/z 284 [M+H]⁺ @ Rt 0.98 min.

25

(2*S*)-2-[(*R*)-Bromo(phenyl)methyl]-4-(phenylmethyl)morpholine (5Ca)

and

(2*R*)-2-[(*S*)-Bromo(phenyl)methyl]-4-(phenylmethyl)morpholine (5Cb)



5 To a solution of **4Ca,4Cb** (10.27 g, 36.29 mmol) in anhydrous dichloromethane (150 ml) under nitrogen at room temperature was added freshly recrystallised triphenylphosphine (13.32 g, 50.80 mmol, 1.4 eq) followed by carbon tetrabromide (16.85 g, 50.8 mmol, 1.4 eq) as a solution in anhydrous dichloromethane (50 ml). After 15 minutes the reaction mixture was diluted with dichloromethane (100 ml) and washed


10 with saturated aqueous solution of sodium hydrogencarbonate, brine, dried over magnesium sulphate and concentrated *in vacuo* to give an orange oil (42.0 g). To the orange oil was added diethyl ether (200 ml) and the resulting suspension was sonicated for 30 minutes. The solvent was decanted and the process repeated with a further portion of diethyl ether. The combined organic extracts were concentrated *in vacuo* to yield an

15 orange solid (22.0 g) which was purified by flash column chromatography (eluent: ethyl acetate/hexane/triethylamine 10/89.5/0.5 [v/v/v]) **5Ca,5Cb** was obtained as a white solid (7.20 g, 57%). Alternative Work-up: The reaction mixture was poured onto a silica (160 g) filtration pad which was washed with dichloromethane (14 x 250 ml). After removal of solvents *in vacuo* and purification by flash column chromatography (eluent: ethyl

20 acetate/hexane/triethylamine gradient 5/94.5/0.5 to 10/89.5/0.5 [v/v/v]) to give a white solid (6.05 g, 48%); MW 346.27; C₁₈H₂₀BrNO; ¹H NMR (CDCl₃): 7.14-7.39 (10H, m), 4.83 (1H, d, 7 Hz), 4.01 (1H, br, t, 8 Hz), 3.73 (1H, br, d, 11 Hz), 3.48-3.60 (2H, m), 3.39 (1H, d, 12 Hz), 3.20 (1H, d, 11 Hz), 2.50 (1H, d, 10 Hz), 2.07 (2H, t, 11 Hz); LCMS: m/z 348/346 [M+H]⁺ @ Rt 1.20 min.

4-[(1*R*)-1-Phenylethyl]morpholine-(2*S*)-carbonitrile (47Ca)

and

4-[(1*R*)-1-Phenylethyl]morpholine-(2*R*)-carbonitrile (47Cb)


- 5 To (*R*)-(-)-2-hydroxyethyl- α -phenethylamine (1.65 g, 10.0 mmol) in diethyl ether (10ml) was added at room temperature 2-chloroacrylonitrile (0.80 ml, 10.0 mmol) with stirring. The mixture was stirred at room temperature for 4.5 days when additional 2-chloroacrylonitrile (0.8 ml, 10.0 mmol) was added. After stirring another 3.5 days, the reaction mixture was concentrated *in vacuo* to give an oil. The oil was dissolved in dry
- 10 tetrahydrofuran (30 ml), cooled under nitrogen to 0°C and potassium *tert*-butoxide (1.23 g, 11.0 mmol) added. The solution was stirred at 0°C for 2 hours then at reflux for 1.5 hours, cooled, diluted with diethyl ether and washed with aqueous saturated sodium bicarbonate. The organic phase was extracted with 2N hydrochloric acid and the aqueous made basic by addition of solid sodium bicarbonate and extracted with diethyl ether. The
- 15 organic phase was dried over magnesium sulphate, filtered and evaporated to a brown oil. The crude product was purified by flash chromatography (eluent: ethyl acetate/hexane gradient 100% ethyl acetate to 50/50 [v/v]) to give **47Ca,47Cb** as a colourless oil (0.58g, 27%); MW 216.29; C₁₃H₁₆N₂O; ¹H NMR (CDCl₃) 7.25-7.38 (5H, m), 4.6 (1H, dd), 4.54 (1H, dd), 3.91-4.06 (2H, m), 3.66-3.82 (2H, m), 3.39-3.49 (2H, m), 2.30-2.89 (4H, m), 1.39 (3H, d). *m/z* [M+H]⁺ 217.
- 20

Phenyl{(2*S*)-4-[(1*R*)-1-phenylethyl]morpholin-2-yl}methanone (48Ca)

and


Phenyl{(2*R*)-4-[(1*R*)-1-phenylethyl]morpholin-2-yl}methanone (48Cb)

173



To a stirred solution of **47Ca,47Cb** (0.57 g, 2.64 mmol) in dry tetrahydrofuran (10 ml) at 0°C under nitrogen was added a solution of phenylmagnesium chloride in tetrahydrofuran (2.0 M, 2.67 ml) dropwise over 2 minutes. The pale yellow solution was stirred at 0°C for 30 minutes and then allowed to warm to room temperature. After 2 hours the mixture was cooled, quenched with 2M hydrochloric acid and was stirred vigorously for 1 hour at room temperature. After addition of water and extraction with ethyl acetate, the combined organic layers were washed with brine, dried over magnesium sulphate, filtered and evaporated to give an oil (0.63 g). After purification by column chromatography (eluent: ethyl acetate/hexane gradient 0/100 to 20/80 [v/v]) **48Ca** was obtained as an oil (0.15 g, 19%); MW 295.38; C₁₉H₂₁NO₂; ¹H NMR (CDCl₃) 8.00 (2H, d), 7.60 (1H, t), 7.50 (2H, t), 7.20-7.35 (5H, m), 4.96 (1H, d), 3.93-4.00 (1H, m), 3.70-3.80 (1H, m), 3.41 (1H, q), 3.25 (1H, br, d), 2.59 (1H, br, d), 2.13 -2.36 (2H, m), 1.38 (3H, d). *m/z* [M+H]⁺ 296 followed by **48Cb** as an oil (0.27 g, 35%); ¹H NMR (CDCl₃) 7.90 (2H, d), 7.54 (1H, t), 7.45 (2H, t), 7.20-7.38 (5H, m), 4.85 (1H, d), 4.05-4.12 (1H, m), 3.80-3.92 (1H, m), 3.43 (1H, q), 2.86-3.00 (2H, m), 2.29-2.40 (1H, m), 2.21 (1H, t), 1.38 (3H, d). *m/z* [M+H]⁺ 296.


(R)-Phenyl{(2S)-4-[(1R)-1-phenylethyl]morpholin-2-yl}methanol (50C)



To a stirred solution of **48Ca** (0.08 g, 0.26 mmol) and triphenylsilane (0.34 g, 1.31 mmol) in dichloromethane (4 ml) cooled to 0°C was added boron trifluoride etherate

(0.09 g, 0.66 mmol) followed by trifluoroacetic acid (0.36 ml, 63 mmol). The reaction mixture was allowed to warm to room temperature and diluted after three hours with dichloromethane (20 ml) and neutralised with aqueous sodium bicarbonate. The organic phase was dried over magnesium sulphate, filtered and evaporated to give the required product. This was purified as its hydrochloric acid salt crystallising from isopropanol and diethyl ether (0.05 g, 69%); MW 297.4; C₁₉H₂₃NO₂; ¹H NMR (CDCl₃) on free base 7.08-7.29 (10H, m), 4.78 (1H, d), 3.90-4.00 (1H, m), 3.57-3.68 (2H, m), 3.33 (1H, q), 2.53-2.64 (1H, m), 2.37-2.47 (1H, m), 2.09-2.26 (2H, m), 1.29 (3H, d). *m/z* [M+H]⁺ 298.


10 **(R)-Phenyl{(2S)-4-[(1R)-1-phenylethyl]morpholin-2-yl}methyl methanesulphonate (51C)**



To a solution of **50C** (0.05 g, 0.17 mmol) in dichloromethane (1 ml) at room temperature was added polymer supported Hünig's base ((Argonaut, 3.56 mmol/g, 0.089 g, 0.32 mmol, 1.9 eq) and methanesulphonyl chloride (0.02 g, 0.19 mmol). The mixture was stirred under nitrogen for 6 hours then filtered and concentrated *in vacuo*. The crude product was purified by flash column chromatography (eluent: ethyl acetate/heptane 33/67 [v/v]) to give **51C** as a colourless oil (0.035 g, 55%); MW 375.49; C₂₀H₂₅NO₄S ¹H NMR (CDCl₃) 7.20-7.35 (10H, m), 5.46 (1H, d), 3.79-3.88 (2H, m), 3.59 (1H, td), 3.4 (1H, q), 2.68-2.78 (2H, m), 2.68 (3H, s), 2.03-2.24 (2H, m), 1.34 (3H, d). *m/z* [M+H]⁺ 376.

(2S)-4-[(1R)-1-Phenylethyl]-2-((S)-phenyl{2-(trifluoromethyl)phenyl}thio}methyl)morpholine (52C)


175



A mixture of **51C** (0.035 g, 0.093 mmol), potassium carbonate (0.026 g, 0.19 mmol) and 2-trifluoromethylbenzenethiol (0.084 g, 0.47 mmol) in dry, degassed dimethylformamide (0.5 ml) was stirred under nitrogen at room temperature for 3 days. The reaction mixture was diluted with water and extracted with diethyl ether. The extracts was washed with water and brine, dried over magnesium sulphate, filtered and evaporated to give a colourless oil (0.03 g, 71%). Purification by flash column chromatography (eluent: ethyl acetate/heptane 20/80 [v/v]) gave **52C** as a colourless oil (0.03 g, 71%); MW 457.56; C₂₆H₂₆F₃NOS ¹H NMR (CDCl₃) 7.53 (1H, d), 7.10-7.28 (13H, m), 4.39 (1H, d), 3.85-4.04 (2H, m), 3.8 (1H, td), 3.35 (1H, q), 2.70 (1H, d), 2.40 (1H, d), 2.30 (1H, td), 2.10-2.20 (1H, m), 1.29 (3H, d). *m/z* [M+H]⁺ 458.

Example 1C: (2S)-2-((S)-Phenyl){2-(trifluoromethyl)phenyl}thio}methyl)morpholine (9C)


15 **(S)-Phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methyl 2-trifluoromethyl)phenyl sulfide (8C)**



Compound **8C** was obtained from **5Ca** (4.00 g, 11.55 mmol), 2-trifluoromethyl thiophenol (2.47 g, 13.86 mmol, 1.2 eq) and caesium carbonate (4.95 g, 15.24 mmol, 1.1 eq) in dimethylformamide (60 ml) as a brown oil following a modification of **General Procedure 1C** in which the reaction was carried out over 1 hour (6.04 g). The oil was purified by flash column chromatography (eluent: hexane/ethyl acetate gradient 100 to

90/10 [v/v]) to give a yellow oil (4.83 g, 94%); MW 443.54; C₂₅H₂₄F₃NOS; ¹H NMR (CDCl₃): 7.60 (1H, dd, 7 Hz, 1 Hz), 7.17-7.39 (13H, m), 4.50 (1H, d, 7 Hz), 3.97-4.12 (2H, m), 3.73 (1H, dt, 10 Hz, 2 Hz), 3.59 (1H, d, 13 Hz), 3.37 (1H, d, 13 Hz), 2.57-2.68 (2H, m); 2.18-2.38 (2H, m); LCMS (2.5 minute method): m/z 445 [M+H]⁺ @ Rt 1.50 min.

(2S)-2-((S)-Phenyl{[2-(trifluoromethyl)phenyl]thio}methyl)morpholine (9C)



Compound **9C** (**Example 1C**) was obtained from **8C** (5.25 g, 11.84 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 6.64 g, 23.67 mmol, 2 eq) and α-chloroethyl chloroformate (3.83 ml, 35.51 mmol, 3 eq) in anhydrous dichloromethane (75 ml) following **General Procedure 2Ca**. After evaporation of solvents a light brown solid (5.60 g) was obtained which was recrystallised from iso-propanol. The solid was suspended in ethyl acetate and washed with an aqueous solution of sodium hydroxide (50 ml of a 1M solution). The organic layer was washed with brine, dried over magnesium sulphate and concentrated *in vacuo* to yield the free amine as a colourless oil (3.10 g, 74%); MW 353.41; C₁₈H₁₈F₃NOS; ¹H NMR (CDCl₃): 7.46 (1H, d, 8 Hz), 7.24 (1H, d, 7 Hz), 7.05-7.2 (7H, m), 4.28 (1H, d, 8 Hz), 3.92 (1H, d, 11 Hz), 3.80 (1H, q, 7 Hz), 3.58 (1H, dt, 2 Hz and 11 Hz), 2.69-2.87 (2H, m), 2.59 (2H, d, 6 Hz), 2.13-1.90 (1H, br s); LCMS (10 minute method): m/z 354 [M+H]⁺ @ Rt 5.26 min. The hydrochloride salt of **9** was obtained following **General Procedure 3C**.


An alternative method for the preparation of compound **9C** (**Example 1C**), according to **Scheme 6C**, is as follows:

To a suspension of polymer supported Hünig's base (0.11 g, 0.40 mmol) and **52C** (0.03 g, 0.066 mmol) in dry dichloromethane (1 ml) was added α-chloroethyl

chloroformate (0.09 g, 0.066 mmol) at room temperature under nitrogen. The mixture was stirred at room temperature over the weekend then filtered and concentrated *in vacuo*. This was taken up in methanol, heated at 70°C for 2 hours, cooled, and purified by SCX chromatography (eluent: ammonia/methanol 1/1 [v/v]) to give **9C** as a colourless oil (0.01 g, 43%). The spectroscopic data for **9C** obtained by the route outlined here was identical to the data for **9C** obtained as described above.


Example 2C: (2S)-2-((S)-Phenyl{2-(thiomethyl)phenyl}thio)methyl morpholine (11C)

(2S)-2-[(S)-{2-(methylthio)phenyl}thio](phenyl)methyl]-4-(phenylmethyl)morpholine (10C)



Compound **10C** was obtained from **5Ca** (4.0 g, 11.55 mmol), 2-methylsulphenylthiophenol (2.17 g, 13.86 mmol, 1.2 eq) and caesium carbonate (4.42 g, 13.63 mmol, 1.18 eq) in dimethylformamide (35 ml) following a modification of **General Procedure 1C** in which the mixture was heated at 50°C for 1.5 hours, allowed to cool to room temperature, taken up in methanol and treated with SCX-2 (100 g). The SCX-2 was washed with methanol. **10C** was obtained as a white solid (4.92 g) after SCX chromatography (eluent: ammonia/methanol 1/1 [v/v]) and removal of solvents *in vacuo*. Purification by flash column chromatography (eluent: ethyl acetate/isohexane gradient 10/90 to 30/70 [v/v]) gave **10C** as a white solid (4.04 g, 83%); MW 421.63; C₂₇H₂₇NOS₂; ¹H NMR (CDCl₃): 7.03-7.15 (6H, m), 6.93-6.99 (2H, m), 6.74 (1H, td, 7 Hz, 1 Hz), 4.31 (1H, d, 8 Hz), 3.95 (1H, br, d, 12 Hz), 3.83 (1H, td, 8 Hz, 3.8 Hz), 3.59 (1H, td, 11 Hz and 3 Hz), 2.82 (1H, td, 12 Hz and Hz), 2.61-2.75 (3H, m), 2.35 (3H, s), 1.73 (1H, br, s); LCMS (6 minute method): m/z 422 [M+H]⁺ @ Rt 3.36 min.

178


(2S)-2-((S)-Phenyl{[2-(trifluoromethyl)phenyl]thio}methyl)morpholine (11C)

Compound **11C** (**Example 2C**) was obtained from **10C** (4.02 g, 9.53 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 5.02 g, 17.87 mmol, 2 eq) and α -chloroethyl chloroformate (3.09 ml, 28.6 mmol, 3 eq) in anhydrous dichloromethane (75 ml) following **General Procedure 2Ca**. The mixture was heated at 40°C for 1.5 hours then left to stir at room temperature overnight. The reaction mixture was filtered and concentrated *in vacuo* to give a pale orange liquid. This was taken up in methanol (70 ml) and heated at 40°C for 2 hours. A white solid crashed out of the solution which was taken up in methanol and purified by SCX chromatography (eluent: ammonia/methanol 1/1 [v/v]). After evaporation *in vacuo* **11C** was obtained as a pale yellow oil (3.13 g, 99%); MW 331.50; C₁₈H₂₁NOS₂; ¹H NMR (CDCl₃): 7.03-7.15 (6H, m), 6.93-6.99 (2H, m), 6.74 (1H, td, 7 Hz, 2 Hz), 4.31 (1H, d, 8 Hz), 3.95 (1H, br, d, 12 Hz), 3.83 (1H, td, 8 Hz, 4 Hz), 3.59 (1H, td, 11 Hz, 3 Hz), 2.82 (1H, td, 12 Hz, 3 Hz), 2.61-2.75 (3H, m), 2.35 (3H, s), 1.73 (1H, br, s). Compound **11C** was converted into its hydrochloride salt following a modification of **General Procedure 3C** in which the pale yellow oil was taken up in isopropanol (~200 ml) and filtered. Addition of hydrogen chloride (19 ml of a 1M solution in diethyl ether, 19 mmol) gave a white precipitate to which further diethyl ether (~50 ml) was added. The solid was isolated by filtration and washed with diethyl ether to give the hydrochloride salt of **11C** as a white solid (3.03 g, 78%); MW 367.96; C₁₈H₂₂ClNOS₂; ¹H NMR (CDCl₃): 9.94 (2H, br, s), 7.06-7.18 (6H, m), 6.94-7.03 (2H, m), 6.78 (1H, t, 7 Hz), 4.24-4.32 (1H, m), 4.20 (1H, d, 6 Hz), 3.89-4.06 (2H, m), 3.18 (2H, br, t, 12 Hz), 2.99 (2H, br, s), 2.37 (3H, s); LCMS (10 minute method): m/z 332 [M-HCl]⁺ @ Rt 5.07 min.

Example 3C: (2S)-2-[(S)-{2-(1-methylethyl)phenyl}thio}(phenyl)methylmorpholine (13C)

(2S)-2-[(S)-{2-(1-methylethyl)phenyl}thio}(phenyl)methyl]-4-


5 **(phenylmethyl)morpholine (12C)**




Compound **12C** was obtained from **5Ca** (4.04 g, 11.66 mmol), 2-isopropylsulphenyl-thiophenol (2.35 ml, 14 mmol, 1.2 eq) and caesium carbonate (4.56 g, 14 mmol, 1.2 eq) in dimethylformamide (35 ml) following a modification of **General**
10 **Procedure 1C** in which the mixture was heated at 90°C for 20 minutes, allowed to cool to room temperature, taken up in ethyl acetate (50 ml), washed with water and brine, dried over sodium sulphate, filtered and reduced *in vacuo* to give a yellow oil which was purified by SCX chromatography (eluent: ammonia/methanol 1/1 [v/v]). Removal of solvents *in vacuo* gave **12C** as a white solid (4.45, 91%); MW 417.62; C₂₇H₃₁NOS; ¹H
15 NMR (CDCl₃): 7.14-7.26 (7H, m), 7.03-7.1 (6H, m), 6.86-6.92 (1H, m), 4.10 (1H, d, 8 Hz), 3.88-3.94 (2H, m), 3.62 (1H, td, 11 Hz, 2 Hz), 3.37-3.47 (2H, m), 3.22 (1H, d, 13 Hz), 2.50 (2H, d, 11 Hz), 2.12-2.29 (2H, m), 1.05 (3H, d, 7 Hz), 0.92 (3H, d, 7 Hz); LCMS (6 minute method): m/z 418 [M+H]⁺ @ Rt 3.72 min.

20 **(2S)-2-[(S)-{2-(1-methylethyl)phenyl}thio}(phenyl)methyl]morpholine (13C)**

180




Compound **13C** (**Example 3C**) was obtained from **12C** (4.44 g, 10.65 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 6.05 g, 21.54 mmol, 2 eq) and α -chloroethyl chloroformate (3.30 ml, 32.0 mmol, 3 eq) in anhydrous dichloromethane (50 ml) following **General Procedure 2Ca**. The mixture was heated at 40°C for 1.5 hours then left to stir at room temperature overnight. The reaction mixture was filtered and concentrated *in vacuo* to give a pale yellow liquid. This was taken up in methanol (50 ml) and heated at 60°C for 1.5 hours. The reaction mixture was allowed to cool to room temperature and purified by SCX chromatography (eluent: ammonia/methanol 1/1 [v/v]) to give **13C** as a pale yellow oil; MW 327.49; C₂₀H₂₅NOS; ¹H NMR (CDCl₃): 7.22 (1H, d, 8 Hz), 7.03-7.13 (7H, m), 6.87-6.92 (1H, m), 4.04 (1H, d, 8 Hz), 3.94-3.99 (1H, m), 3.79 (1H, td, 9 Hz, 3 Hz), 3.61 (1H, td, 11 Hz, 3 Hz), 3.41 (1H, sept., 7 Hz), 2.82 (1H, td, 12 Hz and 3 Hz), 2.72 (1H, br, d, 12 Hz), 2.52-2.63 (2H, m), 1.70 (1H, br, s), 1.05 (3H, d, 7 Hz), 0.91 (3H, d, 7 Hz). Compound **13C** was converted into its hydrochloride salt following a modification of **General Procedure 3C** in which the pale yellow oil was taken up in ether (50 ml), and filtered. Addition of hydrogen chloride in dry diethyl ether (19 ml of a 1M solution in diethyl ether) gave a white precipitate to which further diethyl ether (50 ml) was added. The reaction mixture was concentrated and the residue washed with diethyl ether to give a white solid (2.76 g, 69% overall yield from **5Ca**); MW 363.95; C₂₀H₂₅NOS.HCl; ¹H NMR (CDCl₃): 9.91 (2H, br, s), 7.05-7.22 (7H, m), 6.91-6.96 (2H, m), 4.23-4.31 (1H, m), 4.08-3.90 (3H, m), 3.31-3.41 (1H, m), 3.04-3.21 (2H, br, m), 2.91-2.99 (2H, br, m), 1.06 (3H, d, 7 Hz), 0.93 (3H, d, 7 Hz); LCMS (10 minute method): m/z 327 [M-HCl]⁺ @ Rt 5.7 min.

Example 4C: (2S)-2-[(S)-([1,1'-Biphenyl]-2-ylthio)(phenyl)methyl]morpholine (15C)**(2S)-2-[(S)-([1,1'-Biphenyl]-2-ylthio)(phenyl)methyl]-4-(phenylmethyl)morpholine (14C)**

5

Compound **14C** was obtained from **5Ca** (2.16 g, 6.24 mmol), 2-phenylsulphenylthiophenol (2.35 ml, 14 mmol, 1.2 eq) and caesium carbonate (2.43 g, 7.5 mmol, 1.2 eq) in dimethylformamide (50 ml) following a modification of **General Procedure 1C** in which the mixture was heated at 90°C for 20 minutes, allowed to cool to room temperature, taken up in ethyl acetate (50 ml), washed with water and brine, dried over sodium sulphate, filtered and reduced *in vacuo* to give a yellow oil. Purification by SCX-chromatography (eluent: ammonia/methanol 1/1 [v/v]) followed by evaporation *in vacuo* gave **14C** as a white solid (0.59 g, 90%); MW 451.64; C₃₀H₂₉NOS; ¹H NMR (CDCl₃): 6.93-7.34 (19H, m), 3.92 (1H, br, d, 6 Hz), 3.63-3.76 (2H, m), 3.45 (1H, t, 10 Hz), 3.33 (1H, d, 13 Hz), 3.17 (1H, d, 12 Hz), 2.39 (1H, d, 12 Hz), 2.20 (1H, d, 11 Hz), 1.97-2.07 (1H, m), 1.82-1.92 (1H, m); LCMS (6 minute method): m/z 452 [M+H]⁺ @ Rt 3.69 min.

15

(2S)-2-[(S)-([1,1'-Biphenyl]-2-ylthio)(phenyl)methyl]morpholine (15C)

Compound **15C** (**Example 4C**) was obtained from **14C** (2.95 g, 6.54 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 13.06 g, 21.54 mmol, 2 eq) and α -chloroethyl chloroformate (2.0 ml, 19.6 mmol, 3 eq) in anhydrous dichloromethane (50 ml) following **General Procedure 2Ca**. The reaction mixture was concentrated *in vacuo* to give a pale yellow liquid. This was taken up in methanol (70 ml) and heated at 40°C for 2 hours. A white solid crashed out of the solution which was taken up in methanol and purified by SCX-chromatography (eluent: ammonia/methanol 1/1 [v/v]). After removal of solvents *in vacuo* **15C** was obtained as a pale yellow oil; MW 361.51; C₂₃H₂₃NOS; ¹H NMR (CDCl₃): 7.0-7.45 (14H, m), 3.95 (1H, d, 8 Hz), 3.65-3.85 (2H, m), 3.35 (1H, d, 12 Hz), 3.2 (1H, d, 12 Hz), 2.45 (1H, d, 10 Hz), 2.20 (1H, d, 10 Hz), 2.0-2.15 (1H, m), 1.8-2.0 (1H, m); LCMS (12 minute method): m/z 363 [M+H]⁺ @ Rt 3.00 min. **15C** was converted into its hydrochloride salt following a modification of **General Procedure 3C** in which the pale yellow oil was taken up in isopropanol (~200 ml), and filtered. Addition of hydrogen chloride (19 ml of a 1M solution in diethyl ether) gave a white precipitate to which further diethyl ether (~50 ml) was added. The solid was isolated by filtration and washed with diethyl ether to give the hydrochloride salt of **15C** as a white solid (1.95 g, 75% overall yield from **5Ca**); MW 397.97; C₂₃H₂₃NOS.HCl; ¹H NMR (CDCl₃): 9.80 (2H, br, s), 7.38-7.03 (12H, m), 6.90-6.96 (2H, m), 3.85-4.00 (2H, m), 3.72-3.82 (1H, m), 3.66 (1H, d, 5 Hz), 2.98-3.10 (1H, m), 2.81 (1H, br, s), 2.62 (2H, br, s); LCMS (12 minute method): m/z 362 [M+H]⁺ @ Rt 2.99 min.


Example 5C: (2S)-2-[(S)-[(2-Fluorophenyl)thio](phenyl)methyl]morpholine (17C)

(2S)-2-[(S)-[(2-Fluorophenyl)thio](phenyl)methyl]-4-phenylmethyl)morpholine (16Ca)

25 and


(2R)-2-[(R)-[(2-Fluorophenyl)thio](phenyl)methyl]-4-phenylmethyl)morpholine (16Cb)

183



Compounds **16Ca,16Cb** were obtained from **5Ca,5Cb** (0.114 g, 0.33 mmol), 2-fluorothiophenol (0.045 g, 0.36 mmol, 1.2 eq) and caesium carbonate (0.12 g, 0.36 mmol, 1.2 eq) in dimethylformamide (50 ml) following **General Procedure 1C** as a pale yellow oil (0.14 g, 65%); MW 393.53; C₂₄H₂₄FNOS; ¹H NMR (CDCl₃): 7.12-7.36 (12H, m), 6.87-6.99 (2H, m), 4.48 (1H, d, 8 Hz), 4.00-4.11 (2H, m), 3.77 (1H, td, 11 Hz, 2 Hz), 3.60 (1H, d, 13 Hz), 3.37 (1H, d, 13 Hz); 2.63 (2H, t, 10 Hz), 2.16-2.31 (2H, m); LCMS (2.5 minute method): *m/z* 394 [M+H]⁺ @ R_t 1.41 min.

10 **(2S)-2-[(S)-[(2-Fluorophenyl)thio](phenyl)methyl]morpholine (17C)**



Compound **17C (Example 5C)** was obtained from **16Ca,16Cb** (0.72 g, 0.18 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 2.0 g, 0.56 mmol, 3 eq) and α-chloroethyl chloroformate (0.62 ml, 0.56 mmol, 3 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a viscous yellow oil (0.046 g, 82%) from which **17C** was obtained as a single isomer after separation by chiral HPLC (0.016 g); Chiral LC (AD): 10.83 min. LC purity = 91% (UV254nm) / 98% (ELS); LCMS (10 minute method): *m/z* 304 [M+H]⁺ @ R_t 5.82 min; HPLC purity = 84% (UV215nm) / 98% (ELS); MW 303.41; C₁₇H₁₈FNOS; ¹H NMR (CDCl₃): 7.13-7.00 (7H, m), 6.87-6.76 (2H, m), 4.29 (1H, d, 9 Hz), 3.98-3.93, (1H, m), 3.78 (1H, td, 9 Hz and 4 Hz), 3.60 (1H, td, 11 Hz and 3 Hz), 2.82 (1H, td, 12 Hz, 3 Hz), 2.76-2.70 (1H, m), 2.57-

2.53, (2H, m), NH signal not observed; LCMS (10 minute method): m/z 304 $[M+H]^+$ @ Rt 5.84 min; HPLC purity = 100%% (ELS). Compound **17C** was converted into its hydrochloride salt following **General Procedure 3C**.


5 **Example 6C: (2S)-2-[(S)-[(2-Ethylphenyl)thio](phenyl)methyl]morpholine (19C)**

(2S)-2-[(S)-[(2-Ethylphenyl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine (18Ca)

and

(2R)-2-[(R)-[(2-Ethylphenyl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine


10 **(18Cb)**



Compounds **18Ca,18Cb** were obtained from **5Ca,5Cb** (0.2 g, 0.58 mmol), 2-ethyl-thiophenol (0.16 g, 1.16 mmol, 2 eq) and caesium carbonate (0.23 g, 0.7 mmol, 1.2 eq) in dimethylformamide (5 ml) following modification of **General Procedure 1C** in which the reaction mixture was heated to 95°C for 2 hours. After purification by flash column chromatography (eluent: ethyl acetate/hexane 9/1 [v/v]) **18Ca,18Cb** was obtained as a white solid (0.15 g, 65%%); MW 403.59; $C_{26}H_{29}NOS$; 1H NMR ($CDCl_3$): 6.96-7.40 (14H, m), 4.22 (1H, d, 7 Hz), 3.96-4.01 (2H, m), 3.72 (1H, td, 11 Hz and 2 Hz), 3.52 (1H, d, 13 Hz), 3.32 (1H, d, 13 Hz), 2.68 (2H, q, 8 Hz), 2.59 (2H, br d, 12 Hz), 2.06-2.21 (2H, m), 1.12 (3H, t, 7 Hz); LCMS (2.5 minute method) m/z 404 $[M+H]^+$ @ Rt 1.49 min.

(2S)-2-[(S)-[(2-Ethylphenyl)thio](phenyl)methyl]morpholine (19C)

185



Compound **19C** (**Example 6C**) was obtained from **18Ca,18Cb** (0.18 g, 0.52 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 3.7 g, 1.04 mmol, 2 eq) and α -chloroethyl chloroformate (0.34 ml, 3.12 mmol, 3 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a viscous yellow oil (0.21 g, 86%) from which **19C** was obtained after separation by chiral HPLC on chiral OD semi-preparative column; chiral LC (OD): 15.95 min. LC purity = 100% (UV254nm) / 100% (ELS); MW 313.47; $C_{19}H_{23}NOS$; 1H NMR ($CDCl_3$): 7.17 (1H, d, 8 Hz), 7.12-7.05 (5H, m), 7.01 (2H, d, 4 Hz), 6.87-6.93 (1H, m), 4.07 (1H, d, 8 Hz), 3.92-3.97 (1H, m), 3.74-3.80 (1H, m), 3.59 (1H, td, 11 Hz, 3 Hz), 2.80 (1H, td, 12 Hz and 3 Hz), 2.71 (1H, br, d, 12 Hz), 2.63-2.54 (4H, m), 1.64 (1H, br, s), 1.04 (3H, t, 8 Hz); LCMS (10 minute method): m/z 314 $[M+H]^+$ @ Rt 5.92 min. **19C** was converted into its hydrochloride salt following **General Procedure 3C**; MW 349.93; $C_{19}H_{23}NOS.HCl$; 1H NMR ($CDCl_3$): 10.10 (2H, br, s), 7.13-7.28 (8H, m), 7.02-7.08 (1H, m), 4.36 (1H, br, s), 4.01-4.17 (3H, br, m), 3.16-3.31 (2H, br, m), 2.92-3.09 (2H, br, m), 2.71 (2H, q, 8 Hz), 1.15 (3H, t, 7 Hz).


Example 7C: (2S)-2-[(S)-{2-(Methyloxy)phenyl}thio}(phenyl)methyl]morpholine (21C)

(2S)-2-[(S)-{2-(Methyloxy)phenyl}thio}(phenyl)methyl]-4-(phenylmethyl)morpholine (20Ca)

and


(2R)-2-[(R)-{2-(Methyloxy)phenyl}thio}(phenyl)methyl]-4-(phenylmethyl)morpholine (20Cb)

186



Compounds **20Ca,20Cb** were obtained from **5Ca,5Cb** (0.18 g, 0.52 mmol), 2-methoxy thiophenol (0.074 ml, 0.57 mmol, 1.2 eq) and caesium carbonate (0.17 g, 0.52 mmol, 1.2 eq) in dimethylformamide (5 ml) following modification of **General**
 5 **Procedure 1C** in which the reaction was heated at 95°C for 2.5 hours. After purification by flash column chromatography (eluent: ethyl acetate/hexane gradient 15/85 to 25/75 [v/v]) **20Ca,20Cb** was obtained as a viscous yellow oil (0.17 g, 83%); MW 405.56; C₂₅H₂₇NO₂S; ¹H NMR (CDCl₃): 7.01-7.26 (12H, m), 6.58-6.63 (2H, m), 4.39 (1H, d, 7 Hz), 3.86-3.91 (2H, m), 3.71 (3H, s), 3.56-3.62 (1H, m), 3.42 (1H, d, 11 Hz); 3.21 (1H, d,
 10 11 Hz), 2.46-2.52 (2H, m), 2.01-2.11 (2H, m); LCMS (10 minute method): *m/z* 406 [M+H]⁺ @ R_T 6.09 min.

(2S)-2-[(S)-{2-(Methoxy)phenyl}thio](phenyl)methylmorpholine (21C)



15 Compound **21C (Example 7C)** was obtained from **20Ca,20Cb** (0.1 g, 0.25 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 1.78 g, 0.5 mmol, 2 eq) and α-chloroethyl chloroformate (0.16 ml, 1.5 mmol, 3 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a viscous yellow oil (0.06 g, 77%) from which **21C** was obtained after separation by chiral HPLC on a Chiralcel OJ semi-preparative column. Chiral LC: 11.45 min. LC purity = 100%; MW 315.44; C₁₈H₂₁NO₂S;
 20

¹H NMR (CDCl₃): 7.14-7.34 (7H, m), 6.74-6.84 (2H, m), 4.50 (1H, d, 8 Hz), 4.10 (1H, d, 11 Hz), 3.85-4.00 (4H, m), 3.74 (1H, dt, 1 Hz, 11 Hz), 2.82-3.02 (2H, m), 2.66-3.02 (3H, m); LCMS (10 minute method): *m/z* 316 [M+H]⁺ @ R_t 4.87 min. **21C** was converted its hydrochloride salt following **General Procedure 3C**.

5


Example 8C: (2*S*)-2-[(*S*)-({2-[(1-

Methylethyl)oxy]phenyl}thio)(phenyl)methyl]morpholine (23C**)**

(2*S*)-2-[(*S*)-({2-[(1-Methylethyl)oxy]phenyl}thio)(phenyl)methyl]-4-(phenylmethyl)morpholine (22Ca**)**

10 and

(2*R*)-2-[(*R*)-({2-[(1-Methylethyl)oxy]phenyl}thio)(phenyl)methyl]-4-(phenylmethyl)morpholine (22Cb**)**




Compounds **22Ca,22Cb** were obtained from **5Ca,5Cb** (0.57 g, 1.7 mmol), 2-isopropoxy-thiophenol (0.94 g, 5.61 mmol) and caesium carbonate (2.18 g, 6.72 mmol, 1.2 eq) in dimethylformamide (15 ml) following modification of **General Procedure 1C** in which the reaction mixture was heated to 95°C for 3 hours. After purification by SCX chromatography (eluent: ammonia/methanol 1/1 [v/v]) **22Ca,22Cb** was obtained as a dark yellow oil (0.56 g, 76%%); MW 433.62; C₂₇H₃₁NO₂S; ¹H NMR (CDCl₃): 7.01-7.24 (7H, m), 6.94-7.09 (5H, m), 6.64 (1H, d, 8 Hz), 6.56 (1H, td, 8 Hz, 1 Hz), 4.42-4.51 (2H, m), 3.83-3.92 (2H, m), 3.56 (1H, td, 11 Hz and 3 Hz), 3.42 (1H, d, 13 Hz), 3.24 (1H, d, 13 Hz), 2.52 (1H, d, 11 Hz), 2.46 (1H, d, 11 Hz), 2.05-2.17 (2H, m), 1.29 (3H, d, 6 Hz), 1.27 (3H, d, 6 Hz); LCMS (2.5 minute method): *m/z* 434 [M+H]⁺ @ R_T 1.44 min.

(2*S*)-2-[(*S*)-({2-[(1-Methylethyl)oxy]phenyl}thio)(phenyl)methyl]morpholine (23C**)**

25

188




Compound **23C** (**Example 8C**) was obtained from **22Ca,22Cb** (0.56 g, 1.3 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.73 g, 2.6 mmol, 2 eq) and α -chloroethyl chloroformate (0.16 ml, 1.5 mmol, 3 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a viscous yellow oil (0.41 g, 93%) after separation using chiral HPLC on a OD semi-preparative column. Chiral LC (OD): 12.51 min. LC purity = 100% (UV254nm) / 100% (ELS); MW 343.49; $C_{20}H_{25}NO_2S$; 1H NMR ($CDCl_3$): 7.13-7.20 (1H, m), 6.96-7.12 (6H, m), 6.67 (1H, d, 8 Hz), 6.59 (1H, td, 7 Hz, 1 Hz), 4.48 (1H, sept., 6 Hz), 4.38 (1H, d, 7 Hz), 3.90-3.95 (1H, m), 3.73 (1H, td, 8 Hz, 4 Hz), 3.54 (1H, td, 11 Hz and 3 Hz), 2.79 (1H, td, 12 Hz and 3 Hz), 2.62-2.72 (3H, m), 1.55 (1H, br, s), 1.32 (3H, d, 6 Hz), 1.29 (3H, d, 6 Hz); LCMS (10 minute method): m/z 344 $[M+H]^+$ @ Rt 6.19 min; HPLC purity = 92% (UV215nm). **23C** was converted into its hydrochloride salt following **General Procedure 3C**; MW 379.95; $C_{20}H_{25}NO_2S.HCl$; 1H NMR ($CDCl_3$): 9.81-10.04 (2H, br, m), 7.03-7.25 (7H, m), 6.71 (1H, d, 8 Hz), 6.63 (1H, t, 7 Hz), 4.51 (1H, sept., 6 Hz), 4.31 (1H, d, 6 Hz), 4.15-4.23 (1H, m), 3.83-4.03 (2H, m), 3.05-3.18 (2H, m), 2.80-3.03 (2H, m), 1.31 (3H, d, 6 Hz), 1.29 (3H, d, 6 Hz).

Example 9C: 2-[(S)-(2S)-Morpholin-2-yl(phenyl)methyl]thio}phenyl trifluoromethyl ether (25C)


(2S)-4-(Phenylmethyl)-2-[(S)-phenyl(2-[(trifluoromethyl)oxy]phenyl)thio)methyl]morpholine (24Ca)
and
(2S)-4-(Phenylmethyl)-2-[(S)-phenyl(2-[(trifluoromethyl)oxy]phenyl)thio)methyl]morpholine (24Cb)

189



Compounds **24Ca,24Cb** were obtained from **5Ca,5Cb** (0.011 g, 0.33 mmol), 2-trifluoromethoxythiophenol (1.2 eq, 0.077g, 0.39 mmol) and caesium carbonate (0.15 g, 0.47 mmol, 1.2 eq) in dimethylformamide (15 ml) following modification of **General**
 5 **Procedure 1C** in which the reaction was heated at 95°C for 1.5 hours. The reaction mixture was allowed to cool to room temperature, diluted with ethyl acetate (20 ml), washed sequentially with water and brine, dried over sodium sulphate and finally concentrated *in vacuo* to give a pale yellow oil (0.14 g, 92%); MW 459.53; C₂₅H₂₄F₃NO₂S; ¹H NMR (CDCl₃): 7.13-7.41 (13H, m), 7.08-7.13 (1H, m), 4.51 (1H, d, 8 Hz), 3.99-4.07 (2H, m), 3.73 (1H, td, 9 Hz, 2.5 Hz), 3.57 (1H, d, 13 Hz), 3.37 (1H, d, 13 Hz); 2.57-2.66 (2H, m), 2.20-2.31 (2H, m); LCMS (10 minute method): *m/z* 460 [M+H]⁺
 10 @ R_t 6.69 min.

2-(((S)-(2S)-Morpholin-2-yl(phenyl)methyl)thio)phenyl trifluoromethyl ether (25C)



15

Compound **25C (Example 9C)** was obtained from **24Ca,24Cb** (0.06 g, 0.13 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.073 g, 0.026 mmol, 2 eq) and α-chloroethyl chloroformate (0.04 ml, 0.39mmol, 3 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a viscous yellow oil (0.021 g, 44%) from which **25C** was obtained after separation using chiral HPLC on a OD semi-preparative column. Chiral LC (OJ): 12.60 min. LC purity = 98% (UV_{254nm}) / 100%
 20

(ELS); MW 369.41; $C_{18}H_{18}F_3NO_2S$; 1H NMR ($CDCl_3$): 7.02-7.21 (8H, m), 6.91-6.96 (1H, m), 4.28 (1H, d, 8 Hz), 3.93 (1H, br, d 11 Hz), 3.75-3.81 (1H, m), 3.60 (1H, td, 11 Hz and 3 Hz), 2.71-2.86 (2H, m), 2.61 (2H, d, 6 Hz), 1.90 (1H br, s); LCMS (10 minute method): m/z 370 $[M+H]^+$ @ R_t 5.86 min.


5

Example 10C: (2S)-2-[(S)-[(2-Methylphenyl)thio](phenyl)methyl]morpholine (27C)

(2S)-2-[(S)-[(2-Methylphenyl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine (26Ca)

and


10 **(2R)-2-[(R)-[(2-Methylphenyl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine (26Cb)**



Compounds **26Ca, 26Cb** were obtained from **5Ca, 5Cb** (0.1 g, 0.29 mmol), 2-methyl thiophenol (0.04 ml, 0.31 mmol) and caesium carbonate (0.125 g, 0.37 mmol, 1.2 eq) in dimethylformamide (15 ml) following **General Procedure 1C** as a colourless oil
 15 (0.13 g, 85%); MW 389.56; $C_{25}H_{27}NOS$; 1H NMR ($CDCl_3$): 6.84-7.24 (14H, m), 4.14 (1H, d, 8 Hz), 3.85-3.95 (2H, m), 3.60 (1H, dt, 10 Hz, 3 Hz), 3.42 (1H, d, 13 Hz); 3.21 (1H, d, 13 Hz), 2.46-2.54 (2H, m), 2.18 (3H, s), 1.97-2.13 (2H, m); LCMS (2.5 minute method): m/z 390 $[M+H]^+$ @ R_T 1.49 min.

20

(2S)-2-[(S)-[(2-Methylphenyl)thio](phenyl)methyl]morpholine (27C)




Compound **27C** (**Example 10C**) was obtained from **26Ca,26Cb** (0.04 g, 0.12 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.89 g, 0.24 mmol, 2 eq) and α -chloroethyl chloroformate (0.04 ml, 0.36 mmol, 3 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a viscous yellow oil (0.03 g, 75%) from which **27C** was obtained after chiral separation. Chiral LC (OJ): 15.84 min. LC purity = 98.57% (UV_{254nm}); MW 299.44; C₁₈H₂₁NOS; ¹H NMR (CDCl₃): 6.86-7.21 (9H, m), 4.08 (1H, d, 7 Hz), 3.75 (1H, br s), 3.58 (1H, br s), 2.34-3.1 (4H, m), 2.20 (3H, s); 1.41-2.04 (2H, m); LCMS (10 minute method): *m/z* 300 [M+H]⁺ @ R_T 5.08 min. **27C** was converted into its hydrochloride salt following **General Procedure 3C**.

Example 11C: (2*S*)-2-[(*S*)-Phenyl[(2-propylphenyl)thio]methyl]morpholine (29C)
(*S*)-Phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methyl-2-propylphenyl sulfide (28Ca)

and

(*R*)-Phenyl[(2*R*)-4-(phenylmethyl)morpholin-2-yl]methyl-2-propylphenyl sulfide (28Cb)




Compounds **28Ca,28Cb** were obtained from **5Ca** (0.53 g, 1.50 mmol), 2-*n*-propyl thiophenol (0.025 g, 1.65 mmol) and caesium carbonate (0.59 g, 1.8 mmol, 1.2 eq) in dimethylformamide (5 ml) following a modification of **General Procedure 1C** in which the reaction was heated at 95°C for 3 hours. After purification by SCX column chromatography (eluent: ammonia/methanol 1/1 [v/v]) **28Ca,28Cb** was obtained as a dark yellow oil (0.56 g, 90%); MW 417.62; C₂₇H₃₁NOS; ¹H NMR (CDCl₃): 7.23-7.12 (6H, m), 7.06-7.11 (5H, m), 6.97-6.99 (2H, m), 6.87-6.92 (1H, m), 4.13 (1H, d, 8 Hz), 3.86-3.94 (2H, m), 3.61 (1H, td, 11 Hz, 2 Hz), 3.44 (1H, d, 13 Hz), 3.23 (1H, d, 13 Hz),

192

2.46-2.59 (4H, m), 2.01-2.14 (2H, m), 1.34-1.52 (2H, m), 0.83 (3H, t, 7 Hz); LCMS (2.5 minute method): m/z 418 $[M+H]^+$ @ R_t 1.55 min.

(2S)-2-((S)-Phenyl[(2-propylphenyl)thio]methyl)morpholine (29C)



5

Compound **29C** (**Example 11C**) was obtained from **28Ca,28Cb** (0.56 g, 1.35 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.75 g, 2.7 mmol, 2 eq) and α -chloroethyl chloroformate (0.44 ml, 4.05 mmol, 3 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a viscous yellow oil (0.41 g, 93%); MW 327.49; $C_{20}H_{25}NOS$; 1H NMR ($CDCl_3$): 7.17 (1H, br, d, 7 Hz), 7.07-7.12 (5H, m), 6.96-7.00 (2H, m), 6.88-6.93 (1H, m), 4.07 (1H, d, 8 Hz), 3.93-3.98 (1H, m), 3.74-3.80 (1H, m), 3.60 (1H, td, 11 Hz, 3 Hz), 2.81 (1H, td, 12 Hz and 3 Hz), 2.72 (1H, br, d, 12 Hz), 2.48-2.62 (4H, m), 1.36-1.59 (3H, m), 0.83 (3H, t, 7 Hz); LCMS (2.5 minute method): m/z 328 $[M+H]^+$ @ R_t 1.40 min (single major peak).

15


Example 12C: Methyl 2-(((S)-(2S)-morpholin-2-yl(phenyl)methyl)thio)benzoate (31C)

Methyl-2-(((S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methyl)thio)benzoate (30Ca)

20 and


Methyl-2-(((R)-phenyl[(2R)-4-(phenylmethyl)morpholin-2-yl]methyl)thio)benzoate (30Cb)

193



Compounds **30Ca,30Cb** were obtained from **5Ca,5Cb** (0.5 g, 1.45 mmol), methyl thiosalicylate (0.49 g, 2.89 mmol) and potassium carbonate (0.21 g, 1.52 mmol) in dry tetrahydrofuran (5 ml) following modification of **General Procedure 1C** in which the solvents were degassed and purged with nitrogen before the addition of methyl thiosalicylate. The reaction mixture was stirred at room temperature for 18 hours after which time the reaction mixture was poured onto water and extracted twice with diethyl ether. The organic layers were washed with water, dried and evaporated *in vacuo*. After purification by SCX column chromatography (eluent: ammonia/methanol 1/1 [v/v]) **30Ca,30Cb** was obtained as a colourless solid (0.18 g, 29%); MW 433.57; $C_{26}H_{27}NO_3S$; 1H NMR ($CDCl_3$): 8.65-8.85 (1H, m), 6.95-7.45 (13H, m), 4.45 (1H, d, 8 Hz), 3.85-4.05 (1H, m), 3.8 (3H, s), 3.65 (1H, dt, 1 Hz and 7 Hz), 3.55 (1H, d, 11 Hz), 3.25 (1H, d, 11 Hz), 2.5-2.6 (2H, m); 2.0-2.15 (2H, m); FIA: m/z 462 $[M+H]^+$.

15 Methyl 2-[(*S*)-(2*S*)-morpholin-2-yl(phenyl)methyl]thio}benzoate (31C)



Compound **31C (Example 12C)** was obtained from **30Ca,30Cb** (0.2 g, 0.46 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.08 g, 2.77 mmol, 6 eq) and α -chloroethyl chloroformate (0.5 ml, 4.62 mmol, 10 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a white solid (0.16 g, 91%) from which **31C** was obtained after separation using chiral HPLC on chiral OJ

semi-preparative column. Chiral LC (OJ): 12.32 min. LC purity = 100% (UV_{254nm}); MW 343.45. **31** was converted into its hydrochloride salt following **General Procedure 3C**; ¹H NMR (d₆-DMSO): 9.30-9.5 (1H, m), 7.75-7.80 (1H, m), 7.1-7.55 (8H, m), 4.82 (1H, d, 8 Hz), 3.95-4.15 (2H, m), 3.65-3.9 (3H, m), 3.55 (3H, s), 2.80-3.25 (2H, m).


5

Example 13C: (2S)-2-((S)-(3-Fluorophenyl){[2-(trifluoromethyl)phenyl]thio}methyl)morpholine (33C)

(2S)-2-((S)-(3-Fluorophenyl){[2-(trifluoromethyl)phenyl]thio}methyl)-4-(phenylmethyl)morpholine (32Ca)

10 and

(2R)-2-((R)-(3-Fluorophenyl){[2-(trifluoromethyl)phenyl]thio}methyl)-4-(phenylmethyl)morpholine (32Cb)




Compounds **32Ca,32Cb** were obtained as outlined in **Scheme 5C** from **38Ca,38Cb** (0.33 g, 0.91 mmol) following **General Procedure 4C** as a white solid after column chromatography (0.28 g, 67%); MW 461.53; C₂₅H₂₃F₄NOS; ¹H NMR (CDCl₃) 6.75-7.65 (1H, m), 6.85-7.33 (12H, m), 4.45 (2H, d, 8 Hz), 3.6-3.75 (2H, m), 3.45 (1H, d 12 Hz), 3.3 (1H, d 12 Hz), 2.45-2.7 (2H, br, m), 2.1-2.3 (2H, br, m); FIA: *m/z* 462 [M+H]⁺.

20

(2S)-2-((S)-(3-Fluorophenyl){[2-(trifluoromethyl)phenyl]thio}methyl)morpholine (33C)


195



Compound **33C (Example 13C)** was obtained from **32Ca,32Cb** (0.28 g, 0.615 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.19 g, 0.68 mmol, 1.1 eq) and α -chloroethyl chloroformate (0.07 ml, 0.68 mmol, 1.1 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a colourless oil (0.22 g, 95%) from which **33C** was obtained after chiral chromatography on a Chiralcel OJ semi-preparative column. Chiral LC (OJ): 13.33 min. LC purity = 98.37% (UV_{254nm}); MW 371.4; C₁₈H₁₇F₄NOS. LCMS (12 minute method): m/z 372 [M+H]⁺ @ Rt 5.2 min. **33C** was converted into its hydrochloride salt following **General Procedure 3C**; MW 407.86; C₁₈H₁₇F₄NOS.HCl; ¹H NMR (CDCl₃) 9.8-10.2 (1H, br), 7.4-7.6 (1H, m), (6.85-7.45 (8H, m), 4.05-4.45 (4H, br, m), 2.90-3.41 (4H, br, m).


Example 14C: (2S)-2-((S)-(4-Chlorophenyl){[2-(trifluoromethyl)phenyl]thio}methyl)morpholine (35C)

(2S)-2-((S)-(4-Chlorophenyl){[2-(trifluoromethyl)phenyl]thio}methyl)-4-(phenylmethyl)morpholine (34Ca)
and
(2R)-2-((R)-(4-Chlorophenyl){[2-(trifluoromethyl)phenyl]thio}methyl)-4-(phenylmethyl)morpholine (34Cb)



Compounds **34Ca,34Cb** were obtained as outlined in **Scheme 5C** from **39Ca,39Cb** (0.4 g, 1.06 mmol, 1.1 eq), cesium carbonate (0.33 g, 1.0 mmol, 1.1 eq), and 2-trifluoromethyl benzene thiol (0.19 g, 1.06 mmol, 1.1 eq) following a modification of **General Procedure 1C** in which the reaction was stirred at room temperature for 1.5 hours as a white solid after column chromatography (eluent: gradient hexane/ethyl acetate 10/90 to 25/75[v/v]) (0.409g, 80%); MW 477.98; C₂₅H₂₃F₃ClNOS; ¹H NMR (CDCl₃) 7.1-7.65 (13H, m), 4.45 (1H, d, 8 Hz), 3.85-4.0 (2H, m), 3.55 (1H, m), 3.3 (1H, d 12 Hz), 3.3 (1H, d 12 Hz), 2.45-2.65 (2H, br,), 2.1-2.3 (2H, br, m); FIA: *m/z* 478 [M+H]⁺.

10 **(2S)-2-((S)-(4-Chlorophenyl){2-(trifluoromethyl)phenyl}thio)methyl)morpholine (35C)**




Compound **35C (Example 14C)** was obtained from **34Ca,34Cb** (0.41 g, 0.86 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.27 g, 0.94 mmol, 1.1 eq) and α -chloroethyl chloroformate (0.10 ml, 0.94 mmol, 1.1 eq) in anhydrous dichloromethane (5 ml) following **General Procedure 2Ca** as a colourless oil (0.28 g, 84% yield) from which **35C** was obtained after separation using chiral HPLC on a ChiralPak-AD OJ semi-preparative column; MW 387.85; C₁₈H₁₇ClF₃NOS; LCMS (12 minute method): *m/z* 372 [M+H]⁺ @ Rt 5.2 min. **35C** was converted into its hydrochloride salt following **General Procedure 3C**; MW 423.96; C₁₈H₁₇ClF₃NOS.HCl; ¹H NMR (CDCl₃): 9.8-10.2 (1H, br), 7.4-7.6 (1H, m), 7.07-7.35 (7H, m), 3.8-4.45 (4H, br, m), 2.85-3.45 (4H, br, m).

25 **Example 15C: (2S)-2-((S)-(2-Fluorophenyl){2-(methoxy)phenyl}thio)methyl)morpholine (37C)**
(2S)-2-((S)-(2-Fluorophenyl){2-(methoxy)phenyl}thio)methyl)-4-(phenylmethyl)morpholine (36Ca)


and

(2*R*)-2-((*R*)-(2-Fluorophenyl){[2-(methyloxy)phenyl]thio}methyl)-4-(phenylmethyl)morpholine (36Cb)



- 5 Compounds **36Ca,36Cb** were obtained from **7Ca,7Cb** (0.45 g, 1.17 mmol), cesium carbonate (0.42 g, 1.29 mmol, 1.1 eq), and 2-methoxy-thiophenol (0.82 g, 5.87 mmol) following a modification of **General Procedure 1C** in which the reaction mixture was heated to 95°C for 2 hours and then stirred at room temperature for 18 hours. After purification by flash column chromatography (eluent: heptane/ethyl acetate 80/20 [v/v])
- 10 **36Ca,36Cb** was obtained as a colourless oil (0.36 g, 72%); MW 423.55; C₂₅H₂₆FNOS; ¹H NMR (CDCl₃): 6.65-7.5 (13H, m), 4.9 (1H, d, 7 Hz), 3.9-4.05 (2H, m), 3.8 (3H, s), 3.6 (1H, dt, 8 Hz and 1 Hz), 3.45 (1H, d, 13 Hz), 3.15 (1H, d, 13 Hz), 2.60 (2H, t, 8 Hz), 2.05-2.2 (2H, m); FIA: *m/z* 424 [M+H]⁺.

- 15 (2*S*)-2-((*S*)-(2-Fluorophenyl){[2-(methyloxy)phenyl]thio}methyl)morpholine (37C)




- Compound **37C (Example 15C)** was obtained from **36Ca,36Cb** (0.43 g, 1.02 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.37 g, 1.12 mmol, 1.1 eq) and α-chloroethyl chloroformate (1.08 ml, 10.12 mmol, 10 eq) in anhydrous
- 20 dichloromethane (5 ml) following **General Procedure 2Ca** as a colourless oil (0.34 g, 99%) after separation by chiral HPLC on a ChiralPak-AD semi-preparative column. Chiral LC: 12.86 min. LC purity = 99.1 (UV_{254nm}); MW 369.89; C₁₈H₂₀FNOS; FIA: *m/z*

334 $[M+H]^+$. **37C** was converted into its hydrochloride salt following **General Procedure 3C**; MW 333.43; $C_{18}H_{20}FNOS$; 1H NMR ($CDCl_3$): 7.2-7.3 (1H, m), 6.85-7.2 (8H, m), 4.85 (1H, d, 8 Hz), 3.95-4.15 (2H, m), 3.85-3.9 (3H, m), 3.7 (1H, dt, 1 Hz and 7 Hz), 2.6-3.0 (4H, m).

5


Example 16C: 2-[2-Methyl-1-(2-trifluoromethyl-phenylsulfanyl)-propyl]-morpholine (56C)

4-Benzyl-2-(1-hydroxy-2-methyl-propyl)-morpholin-3-one (53C)



10 To a stirred solution of **2C** (5.05 g, 26.4 mmol) in tetrahydrofuran (25 ml) at –78°C under nitrogen was added lithium diisopropylamide (14.5 ml of a 2M solution, 29.0 mmol) dropwise over 40 minutes. The reaction mixture was stirred at the same temperature over 30 minutes after which time a solution of isobutyraldehyde (2.63 ml, 29.0 mmol) in tetrahydrofuran (15 ml) was added dropwise over 30 minutes. After one
15 hour, the reaction mixture was allowed to warm to room temperature and quenched by addition of saturated ammonium chloride solution. Extraction with dichloromethane and drying over magnesium sulphate gave **53C** as a mixture of diastereomers. Upon concentration *in vacuo* one diastereomer precipitated as a white solid (**53Ca**: 0.99 g). The remaining mother liquors were purified by column chromatography (30% ethyl acetate in
20 hexane [v/v]) to give **53C** (2.06 g). MW 263.34; $C_{15}H_{21}NO_3$; LCMS (6 min method): m/z 286 $[M+Na]^+$; RT = 2.748.


1-(4-Benzyl-morpholin-2-yl)-2-methyl-propan-1-ol (54C)




25 To a stirred solution of **53C** (1.97 g, 7.47 mmol) in tetrahydrofuran (50 ml) at room temperature under nitrogen was added borane-tetrahydrofuran complex (30 ml of a

1M solution, ca 4 eq.). The reaction was heated to 60°C and followed by TLC-analysis. When all starting material had been consumed a few drops of methanol were added followed by a similar amount of 1N hydrochloric acid and heating was continued for another hour. Organic solvents were removed *in vacuo* and the remaining solution was poured onto 1M potassium carbonate solution (30 ml), extracted with diethyl ether. The organic layers were dried over magnesium sulphate and purified by column chromatography (gradient from 15% ethyl acetate in hexane [v/v]) gave **54C** (1.8 g, 97%). MW 249.36; C₁₅H₂₃NO₂; LCMS (6 min method): *m/z* 250 [M+H]⁺; RT = 0.838.

10 **4-Benzyl-2-[2-methyl-1-(2-trifluoromethyl-phenylsulfanyl)-propyl]-morpholine (55C)**




Compound **55C** was obtained from **54C** in a two-step procedure. To a stirred solution of **54C** (1.8 g, 7.2 mmol) in dichloromethane (50 ml) at room temperature was added solid solid supported Hünig's base (Argonaut, 3.56 mmol/g, 6.2 g, 22 mmol, 3 eq) followed methanesulphonyl chloride (1.12 ml, 14 mmol). After stirring for one hour, the reaction mixture was filtered and the filtrates washed with brine and dried over magnesium sulphate to give the intermediate mesylate as a yellow oil (2.93 g of isolated crude product). The crude product was taken up in dry dimethylformamide (50 ml), 2-trifluoromethyl benzenethiol (2.1 ml, 14 mmol) and solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.55 g, 1.95 mmol) were added and the mixture heated to 70°C and stirred for 72 hours. The reaction was quenched by addition of water (50 ml) and sodium hydroxide solution (70 ml of a 2N solution). The aqueous layer was extracted with diethyl ether (3x50 ml), washed with brine and dried over magnesium sulphate. Purification by ion-exchange chromatography followed by preparative HPLC gave **55C**. MW 409.52; C₂₂H₂₆F₃NOS; LCMS (6 min method): *m/z* 410 [M+H]⁺; RT = 3.398.

2-[2-Methyl-1-(2-trifluoromethyl-phenylsulfanyl)-propyl]-morpholine (56C)

Compound **56C** (**Example 16C**) was obtained from **55C** (0.8 g, 1.95 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 1.65 g, 5.85 mmol, 3 eq) and α -chloroethyl chloroformate (0.4 ml, 3.9 mmol, 2 eq) in anhydrous dichloromethane (20 ml) following **General Procedure 2Ca** as a colourless oil (0.5 g, 85% yield). Chiral HPLC on a ChiralCel-OD(3671) column using 50% heptane in ethanol [v/v] gave 2 fractions (R_t = 8.793 min and 10.443 min). Conversion into fumarate salt **56C** was carried out by dissolving in diethyl ether and addition of small amount of methanol. Data for **56C** derived from fraction with R_t = 8.793 min: MW 435.46; $C_{19}H_{24}F_3NO_5S$; 1H NMR (d_3 -MeOD): 6.2-6.3 (2H, m), 6.1-6.2 (1H, m), 5.2 (1H, s), 2.6-2.7 (2H, m), 2.2-2.4 (1H, m), 1.6-1.9 (4H, m), 1.6-1.7 (1H, m), -0.4- -0.5 (6H, m).

Example 17C: 2-[2-Methyl-1-(2-trifluoromethyl-phenoxy)-propyl]-morpholine (58C)


4-Benzyl-2-[2-methyl-1-(2-trifluoromethyl-phenoxy)-propyl]-morpholine (57C)



To a solution of **53Ca** (0.146 g, 0.585 mmol) in dry dimethylformamide (2 ml) under nitrogen and ice-cooling was added sodium hydride (26 mg of a 60% dispersion in oil, 0.644 mmol) portionwise. The reaction was allowed to warm to room temperature for 30 minutes before addition of 2-fluoro-benzotrifluoride (0.07 ml, 0.66 mmol). After stirring for 12 hours, another 0.5 equivalents of reagents were added and the reaction mixture heated to 40°C for 30 minutes and then to 60°C for another 2 hours. The crude reaction mixture was purified by ion-exchange column chromatography followed by preparative

HPLC to give **57C** (0.208 g, 92% yield) MW 393.45; C₂₂H₂₆F₃NO₂; LCMS (6 min method): *m/z* 394 [M+H]⁺; RT = 3.150.

2-[2-Methyl-1-(2-trifluoromethyl-phenoxy)-propyl]-morpholine (58C)



Compound **58C** (**Example 17C**) was obtained from **57C** (0.21 g, 0.53 mmol), solid supported Hünig's base (Argonaut, 3.56 mmol/g, 0.45 g, 1.5 mmol, 3 eq) and α -chloroethyl chloroformate (0.11 ml, 1.06 mmol, 2 eq) in anhydrous dichloromethane (10 ml) following **General Procedure 2C** as a colourless oil (0.147 g, 92% yield) MW 303.33; C₁₅H₂₀F₃NO₂; ¹H NMR (CDCl₃): 7.5-7.6 (1H, m), 7.2-7.4 (1H, m), 7.0-7.1 (1H, m), 6.8-6.95 (1H, m), 4.15-4.25 (1H, m), 3.6-3.9 (2H, m), 3.4-3.6 (1H, m), 2.6-2.9 (4H, m), 2.15 (1H, br, s) 1.8-2.1 (1H, m), 1.1-1.2 (6H, m); LCMS (12 min method): *m/z* 304 [M+H]⁺; RT = 4.862.

The following examples illustrate compounds of of Formulae (ID) above and methods for their preparation.

Scheme 1D - Preparation of Intermediates

1-Phenyl-3,4-dihydro-1H-quinolin-2-one (2Da)

A stirred mixture of 3,4-Dihydro-1H-quinolin-2-one (**1Da**) (1.47 g, 10 mmol), K₂CO₃ (2.9 g, 21 mmol), *trans*-cyclohexane-1,2-diamine (240 μ L, 2 mmol) and bromobenzene (3.16 mL, 30 mmol) in 1,4-dioxane (10 mL) was heated under a nitrogen atmosphere at 125°C for 5 min to deoxygenate the reaction mixture. Copper (I) iodide (380 mg, 2 mmol) was added in one portion and the reaction mixture was refluxed overnight at 125°C. After cooling to rt, the reaction mixture was poured into ethyl acetate (100 mL) and extracted with water. The organic layer was separated, dried over MgSO₄ and concentrated. Treatment of the residue with ether (100 mL) and cooling (ice bath) gave the product as a white solid after filtration (1.77 g, 79%).

6-Fluoro-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (2Db)

This was prepared using the method described for (2Da) using 6-Fluoro-3,4-dihydro-1*H*-quinolin-2-one (1Db) (617 mg, 3.7 mmol) and 4-bromotoluene (1.91 g, 11 mmol) to give
5 the crude product, which was purified using automated chromatography (silica) (0 to 60% ethyl acetate\cyclohexane gradient) to provide the product as a light brown solid (880 mg, 92%).

3-Methyl-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (3Da)

10 To a soln of (2Da) (892 mg, 4 mmol) in anhydrous THF (40 mL) at -78°C under nitrogen was added LiHMDS (4.4 mL, 1M soln in hexanes, 4.4 mmol) dropwise over 10 min. The reaction mixture was left at -78°C for 30 min and then a solution of methyl iodide (298 µL, 4.8 mmol) in THF (1 mL) was added dropwise. The reaction mixture was warmed slowly to rt, quenched with water (2 mL) and extracted with ethyl acetate (100 mL). The
15 organic layer was separated, dried over MgSO₄ and concentrated. The residue was purified by column chromatography (silica, gradient 100% hexane to ethyl acetate\hexane 3:10) giving the product as an oil (667 mg, 70%).

3-Ethyl-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (3Db)

20 This was prepared in a similar manner to (3Da) on a 1.5 mmol scale using 1-iodoethane (125 µL, 1.1 eq.) as the alkylating agent. The crude product (378 mg) was used directly in the next step.

3-(3-Chloro-propyl)-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (4Da)

25 To a soln of (2Da) (892 mg, 4 mmol) in anhydrous THF (40 mL) at -78°C under nitrogen was added LiHMDS (4.4 mL, 1M soln in hexanes, 4.4 mmol) dropwise over 10 min. The reaction mixture was left at -78°C for 30 min and then a solution of 1-bromo-3-chloropropane (405 µL, 4.4 mmol) in THF (1 mL) was added dropwise. The reaction mixture was warmed slowly to rt, quenched with water (2 mL) and extracted with ethyl
30 acetate (100 mL). The organic layer was separated, dried over MgSO₄ and concentrated. The crude product (1.2 g) was used directly in the next step.

3-(3-Chloro-propyl)-6-fluoro-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (4Db)

This was prepared from (2Db) (300 mg, 1.17 mmol) using the method described for (4Da) using 1-bromo-3-chloropropane (140 μ L, 1.4 mmol) as the alkylating agent. The crude product (399 mg) was used directly in the next step.

5

3-(2-Chloro-ethyl)-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (4Dc)

This was prepared from (2Da) (892 mg, 4.0 mmol) using the method described for (4Da) using 1-bromo-2-chloroethane (365 μ L, 4.4 mmol) as the alkylating agent. The crude product (1 g) was used directly in the next step.

10

3-(3-Chloro-propyl)-3-methyl-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (5Da)

This was prepared from (3Da) (462 mg, 1.95 mmol) using the method described for (4Da) using 1-bromo-3-chloropropane (270 μ L, 2.7 mmol) as the alkylating agent. The crude product (650 mg) was used directly in the next step.

15

3-(3-Chloro-propyl)-3-ethyl-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (5Db)

This was prepared from (3Db) (378 mg, 1.5 mmol) using the method described for (4Da) using 1-bromo-3-chloropropane (179 μ L, 1.8 mmol) as the alkylating agent. The crude product (528 mg) was used directly in the next step.

20

Scheme 1D - Examples**Example 1D: 3-(3-Methylamino-propyl)-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (6Da)**

25 A soln of (4Da) (1.2 g, 4 mmol), potassium iodide (200 mg, 1.2 mmol) and aqueous 40% methylamine (12 mL) in ethanol (30 mL) was refluxed at 100°C under nitrogen for 3 h. The reaction mixture was cooled, poured into water and extracted with ethyl acetate (100 mL). The organic layer was separated, dried over MgSO₄ and concentrated. The product was purified by preparative LCMS to give 500 mg of the racemate. The racemate was
30 separated into its individual enantiomers using chiral HPLC. ¹H NMR (300 MHz, CDCl₃) (racemate & isomer) δ 1.5-1.73 (m, 4H), 1.88-1.97 (m, 1H), 2.43 (s, 3H), 2.62 (t, J= 6.69

Hz, 2H), 2.70-2.79 (m, 1H), 2.84-2.92 (m, 1H), 3.15 (dd, J= 15.45, 5.28 Hz, 1H), 6.33 (d, J= 7.73 Hz, 1H), 6.95-7.06 (m, 2H), 7.19-7.22 (m, 3H), 7.38-7.43 (m, 1H), 7.47-7.52 (m, 2H). LCMS (12 minute method) $[M+H]^+ = 295$ @ Rt 4.0 min (100%).

5 **Example 2D: 6-Fluoro-3-(3-methylamino-propyl)-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (6Db)**

This was prepared in an identical manner to (6Da) using crude (4Db) (399 mg) to give the crude product, which was purified by preparative LCMS to give the product (35 mg).

¹H NMR (300 MHz, CDCl₃) (racemate) δ 1.40-1.70 (m, 3H), 1.75-1.90 (m, 4H), 2.34 (s, 3H), 2.36 (s, 3H), 2.50-2.83 (m, 2H), 3.01-3.08 (m, 1H), 6.21-6.26 (m, 1H), 6.62-6.68 (m, 1H), 6.82-6.86 (m, 1H), 6.99 (d, J= 8.1 Hz, 2H), 7.22 (d, J= 8.1 Hz, 2H). LCMS (12 minute method) $[M+H]^+ = 327$ @ Rt 4.8 min (100%).

15 **Example 3D: 3-(2-Methylamino-ethyl)-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (6Dc)**

This was prepared in an identical manner to (6Da) using crude (4Dc) (1g) to give the racemate (80 mg). The racemate was separated into its individual enantiomers using chiral HPLC. ¹H NMR (300 MHz, CDCl₃) (racemate & isomer) δ ppm 1.64-1.76 (m, 1H), 1.79 (br, 1H), 2.03-2.18 (m, 1H), 2.44 (s, 3H), 2.71-2.82 (m, 2H), 2.82-2.94 (m, 2H), 3.09-3.21 (m, 1H), 6.33 (dd, J= 7.91, 1.32 Hz, 1H), 6.94-7.07 (m, 2H), 7.18-7.24 (m, 3H), 7.37-7.44 (m, 1H), 7.47-7.54 (m, 2H). LCMS (12 minute method) $[M+H]^+ = 281$ @Rt 3.82 min (100%).

25 **Example 4D: 3-Methyl-3-(3-methylamino-propyl)-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (7Da)**

This was prepared in an identical manner to (6Da) using crude (5Da) (650 mg) to give the crude product (198 mg), which was purified by preparative LCMS. The purified racemate was then separated into its individual enantiomers using chiral HPLC. ¹H NMR (300 MHz, CDCl₃) (isomer) δ ppm 1.27 (s, 3H), 1.43 (br, 1H), 1.53-1.66 (m, 4H), 2.39 (s, 3H), 2.54 (t, J= 6.12 Hz, 2H), 2.91 (d, J= 15.64 Hz, 1H), 2.98 (d, J= 15.64 Hz, 1H), 6.28 (dd, J= 7.91, 1.32 Hz, 1H), 6.97 (td, J= 7.21, 1.41 Hz, 1H), 7.03 (td, J= 7.68, 1.98 Hz, 1H),

7.14-7.22 (m, 3H), 7.36-7.44 (m, 1H), 7.46-7.53 (m, 2H). LCMS (12 minute method) $[M+H]^+ = 309$ @Rt 4.21 min (100%).

Example 5D: 3-Ethyl-3-(3-methylamino-propyl)-1-phenyl-3,4-dihydro-1H-quinolin-

5 **2-one (7Db)**

This was prepared in an identical manner to (6Da) using crude (5Db) (528 mg) to give the crude product (105 mg), which was purified by preparative LCMS. The purified racemate was then separated into its individual enantiomers using chiral HPLC.

^1H NMR (300 MHz, CDCl_3) (racemate) δ 0.93 (t, $J = 7.53$ Hz, 3H), 1.56-1.75 (m, 6H),
10 1.91 (bs, 1H), 2.41 (s, 3H), 2.55-2.60 (m, 2H), 2.91 (d, $J = 15.82$, 1H), 3.02 (d, $J = 15.82$, 1H), 6.25-6.28 (m, 1H), 6.94-7.05 (m, 2H), 7.16-7.19 (m, 3H), 7.38-7.43 (m, 1H), 7.4-7.52 (m, 2H). ^1H NMR (300 MHz, MeOD-d_4) (isomer D-tartrate salt) δ 0.85 (t, $J = 7.53$ Hz, 3H), 1.45-1.75 (m, 6H), 2.57 (s, 2H), 2.83-2.89 (m, 2H), 3.01-3.06 (d, $J = 16.01$, 1H), 4.32 (s, 2H), 6.11-6.14 (m, 1H), 6.89-6.97 (m, 2H), 7.09 (d, $J = 7.16$ Hz, 2H), 7.15-7.18
15 (m, 1H), 7.37 (t, $J = 7.35$ Hz, 1H), 7.46 (t, $J = 7.35$ Hz, 2H). LCMS (12 minute method) $[M+H]^+ = 323$ @ Rt 4.9 min (98%).

Scheme 2D - Preparation of Intermediates

20 **1-*p*-Tolyl-3,4-dihydro-1H-quinolin-2-one (2Dc)**

A stirred mixture of 3,4-Dihydro-1H-quinolin-2-one (1Da) (4.41 g, 30 mmol), K_2CO_3 (8.7 g, 63 mmol), *trans*-cyclohexane-1,2-diamine (720 μL , 2 mmol) and 4-bromotoluene (15.4 g, 90 mmol) in 1,4-dioxane (30 mL) was heated under a nitrogen atmosphere at 125°C for 5 min to deoxygenate the reaction mixture. Copper (I) iodide (1.14 g, 2 mmol)
25 was added in one portion and the reaction mixture was refluxed overnight at 125°C. After cooling to rt, the reaction mixture was filtered through celite, poured into ethyl acetate (100 mL) and extracted with water. The organic layer was separated, dried over MgSO_4 and concentrated. Treatment of the residue with ether (200 mL) and cooling (ice bath) gave the product as a white solid after filtration (6.2 g, 87%).

30

1-Phenyl-3-propyl-3,4-dihydro-1H-quinolin-2-one (3Dc)

This was prepared from (**2Da**) (669 mg, 3 mmol) and 1-iodopropane (352 μ l, 1.2 eq.) as the alkylating agent. The crude product (780 mg) was used directly in the next step.

3-Ethyl-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (3Dd)

- 5 This was prepared from (**2Dc**) (711 mg, 3 mmol) and 1-iodoethane (265 μ l, 1.2 eq.) as the alkylating agent. The crude product (800 mg) was used directly in the next step.

3-Propyl-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (3De)

- 10 This was prepared from (**2Dc**) (711 mg, 3 mmol) and 1-iodopropane (352 μ l, 1.2 eq.) as the alkylating agent. The crude product (840 mg) was used directly in the next step.

3-Butyl-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (3Df)

- 15 This was prepared from (**2Dc**) (711 mg, 3 mmol) and 1-iodobutane (354 μ l, 1.1 eq.) as the alkylating agent. The crude product (790 mg) was used directly in the next step.

3-Isopropyl-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (3Dg)

This was prepared from (**2Dc**) (711 mg, 3 mmol) and 2-iodopropane (330 μ l, 1.1 eq.) as the alkylating agent. The crude product (806 mg) was used directly in the next step.

20 **3-Allyl-3-ethyl-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (11Db)**

- To a soln of (**3Dd**) (800 mg, 2.7 mmol) in anhydrous THF (30 mL) at -78°C under nitrogen was added LiHMDS (3 mL, 1M soln in hexanes, 3 mmol) dropwise over 10 min. The reaction mixture was left at -78°C for 30 min and then a solution of allyl bromide (280 μ L, 3.2 mmol) in THF (1 mL) was added dropwise. The reaction mixture was
25 warmed slowly to rt, quenched with water (2 mL) and extracted with ethyl acetate (100 mL). The organic layer was separated, dried over MgSO_4 and concentrated. The crude product (920 mg) was used directly in the next step.

3-Ethyl-3-(3-hydroxypropyl)-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (12Db)

- 30 To a soln of (**11Db**) (732 mg, 2.4 mmol) in anhydrous THF (25 mL) at 0°C under nitrogen was added 9-BBN (12 mL, 0.5M soln in THF, 6 mmol, 2.5 eq.) dropwise over

10 min. The reaction mixture was warmed to rt and left to stir overnight. The resultant yellow soln was cooled to 0°C and then quenched carefully with ethanol (3 mL), followed by aq. NaOH (1.8 mL, 3N soln). Finally, aq. H₂O₂ (1.8 mL, 37% soln) was added dropwise maintaining the internal reaction mixture temp between 5 and 10 °C. The reaction mixture was warmed to rt and then refluxed for 90 min. The reaction mixture was cooled to rt, poured into ethyl acetate and water and extracted. The organic layer was separated, dried over MgSO₄ and concentrated. The crude product was purified using automated chromatography (silica) (0 to 60% ethyl acetate/cyclohexane gradient) to provide (12Db) as a clear oil (540 mg, 70%).

10

Scheme 2D - Examples

Example 6D: 3-Ethyl-3-(3-methylamino-propyl)-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (13Db)

15 To a soln of (12Db) (540 mg, 1.67 mmol) and triethylamine (350 µL, 2.5 mmol) in anhydrous THF (20 mL) at 0°C under nitrogen was added dropwise a soln of methanesulfonyl chloride (142 µL, 1.8 mmol) in THF (1 mL). The reaction mixture was warmed to rt and stirred for 3 h. The reaction mixture was poured into ethyl acetate and water and extracted. The organic layer was separated, dried over MgSO₄ and concentrated. The crude mesylate (670 mg, 100%) was dissolved in ethanol (10 mL) and aqueous 40% methylamine (5 mL) and heated at 65°C under nitrogen for 2 h. The reaction mixture was cooled, poured into water and extracted with ethyl acetate (100 mL). The organic layer was separated, dried over MgSO₄ and concentrated. The product was purified by SCX-2 to give 384 mg of the racemate. The racemate was separated into its individual enantiomers using chiral HPLC. Each enantiomer was dissolved in CH₂Cl₂ (2 mL) and treated with 1 equivalent of D-tartaric acid dissolved in a minimum volume of warm methanol. The resultant soln was concentrated and the solid was dried under vacuo to provide the D-tartrate salt of the amine. ¹H NMR (300 MHz, CDCl₃) (racemate) δ 0.92 (t, J= 7.44 Hz, 3H), 1.49-1.75 (m, 6H), 1.81 (br, 1H), 2.40 (s, 6H), 2.57 (t, J= 6.59 Hz, 2H), 2.89 (d, J= 15.82 Hz, 1H), 3.00 (d, J= 15.82 Hz, 1H), 6.29 (d, J= 7.91 Hz, 1H), 6.92-

20

25

30

7.08 (m, 4H), 7.16 (d, J= 7.16 Hz, 1H), 7.29 (d, J= 7.91 Hz, 2H). ¹H NMR (300 MHz, MeOD-d₄) (isomer D-tartrate salt) δ 0.93 (t, J= 7.44 Hz, 3H), 1.54-1.84 (m, 6H), 2.42 (s, 3H), 2.66 (s, 3H), 2.91-3.00 (m, 3H), 3.11 (d, J= 15.83 Hz, 1H), 4.41 (s, 2H), 6.22-6.27 (m, 1H), 6.80-7.07 (m, 4H), 7.21-7.27 (m, 1H), 7.36 (d, J= 7.91 Hz, 2H). LCMS (12 minute method) [M+H]⁺ = 337 @Rt 5.21 min (100%).

Example 7D: 3-(3-Methylamino-propyl)-1-phenyl-3-propyl-3,4-dihydro-1H-quinolin-2-one (13Da)

This was prepared from (3Dc) (780 mg, 2.9 mmol) using the same synthetic sequence described above (3Dd to 13Db) to give 233 mg of the racemate. The racemate was separated into its individual enantiomers using chiral HPLC and each enantiomer was converted into its D-tartrate salt as described for (13Db). ¹H NMR (300 MHz, CDCl₃) (racemate) δ 0.88 (t, J= 7.16 Hz, 3H), 1.26-1.48 (m, 2H), 1.50-1.78 (m, 7H), 2.40 (s, 3H), 2.56 (t, J= 6.59 Hz, 2H), 2.92 (d, J= 15.83 Hz, 1H), 3.01 (d, J= 15.83 Hz, 1H), 6.25-6.28 (m, 1H), 6.94-7.05 (m, 2H), 7.16-7.19 (m, 3H), 7.37-7.42 (m, 1H), 7.47-7.52 (m, 2H). ¹H NMR (300 MHz, MeOD-d₄) (isomer D-tartrate salt) δ 0.77-0.82 (t, J= 7.06 Hz, 3H), 1.24-1.35 (m, 2H), 1.44-1.51 (m, 2H), 1.69 (bs, 3H), 2.56 (s, 3H), 2.84-2.89 (m, 3H), 3.01-3.06 (d, J= 15.83 Hz, 1H), 3.20-3.22 (q, J=1.55 Hz, 2H), 4.30 (s, 2H), 6.11-6.14 (dd, J= 7.72, 2.26 Hz, 1H), 6.89-6.97 (m, 2H), 7.07-7.10 (m, 2H), 7.14-7.17 (m, 1H), 7.34-7.39 (t, J= 7.35 Hz, 1H), 7.43-7.48 (t, J= 7.35 Hz, 2H). LCMS (12 minute method) [M+H]⁺ = 337 @ Rt 5.2 min (100%).

Example 8D: 3-(3-Methylamino-propyl)-3-propyl-1-p-tolyl-3,4-dihydro-1H-quinolin-2-one (13Dc)

This was prepared from (3De) (840 mg, 2.6 mmol) using the same synthetic sequence described above (3Dd to 13Db) to give 393 mg of the racemate. The racemate was separated into its individual enantiomers using chiral HPLC and each enantiomer was converted into its D-tartrate salt as described for (13Db). ¹H NMR (300 MHz, CDCl₃) (racemate) δ 0.88 (t, J= 7.16 Hz, 3H), 1.20-1.75 (m, 11H), 2.39 (s, 3H), 2.40 (s, 3H), 2.90 (d, J= 15.64 Hz, 1H), 2.99 (d, J= 15.64 Hz, 1H), 6.29 (d, J= 7.72 Hz, 1H), 6.93-7.07 (m, 4H), 7.14-7.16 (m, 1H), 7.25-7.31 (m, 2H). ¹H NMR (300 MHz, MeOD-d₄) (isomer D-

tartrate salt) δ 0.91 (t, J = 7.06 Hz, 3H), 1.28-1.85 (m, 8H), 2.44 (s, 3H), 2.68 (s, 3H), 2.94-2.99 (m, 3H), 3.14 (d, J = 15.82 Hz, 1H), 4.41 (s, 2H), 6.25-6.28 (m, 1H), 7.02-7.07 (m, 4H), 7.25-7.28 (m, 1H), 7.38 (d, J = 7.91 Hz, 2H). LCMS (12 minute method) $[M+H]^+$ = 351 @ Rt 5.6 min (100%).

5

Example 9D: 3-Butyl-3-(3-methylamino-propyl)-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (13Dd)

This was prepared from (3Df) (790 mg, 2.7 mmol) using the same synthetic sequence described above (3Dd to 13Db) to give 334 mg of the racemate. The racemate was separated into its individual enantiomers using chiral HPLC and each enantiomer was converted into its D-tartrate salt as described for (13Db). ^1H NMR (300 MHz, CDCl_3) (racemate) δ 0.87 (t, J = 6.97 Hz, 3H), 1.20-1.40 (m, 4H), 1.55-1.74 (m, 6H), 2.40 (s, 3H), 2.40 (s, 3H), 2.55 (t, J = 6.78 Hz, 3H), 2.91 (d, J = 15.63 Hz, 1H), 2.99 (d, J = 15.63 Hz, 1H), 6.28-6.31 (m, 1H), 6.93-7.00 (m, 2H), 7.02-7.06 (m, 2H), 7.14-7.16 (m, 1H), 7.29 (d, J = 8.07 Hz, 2H). ^1H NMR (300 MHz, MeOD-d_4) (isomer D-tartrate salt) δ 0.90 (t, J = 6.97 Hz, 3H), 1.20-1.85 (m, 10H), 2.44 (s, 3H), 2.68 (s, 3H), 2.94-2.99 (m, 3H), 3.14 (d, J = 15.82 Hz, 1H), 4.42 (s, 2H), 6.25-6.28 (m, 1H), 7.00-7.07 (m, 4H), 7.25-7.28 (m, 1H), 7.38 (d, J = 7.91 Hz, 2H). LCMS (12 minute method) $[M+H]^+$ = 365 @ Rt 5.9 min (100%).

20

Example 10D: 3-Isopropyl-3-(3-methylamino-propyl)-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (13De)

This was prepared from (3Dg) (806 mg, 2.89 mmol) using the same synthetic sequence described above (3Dd to 13Db) to give 307 mg of the racemate. ^1H NMR (300 MHz, CDCl_3) (racemate) δ ppm 0.92 (dd, J = 8.95, 6.88 Hz, 6H), 1.39-1.88 (m, 5H), 2.12-2.23 (m, 1H), 2.39 (s, 3H), 2.40 (s, 3H), 2.56 (t, J = 6.78 Hz, 2H), 2.94 (d, J = 15.92 Hz, 1H), 3.00 (d, J = 15.92 Hz, 1H), 6.28 (dd, J = 7.82, 1.04 Hz, 1H), 6.92-7.06 (m, 4H), 7.16 (dd, J = 6.97, 1.13 Hz, 1H), 7.29 (d, J = 7.91 Hz, 2H). LCMS (12 minute method) $[M+H]^+$ = 351 @ Rt 5.55 min (100%).

30

Example 11D: 6-Chloro-3-ethyl-3-(3-methylamino-propyl)-1-*p*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (13Df)

This was prepared from (1Dc) using the same synthetic sequence described above to give 205 mg of the racemate. The racemate was separated into its individual enantiomers using
5 chiral HPLC and each enantiomer was converted into its D-tartrate salt as described for (13Db). ¹H NMR (300 MHz, CDCl₃) (racemate) δ ppm 0.91 (t, J= 7.44 Hz, 3H), 1.50-1.75 (m, 6H), 2.15 (br, 1H), 2.40 (s, 3H), 2.41 (s, 3H), 2.55-2.64 (m, 2H), 2.85 (d, J= 16.01 Hz, 1H), 2.97 (d, J= 16.01 Hz, 1H), 6.23 (d, J= 8.85 Hz, 1H), 6.97 (dd, J= 8.67, 2.45 Hz, 1H), 7.02 (d, J= 8.29 Hz, 2H), 7.14 (d, J= 2.26 Hz, 1H), 7.29 (d, J= 8.10 Hz, 2H). ¹H NMR (300 MHz, MeOD-d₄) (isomer, D-tartrate salt) δ ppm 0.84 (t, J= 7.35 Hz, 3H), 1.40-1.75 (m, 6H), 2.32 (s, 3H), 2.57 (s, 3H), 2.80-2.92 (m, 3H), 3.01 (d, J= 16.20 Hz, 1H), 4.31 (s, 2H), 6.13 (d, J= 8.67 Hz, 1H), 6.92-6.98 (m, 3H), 7.19 (d, J= 2.26 Hz, 1H), 7.26 (d, J= 7.91 Hz, 2H). LCMS (12 minute method) [M+H]⁺ = 371/373 @Rt 5.75 min (100%).

15

Example 12D: 6-Chloro-1-(4-chloro-phenyl)-3-ethyl-3-(3-methylamino-propyl)-3,4-dihydro-1*H*-quinolin-2-one (13Dg)

This was prepared from (1Dc) using the same synthetic sequence described above to give 222 mg of the racemate, which was purified by preparative LCMS. ¹H NMR (300 MHz, CDCl₃) (racemate) δ ppm 0.84 (t, J= 7.44 Hz, 3H), 1.40-1.70 (m, 6H), 2.35 (br, 4H), 2.49-2.56 (m, 2H), 2.80 (d, J= 16.01 Hz, 1H), 2.90 (d, J= 16.01 Hz, 1H), 6.14 (d, J= 8.67 Hz, 1H), 6.93 (dd, J= 8.67, 2.26 Hz, 1H), 7.04 (ddd, J= 9.04, 2.83, 2.45 Hz, 2H), 7.09 (d, J= 2.26 Hz, 1H), 7.36-7.43 (m, 2H). LCMS (12 minute method) [M+H]⁺ = 391/393 @Rt 5.67 min (92%).

25

Scheme 3D - Preparation of intermediates

1-(4-Methoxy-benzyl)-3,4-dihydro-1*H*-quinolin-2-one (14D)

A 5 litre flange-neck flask equipped with an air stirrer and paddle, thermometer, nitrogen
30 bubbler and pressure equalising dropping funnel was charged with sodium hydride

(25.5g, 60% oil dispersion, 0.637 mol) and 40-60 pet. ether (100 ml). The mixture was stirred briefly and then allowed to settle under nitrogen. After decanting the supernatant liquid, the vessel was charged with dimethylformamide (2 litres). The well stirred suspension was cooled to 7-8°C using an external ice-bath. Then a soln of 3,4-dihydro-1H-quinolin-2-one (**1a**) (73.6g, 0.5 mole) in anhydrous dimethylformamide (500 ml) was added dropwise over 25 min. The mixture was stirred at 7-8°C for 30 min. then 4-methoxybenzyl chloride (102 g, 0.65 mole, 1.3 eq.) was added over 10 min. The reaction mixture was left to stir for 2 h. at <10°C then allowed to warm-up to room temperature and stirred overnight. The stirred reaction mixture was quenched with ice/water (2.5 litres) and cooled to 15 °C using an external ice-bath. The white solid was isolated by filtration and washed with water. After drying in vacuo at 40°C overnight the product was obtained (113.4g, 85%).

1-(4-Methoxy-benzyl)-3-methyl-3,4-dihydro-1H-quinolin-2-one (15D)

To a soln of (**14**) (20 g, 75 mmol) in anhydrous THF (400 mL) at -78°C under nitrogen was added LiHMDS (78.6 mL, 1M soln in hexanes, 78.6 mmol) dropwise over 10 min. The reaction mixture was left at -78°C for 30 min and then a solution of methyl iodide (5.13 mL, 83 mmol) in THF (5 mL) was added dropwise. The reaction mixture was warmed slowly to rt, quenched with water (50 mL) and extracted with ethyl acetate (400 mL). The organic layer was separated, dried over MgSO₄ and concentrated to give the product as a yellow solid (21 g, 100%) that was used directly in the next step.

3-Allyl-1-(4-methoxy-benzyl)-3-methyl-3,4-dihydro-1H-quinolin-2-one (16Db)

To a soln of (**15D**) (20.5 g, 73 mmol) in anhydrous THF (400 mL) at -78°C under nitrogen was added LiHMDS (80 mL, 1M soln in hexanes, 80 mmol) dropwise over 10 min. The reaction mixture was left at -78°C for 30 min and then a solution of allyl bromide (7.6 mL, 87 mmol) in THF (5 mL) was added dropwise. The reaction mixture was warmed slowly to rt, quenched with water (100 mL) and extracted with ethyl acetate (400 mL). The organic layer was separated, dried over MgSO₄ and concentrated to give the product as an orange oil (23.9 g, 100%) that was used directly in the next step.

3-(3-Hydroxy-propyl)-1-(4-methoxy-benzyl)-3-methyl-3,4,4a,8a-tetrahydro-1H-quinolin-2-one (17Db)

To a soln of (16Db) (23.9 g, 74 mmol) in anhydrous THF (400 mL) at 0°C under nitrogen was added 9-BBN (370 mL, 0.5M soln in THF, 185 mmol, 2.5 eq.) dropwise over 10 min. The reaction mixture was warmed to rt and left to stir overnight. The resultant yellow soln was cooled to 0°C and then quenched carefully with ethanol (95 mL), followed by aq. NaOH (60 mL, 3N soln). Finally, aq. H₂O₂ (60 mL, 37% soln) was added dropwise maintaining the internal reaction mixture temp between 5 and 10 °C. The reaction mixture was warmed to rt and then refluxed for 90 min. The reaction mixture was cooled to rt, poured into ethyl acetate and water and extracted. The organic layer was separated, dried over MgSO₄ and concentrated. The crude product was purified using automated chromatography (silica) (0 to 80% ethyl acetate/cyclohexane gradient) to provide the product as a clear oil (21.3 g, 84%).

1-(4-Methoxy-benzyl)-3-methyl-3-(3-methylamino-propyl)-3,4,4a,8a-tetrahydro-1H-quinolin-2-one (18Db)

To a soln of (17Db) (18 g, 53 mmol) and triethylamine (11.1 mL, 79 mmol) in anhydrous THF (450 mL) at 0°C under nitrogen was added dropwise a soln of methanesulfonyl chloride (4.52 mL, 58 mmol) in THF (50 mL). The reaction mixture was warmed to rt and stirred for 3 h. The reaction mixture was poured into ethyl acetate and water and extracted. The organic layer was separated, dried over MgSO₄ and concentrated. The crude mesylate (22 g, 99%) was dissolved in ethanol (500 mL) and aqueous 40% methylamine (200 mL) and heated at 65°C under nitrogen for 2 h. The reaction mixture was cooled, concentrated and then extracted with ethyl acetate (300 mL). The organic layer was washed with water, brine, dried over MgSO₄ and concentrated to give the crude product (17.8 g, 96%).

Methyl-[3-(3-methyl-2-oxo-1,2,3,4,4a,8a-hexahydro-quinolin-3-yl)-propyl]-carbamic acid *tert*-butyl ester (19Db)

A mixture of (18Db) (17.8 g, 50.5 mmol) and anisole (5.5 mL, 50.5 mmol) in trifluoroacetic acid (250 mL) was heated at 65°C under nitrogen for 2 h. The reaction

mixture was concentrated under vacuo and the residue was dissolved in methanol (10 mL). The methanol soln was applied to an SCX-2 column (300 g, pre-washed with methanol) and the column washed with methanol (approx 1 litre) until the soln became colourless. The product was eluted with 2N NH₃ in methanol (500 mL) and the basic soln
5 was concentrated to provide 3-Methyl-3-(3-methylamino-propyl)-3,4-dihydro-1H-quinolin-2-one (9 g, 77%). To a soln of this amine (8.6 g, 37 mmol) in anhydrous THF (350 mL) at 0°C was added a soln of di-*tert*-butyl dicarbonate (8.34 g, 97%, 50.5 mmol) in THF (20 mL) dropwise. The reaction mixture was warmed to rt and stirred for 3 h. The reaction mixture was poured into ethyl acetate (400 mL) and water (200 mL) and
10 extracted. The organic layer was separated, dried over MgSO₄ and concentrated to give the product as a yellow solid (12.26 g, 100%). This material was used without further purification.

Methyl-[3-(2-oxo-1,2,3,4-tetrahydro-quinolin-3-yl)-propyl]-carbamic acid *tert*-butyl ester (19Da)
15

This was prepared from (14D) using the same synthetic sequence described above.

[3-(6-Chloro-1,2,3,4-tetrahydro-quinolin-3-yl)-propyl]-methyl-carbamic acid *tert*-butyl ester (20Da)
20 To a soln of (19Da) (2.75 g, 8.6 mmol) in anhydrous DMF (25 mL) at 0°C was added dropwise a soln of N-chlorosuccinimide (1.17 g, 8.7 mmol) in anhydrous DMF (3 mL). The reaction mixture was warmed to rt, stirred overnight and then poured into ethyl acetate (100 mL) and water (50 mL) and extracted. The organic layer was separated, dried over MgSO₄ and concentrated to provide the product as a yellow oil 3 g, 98%) that was
25 used without further purification.

Scheme 3D - Examples

Example 13D: 3-(3-Methylamino-propyl)-1-*p*-tolyl-3,4-dihydro-1H-quinolin-2-one (21Da)
30

A stirred mixture of (19Da) (100 mg, 0.31 mmol), K₂CO₃ (92 mg, 0.66 mmol), *trans*-cyclohexane-1,2-diamine (8 μ L, 0.06 mmol) and 4-bromotoluene (162 mg, 0.94 mmol) in 1,4-dioxane (0.5 mL) was heated under a nitrogen atmosphere at 125°C for 5 min to deoxygenate the reaction mixture. Copper (I) iodide (12 mg, 0.06 mmol) was added in one portion and the reaction mixture was refluxed overnight at 125°C. After cooling to rt, the reaction mixture was poured into ethyl acetate (100 mL) and extracted with water. The organic layer was separated, dried over MgSO₄ and concentrated. The crude product was purified using automated chromatography (silica) (0 to 80% ethyl acetate/cyclohexane gradient) to provide the Boc protected product (70 mg, 54%). To a soln of this material (70 mg, 0.17 mol) in DCM (2 mL), was added trifluoroacetic acid (197 μ L, 2.55 mmol, 15 eq.). The reaction mixture was left to stir at room temperature for 90 min, concentrated under vacuo poured into ethyl acetate (50 mL) and aq. NaHCO₃ (20 mL) and extracted. The organic layer was separated, dried over MgSO₄, concentrated and the crude product was purified by SCX-2 to provide the racemate (40 mg, 75%). The racemate was separated into its individual enantiomers using chiral HPLC. ¹H NMR (300 MHz, CDCl₃) (racemate) δ 1.49-1.77 (m, 3H), 1.86-1.96 (m, 1H), 2.34 (bs, 1H), 2.40 (s, 3H), 2.43 (s, 3H), 2.61-2.66 (t, J= 6.88 Hz, 2H), 2.68-2.78 (m, 1H), 2.83-2.90 (m, 1H), 3.09-3.17 (m, 1H), 6.36 (dd, J= 7.7 Hz, 1.0 Hz, 1H), 6.94-7.03 (m, 2H), 7.08 (d, J= 8.2 Hz, 2H), 7.13-7.17 (m, 1H), 7.29 (d, J= 8.1 Hz, 2H); ¹H NMR (300 MHz, MeOD-d₄) (isomer, D-tartrate salt) δ 1.64 (bs, 1H), 1.89 (bs, 3H), 2.41(s, 3H), 2.70 (s, 3H), 2.75-2.87 (m, 1H), 2.91-3.06 (m, 3H), 3.20 (dd, J= 5.9, 15.26 Hz, 1H), 4.45 (s, 2H), 6.32-6.35 (m, 1H), 7.00-7.12 (m, 4H), 7.28-7.30 (m, 1H), 7.37 (d, J= 8.1 Hz, 2H). LCMS (12 minute method) [M+H]⁺ = 309 @ Rt 4.7 min (100%).

25

Example 14D: 6-Chloro-3-(3-methylamino-propyl)-1-*p*-tolyl-3,4-dihydro-1H-quinolin-2-one (21Dn)

This was prepared from (20Da) (132 mg, 0.29 mmol) using the same methods described for (21Da) to provide the racemate (86 mg). ¹H NMR (300 MHz, CDCl₃) (racemate & isomer) δ 1.50-1.57 (m, 1H), 1.62-1.90 (m, 3H), 2.34 (s, 3H), 2.41 (s, 3H), 2.63-2.82 (m, 5H), 3.00-3.07 (m, 1H), 6.22 (d, J= 8.6 Hz, 1H), 6.92 (dd, J= 2.45, 8.66 Hz, 1H), 6.99 (d,

J= 8.1 Hz, 2H), 7.11 (d, J= 2.25 Hz, 1H), 7.23 (d, J= 8.1 Hz, 2H). LCMS (12 minute method) $[M+H]^+ = 343/345$ @ Rt 5.2 min (96%).

Example 15D: 1-(3-Fluorophenyl)-3-(3-methylamino-propyl)-3,4-dihydro-1H-

5 **quinolin-2-one (21Db)**

This was prepared from (19Da) (200 mg, 0.63 mmol) using the same two-step procedure described for (21Da) to provide the racemate (83 mg). ^1H NMR (300 MHz, CDCl_3) (racemate) δ 1.60-1.70 (m, 1H), 1.92 (br, 3H), 2.64 (bs, 3H), 2.72-2.74 (m, 1H), 2.86-3.09 (m, 4H), 6.35 (dd, J= 7.72, 1.510 Hz, 1H), 6.94-7.23 (m, 6H), 7.43-7.51 (m, 1H). LCMS
10 (12 minute method) $[M+H]^+ = 313$ @ Rt 4.4 min (100%).

Example 16D: 1-(4-Chlorophenyl)-3-(3-methylamino-propyl)-3,4-dihydro-1H-

quinolin-2-one (21Dc)

This was prepared from (19Da) (122 mg, 0.38 mmol) using the same two-step procedure
15 described for (21Da) to provide the crude product, which was purified by SCX-2 to give the racemate (70 mg). ^1H NMR (300 MHz, CDCl_3) (racemate) δ 1.49-1.73 (m, 3H), 1.89 (m, 2H), 2.43 (s, 3H), 2.62 (t, J= 6.79, 7.15 Hz, 2H), 2.68-2.78 (m, 1H), 2.83-2.93 (m, 1H), 3.14 (dd, J= 15.43, 5.37 Hz, 1H), 6.34 (dd, J= 7.73, 1.14 Hz, 1H), 6.96-7.09 (m, 2H), 7.14-7.21 (m, 3H), 7.45-7.48 (m, 2H). LCMS (12 minute method) $[M+H]^+ = 329/331$ @
20 Rt 5.1 min (90%).

Example 17D: 1-(3,4-Dichlorophenyl)-3-(3-methylamino-propyl)-3,4-dihydro-1H-

quinolin-2-one (21Dd)

This was prepared from (19Da) (150 mg, 0.47 mmol) using the same two-step procedure
25 described for (21Da) to provide the crude product, which was purified by SCX-2 to give the racemate (111 mg). ^1H NMR (300 MHz, CDCl_3) (racemate) δ 1.49-1.75 (m, 3H), 1.83 (bs, 1H), 1.85-1.97 (m, 1H), 2.43 (s, 3H), 2.63 (t, J= 13.56, 6.59 Hz, 2H), 2.68-2.77 (m, 1H), 2.83-2.94 (m, 1H), 3.13 (dd, J= 15.45, 5.28 Hz, 1H), 6.36 (dd, J= 7.73, 0.93 Hz, 1H), 6.99-7.11 (m, 3H), 7.20-7.21 (m, 1H), 7.35 (d, J= 2.26 Hz, 1H), 7.57 (d, J= 8.48 Hz, 1H).
30 LCMS (12 minute method) $[M+H]^+ = 363/365$ @Rt 5.4 min (92%).

Example 18D: 1-(3-Chlorophenyl)-3-(3-methylamino-propyl)-3,4-dihydro-1H-quinolin-2-one (21De)

This was prepared from (19Da) (200 mg, 0.63 mmol) using the same two-step procedure described for (21Da) to provide the crude product, which was purified by SCX-2 to give
5 the racemate (138 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ 1.50-1.77 (m, 3H), 1.89-1.96 (m, 2H), 2.44 (s, 3H), 2.64 (t, J= 6.89 Hz, 2H), 2.69-2.78 (m, 1H), 2.84-2.93 (m, 1H), 3.10-3.17 (m, 1H), 6.33-6.36 (m, 1H), 6.97-7.10 (m, 2H), 7.11-7.15 (m, 1H), 7.21-7.24 (m, 2H), 7.37-7.47 (m, 2H). LCMS (12 minute method) [M+H]⁺ = 329/331 @ Rt 5.01 min (90%).

Example 19D: 1-(4-Fluorophenyl)-3-(3-methylamino-propyl)-3,4-dihydro-1H-quinolin-2-one (21Df)

This was prepared from (19Da) (200 mg, 0.63 mmol) using the same two-step procedure described for (21Da) to provide the crude product, which was purified by SCX-2 to give
15 the racemate (48 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ 1.26-1.28 (m, 1H), 1.92 (m, 2H), 2.63 (bs, 1H), 2.72 (m, 1H), 2.85-3.08 (m, 2H), 3.48-3.51 (m, 5H), 6.32-6.34 (d, J= 7.91 Hz, 1H), 7.01-7.70 (m, 2H), 7.16-7.19 (d, J= 7.16 Hz, 5H), 9.46 (bs, 1H). LCMS (12 minute method) [M+H]⁺ = 313 @ Rt 4.5 min (100%).

Example 20D: 1-(4-Ethylphenyl)-3-(3-methylamino-propyl)-3,4-dihydro-1H-quinolin-2-one (21Dg)

This was prepared from (19Da) (148 mg, 0.46 mmol) using the same two-step procedure described for (21Da) to provide the racemate (61 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ 1.25-1.30 (m, 1H), 1.52-1.67(m, 1H), 1.69-1.80 (m, 2H), 1.87-1.98 (m, 1H),
25 2.46 (s, 3H), 2.67-2.92 (m, 9H), 3.11-3.16 (m, 1H), 6.34-6.37 (m, 1H), 6.94-7.06 (m, 2H), 7.09-7.11 (d, J= 8.1 Hz, 2H), 7.17-7.20 (d, J= 7.35 Hz, 1H), 7.30-7.33 (d, J= 8.28 Hz, 2H). LCMS (12 minute method) [M+H]⁺ = 323 @ Rt 5.4 min (98%).

Example 21D: 3-Methyl-3-(3-methylamino-propyl)-1-p-tolyl-3,4-dihydro-1H-quinolin-2-one (21Dh)

This was prepared from (19Db) (806 mg, 2.89 mmol) using the same methods described for (21Da) to provide the racemate. The racemate was separated into its individual enantiomers using chiral HPLC. ¹H NMR (300 MHz, CDCl₃) (racemate & isomer) δ 1.24 (s, 3H), 1.60-1.65 (m, 4H), 2.40 (s, 3H), 2.43 (s, 3H), 2.60-2.65 (m, 2H), 2.87 (d, J= 15.73 Hz, 1H), 2.98 (d, J= 15.73 Hz, 1H), 3.46 (br, 1H), 6.30 (dd, J= 7.91, 1.13 Hz, 1H), 6.90-7.05 (m, 2H), 7.05 (d, J= 8.29 Hz, 2H), 7.10-7.20 (m, 1H), 7.29 (d, J= 7.91 Hz, 2H). LCMS (12 minute method) [M+H]⁺ = 323 @Rt 5.06 min (100%).

Example 22D: 1-(4-Chlorophenyl)-3-methyl-3-(3-methylamino-propyl)-3,4-dihydro-1H-quinolin-2-one (21Di)

This was prepared from (19Db) (100 mg, 0.30 mmol) using the same methods described for (21Da) to provide the racemate (97 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ ppm 1.25 (s, 3H), 1.55-1.65 (m, 4H), 2.41 (s, 3H), 2.58 (m, 2H), 2.89 (d, J= 15.82 Hz, 1H), 2.98 (d, J= 15.82 Hz, 1H), 3.12 (br, 1H), 6.29 (dd, J= 7.91, 0.94 Hz, 1H), 6.95-7.10 (m, 2H), 7.14 (d, J= 8.67 Hz, 2H), 7.15 (m, 1H), 7.45 (d, J= 8.67 Hz, 2H). LCMS (12 minute method) [M+H]⁺ = 343/345 @Rt 5.09 min (100%).

Example 23D: 1-(3,4-Difluorophenyl)-3-methyl-3-(3-methylamino-propyl)-3,4-dihydro-1H-quinolin-2-one (21Dj)

This was prepared from (19Db) (100 mg, 0.30 mmol) using the same two-step procedure described for (21Da) to provide the crude product, which was purified by SCX-2 to give the racemate (100 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ ppm 1.25 (s, 3H), 1.55-1.65 (m, 4H), 2.41 (s, 3H), 2.50-2.60 (m, 2H), 2.89 (d, J= 15.45 Hz, 1H), 2.90 (s, 1H), 2.98 (d, J= 15.45 Hz, 1H), 6.30 (dd, J= 7.91, 1.13 Hz, 1H), 6.90-7.10 (m, 4H), 7.18 (dd, J= 7.16, 1.32 Hz, 1H), 7.22-7.35 (m, 1H). LCMS (12 minute method) [M+H]⁺ = 345 @Rt 4.85 min (97%).

Example 24D: 3-Methyl-3-(3-methylamino-propyl)-1-*m*-tolyl-3,4-dihydro-1*H*-quinolin-2-one (21Dk)

This was prepared from (19Db) (100 mg, 0.30 mmol) using the same two-step procedure described for (21Da) to provide the crude product, which was purified by SCX-2 to give the racemate (90 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ ppm 1.26 (s, 3H), 1.50-1.70 (m, 4H), 1.75 (s, 1H), 2.38 (s, 3H), 2.39 (s, 3H), 2.50-2.60 (m, 2H), 2.89 (d, J= 15.64 Hz, 1H), 2.98 (d, J= 15.64 Hz, 1H), 6.30 (dd, J= 7.82, 1.04 Hz, 1H), 6.90-7.07 (m, 4H), 7.18 (dd, J= 13.66, 7.63 Hz, 2H), 7.37 (t, J= 7.63 Hz, 1H). LCMS (12 minute method) [M+H]⁺ = 323 @Rt 5.09 min (98%).

10

Example 25D: 1-(3,5-Difluorophenyl)-3-methyl-3-(3-methylamino-propyl)-3,4-dihydro-1*H*-quinolin-2-one (21Dl)

This was prepared from (19Db) (100 mg, 0.30 mmol) using the same two-step procedure described for (21Da) to provide the crude product, which was purified by SCX-2 to give the racemate (95 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ ppm 1.26 (s, 3H), 1.50-1.65 (m, 4H), 2.40 (s, 3H), 2.50-2.60 (m, 2H), 2.82 (br, 1H), 2.89 (d, J= 15.82 Hz, 1H), 2.97 (d, J= 15.82 Hz, 1H), 6.34 (dd, J= 8.01, 1.04 Hz, 1H), 6.74-6.83 (m, 2H), 6.83-6.92 (m, 1H), 6.97-7.13 (m, 2H), 7.19 (dd, J= 7.06, 1.22 Hz, 1H). LCMS (12 minute method) [M+H]⁺ = 345 @ Rt 4.87 min, (97%).

20

Example 26D: 6-Chloro-3-(3-methylamino-propyl)-1-phenyl-3,4-dihydro-1*H*-quinolin-2-one (21Dm)

This was prepared from (20Da) (285 mg, 0.8 mmol) using the same two-step procedure described for (21Da) to provide the crude product, which was purified by preparative LCMS to give the racemate (62 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ 1.49-1.76 (m, 3H), 1.86-1.95 (m, 1H), 2.33 (bs, 1H), 2.44 (s, 3H), 2.61-2.95 (m, 4H), 3.09-3.16 (m, 1H), 6.24-6.27 (d, J= 8.67 Hz, 1H), 6.99 (dd, J= 8.67, 2.26 Hz, 1H), 7.17-7.19 (m, 3H), 7.39-7.44 (m, 1H), 7.47-7.52 (m, 2H). LCMS (12 minute method) [M+H]⁺ = 329/331 @ Rt 5.04 min (93%).

30

Example 27D: 6-Chloro-1-(4-chlorophenyl)-3-(3-methylamino-propyl)-3,4-dihydro-1H-quinolin-2-one (21Do)

This was prepared from (20Da) (160 mg, 0.45 mmol) using the same two-step procedure described for (21Da) to provide the crude product, which was purified by preparative LCMS to give the racemate (52 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ 1.57-1.67 (m, 1H), 1.73-1.75 (m, 2H), 1.87-1.9 (m, 1H), 2.47 (s, 2H), 2.64 (s, 1H), 2.68-2.73 (m, 2H), 2.81-2.89 (m, 1H), 3.07-3.13 (m, 3H), 6.27 (d, J= 8.48 Hz, 1H), 7.02 (d, J= 8.48 Hz, 1H), 7.14 (d, J= 8.29 Hz, 2H), 7.19 (s, 1H), 7.47 (d, J= 8.29 Hz, 2H). LCMS (12 minute method) [M+H]⁺ = 363/365 @ Rt 5.4 min (72%).

Example 28D: 6-Chloro-3-methyl-3-(3-methylamino-propyl)-1-p-tolyl-3,4-dihydro-1H-quinolin-2-one (21Dp)

This was prepared from (20Db) (490 mg, 1.34 mmol) using the same methods described for (21Da) to provide the racemate (470 mg). The racemate was separated into its individual enantiomers using chiral HPLC. ¹H NMR (300 MHz, CDCl₃) (racemate) δ 1.25 (s, 3H), 1.50-1.65 (m, 4H), 2.39 (s, 3H), 2.40 (s, 3H), 2.50-2.60 (m, 3H), 2.86 (d, J= 16.01 Hz, 1H), 2.94 (d, J= 16.01 Hz, 1H), 6.24 (d, J= 8.67 Hz, 1H), 6.97 (dd, J= 8.76, 2.35 Hz, 1H), 7.03 (d, J= 8.10 Hz, 2H), 7.14 (d, J= 2.26 Hz, 1H), 7.29 (d, J= 7.91 Hz, 2H); ¹H NMR (300 MHz, MeOD-d₄) (isomer hemi-D-tartrate salt) δ 1.15 (s, 3H), 1.50-1.75 (m, 4H), 2.32 (s, 3H), 2.51 (s, 3H), 2.78 (br, 2H), 2.84 (d, J= 16.20 Hz, 1H), 2.98 (m, 1H), 3.15-3.25 (m, 2H), 4.22 (s, 1H), 6.14 (d, J= 8.85 Hz, 1H), 6.90-6.70 (m, 3H), 7.19 (d, J= 2.26 Hz, 1H), 7.25 (d, J= 7.91 Hz, 2H). LCMS (12 minute method) [M+H]⁺ = 357/359 @Rt 5.43 min (100%).

Example 29D: 6-Chloro-1-(4-chlorophenyl)-3-methyl-3-(3-methylamino-propyl)-3,4-dihydro-1H-quinolin-2-one (21Dq)

This was prepared from (20Db) (490 mg, 1.34 mmol) using the same methods described for (21Da) to provide the racemate (425 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ ppm 1.25 (s, 3H), 1.50-1.65 (m, 4H), 2.39 (s, 3H), 2.40 (br, 1H), 2.50-2.60 (m, 2H), 2.87 (d, J= 16.20 Hz, 1H), 2.95 (d, J= 16.20 Hz, 1H), 6.23 (d, J= 8.85 Hz, 1H), 7.00 (dd, J=

8.57, 2.35 Hz, 1H), 7.05-7.20 (m, 3H), 7.40-7.50 (m, 2H). LCMS (12 minute method) $[M+H]^+ = 377/379$ @Rt 5.26 min (94%).

Example 30D: 3-Methyl-3-(3-methylamino-propyl)-1-thiophen-2-yl-3,4-dihydro-1H-quinolin-2-one (22Da)

This was prepared from (19Db) (200 mg, 0.60 mmol) using the same two-step procedure described for (21Da) to provide the crude product, which was purified by SCX-2 to give the racemate (125 mg). ^1H NMR (300 MHz, CDCl_3) (racemate) δ ppm 1.25 (s, 3H), 1.50-1.65 (m, 4H), 2.39 (s, 3H), 2.50-2.60 (br, 2H), 2.88 (d, $J = 16.20$ Hz, 1H), 2.97 (d, $J = 16.20$ Hz, 1H), 3.17 (br, 1H), 6.58 (dd, $J = 8.01, 0.85$ Hz, 1H), 6.89 (dd, $J = 3.58, 1.32$ Hz, 1H), 6.95-7.15 (m, 3H), 7.16 (d, $J = 7.16$ Hz, 1H), 7.32 (dd, $J = 5.65, 1.32$ Hz, 1H). LCMS (12 minute method) $[M+H]^+ = 315$ @Rt 4.35 min (98%).

Example 31D: 3-Methyl-3-(3-methylamino-propyl)-1-thiophen-3-yl-3,4-dihydro-1H-quinolin-2-one (22Db)

This was prepared from (19Db) (200 mg, 0.60 mmol) using the same two-step procedure described for (21Da) to provide the crude product, which was purified by SCX-2-2 to give the racemate (128 mg). ^1H NMR (300 MHz, CDCl_3) δ 1.24 (s, 3H), 1.50-1.65 (m, 4H), 2.40 (s, 3H), 2.50-2.60 (m, 2H), 2.87 (d, $J = 15.82$ Hz, 1H), 2.96 (d, $J = 15.82$ Hz, 1H), 3.07 (br, 1H), 6.45 (dd, $J = 8.10, 0.94$ Hz, 1H), 6.92 (dd, $J = 5.09, 1.32$ Hz, 1H), 6.98 (td, $J = 7.35, 1.13$ Hz, 1H), 7.07 (td, $J = 7.77, 1.60$ Hz, 1H), 7.16 (d, $J = 7.35$ Hz, 1H), 7.22 (dd, $J = 3.20, 1.32$ Hz, 1H), 7.41 (dd, $J = 5.09, 3.20$ Hz, 1H). LCMS (12 minute method) $[M+H]^+ = 315$ @Rt 4.29 min (100%).

Scheme 4D - Preparation of intermediates

{3-[1-(4-Methoxy-benzyl)-3-methyl-2-oxo-6-phenyl-1,2,3,4-tetrahydro-quinolin-3-yl]-propyl}-methyl-carbamic acid *tert*-butyl ester (23D)

Step (i)

Sodium hydride (340 mg, 60% dispersion in mineral oil, 8.55 mmol, 1.3 eq.) was added portionwise to a soln of (20Dc) (2.7 g, 6.57 mmol) in DMF (40 mL) at 0°C. The reaction

mixture was left for 30 min at this temperature and then 4-methoxybenzyl chloride (1.16 mL, 8.55 mmol, 1.3 eq.) in DMF (1 mL) was added dropwise over 10 min. The reaction mixture was warmed to rt slowly and after 1 h was poured into ethyl acetate (200 mL) and extracted with water (3 x 50 mL). The organic layer was separated, dried over
5 MgSO₄ and concentrated under vacuo. The crude product was purified using automated chromatography (silica) (0 to 80% ethyl acetate/cyclohexane gradient) to provide the 4-methoxybenzyl protected 6-bromo precursor (2.2 g, 63%).

Step (ii)

The product from Step (i) (100 mg, 0.23 mmol), phenylboronic acid (85 mg, 0.70 mmol,
10 3 eq.), K₂CO₃ (138 mg, 1 mmol, 4.3 eq.) and Pd(PPh₃)₄ (11 mg, 0.009 mmol, 0.04 eq.) were suspended in ethanol (1 mL) and water (0.6 mL). The reaction mixture was heated at 80°C overnight, cooled to rt and filtered through celite. The filtrate was poured into ethyl acetate (100 mL) and water (50 mL) and extracted. The organic layer was separated, dried over MgSO₄ and concentrated to provide the product (**23D**) (120 mg, 98%) that was
15 used without further purification.

Methyl-[3-(3-methyl-2-oxo-6-phenyl-1,2,3,4-tetrahydro-quinolin-3-yl)-propyl]-carbamic acid *tert*-butyl ester

Step (iii) & (iv)

20 A mixture of (**23D**) (120 mg, 0.23 mmol) and anisole (25 µL, 0.23 mmol) in trifluoroacetic acid (2.3 mL) was heated at 65°C under nitrogen for 4 h. The reaction mixture was concentrated under vacuo and the residue was dissolved in methanol (2 mL). The methanol soln was applied to an SCX-2 column (5g) and the column washed with methanol (50 mL). The product was eluted with 2N Et₃N in methanol (50 mL) and the
25 basic soln was concentrated to provide 3-Methyl-3-(3-methylamino-propyl)-6-phenyl-3,4-dihydro-1*H*-quinolin-2-one (72 mg, 100%). To a soln of this amine (72 mg, 0.23 mmol) in anhydrous THF (2 mL) at 0°C was added di-*tert*-butyl dicarbonate (53 mg, 97%, 0.24 mmol) in one portion. The reaction mixture was warmed to rt and stirred for 3 h. The reaction mixture was poured into ethyl acetate (25 mL) and water (10 mL) and
30 extracted. The organic layer was separated, dried over MgSO₄ and concentrated to give

the Boc protected precursor (95 mg, 100%). This material was used without further purification.

Scheme 4D - Examples

5

Example 32D: 3-Methyl-3-(3-methylamino-propyl)-6-phenyl-1-*p*-tolyl-3,4-dihydro-1H-quinolin-2-one (24D)

This was prepared from the above Boc protected precursor (95 mg, 0.23 mmol) using the same two-step procedure described above (19Da to 21Da) to provide the crude product, which was purified by SCX-2 to give the racemate (53 mg). ¹H NMR (300 MHz, CDCl₃) (racemate) δ 1.29 (s, 3H), 1.50-1.70 (m, 4H), 2.42 (s, 6H), 2.55-2.65 (m, 2H), 2.94 (d, J= 15.64 Hz, 1H), 3.04 (d, J= 15.64 Hz, 1H), 3.18 (br, 1H), 6.38 (d, J= 8.29 Hz, 1H), 7.09 (d, J= 8.10 Hz, 2H), 7.29 (m, 4H), 7.41 (m, 3H), 7.54 (m, 2H). LCMS (12 minute method) [M+H]⁺ = 399 @Rt 6.06 min (100%).

15

The following examples illustrate compounds of of Formulae (IE) above and methods for their preparation.

Preparation of Intermediates

20

1,1-Dimethylethyl (3S)-3-aminopyrrolidine-1-carboxylate

a) 1,1-Dimethylethyl (3R)-3-hydroxypyrrolidine-1-carboxylate

Solid *tert*-butyldicarbonate (38.8g, 178mmol) was added in portions over 15 minutes to a stirred solution of (3R)-pyrrolidin-3-ol hydrochloride (20g, 162mmol), triethylamine (24.8mL, 178mmol) and 4-(dimethylamino)-pyridine (20mg) in dry dichloromethane (300mL). After stirring for 2 hours at room temperature, the mixture was washed with aqueous citric acid, then brine. The organic extracts were dried (MgSO₄), filtered and evaporated *in vacuo* to give an oil. This was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (20:80 to 60:40), to give the title compound as a solid.

30

b) 1,1-Dimethylethyl (3*R*)-3-[(methylsulfonyl)oxy]-pyrrolidine-1-carboxylate

Methanesulfonyl chloride (5.26mL, 68mmol) was added dropwise over 5 minutes to a stirred solution of 1,1-dimethylethyl (3*R*)-3-hydroxypyrrolidine-1-carboxylate (10.6g, 56.7mmol) and triethylamine (11.8mL, 85mmol) in dichloromethane (250mL) at -10°C. After stirring for 1 hour at 0°C, the reaction was quenched by addition of water. The organic phase was washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo* to give an oil. This was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (25:75 to 50:50), to give the title compound as an oil.

10

c) 1,1-Dimethylethyl (3*S*)-3-azidopyrrolidine-1-carboxylate

Sodium azide (4.4g, 67.4mmol) was added to a solution of 1,1-dimethylethyl (3*R*)-3-[(methylsulfonyl)oxy]-pyrrolidine-1-carboxylate (14.3g, 54mmol) in dry dimethylformamide (75mL) and the resultant suspension heated at 65°C for 8 hours.

After cooling to room temperature, the reaction mixture was diluted with water and extracted into diethyl ether. The organic phase was washed two further times with water, then brine. The organic extracts were dried (MgSO₄), filtered and evaporated *in vacuo* to give an oil. This was purified by flash chromatography on silica, eluting with diethyl ether/cyclohexane (20:80 to 40:60), to give the title compound as an oil.

20

d) 1,1-Dimethylethyl (3*S*)-3-aminopyrrolidine-1-carboxylate

A mixture of 1,1-dimethylethyl (3*S*)-3-azidopyrrolidine-1-carboxylate (9.0g, 2.97mmol) and 5% palladium-on-carbon (0.70g) in methanol (150mL) was hydrogenated in a Parr apparatus at 65 p.s.i. for 4 hours. The catalyst was removed by filtration through Celite and the solvent evaporated *in vacuo* to give an oil. The resultant title compound was used in subsequent reactions without further purification.

25

1,1-Dimethylethyl (3*R*)-3-aminopyrrolidine-1-carboxylate was similarly prepared as described above, from (3*S*)-pyrrolidin-3-ol.

30

1,1-Dimethylethyl (3*S*)-3-[(1-methylethyl)amino]-pyrrolidine-1-carboxylate

A mixture of 1,1-dimethylethyl (3*S*)-3-aminopyrrolidine-1-carboxylate (3.0g) and 5% palladium-on-carbon (0.35g) in methanol (75mL) and acetone (15mL) was hydrogenated in a Parr apparatus at 65 p.s.i. for 3 hours. The catalyst was removed by filtration through
 5 Celite and the solvent evaporated *in vacuo* to give an oil. The resultant title compound was used in subsequent reactions without further purification.

¹H NMR (300 MHz, CDCl₃) δ_H: 1.11-1.19 (m, 6H), 1.45 (s, 9H), 1.55-1.75 (m, 1H), 2.01-2.15 (m, 1H), 2.80-2.92 (m, 1H), 2.93-3.05 (m, 1H), 3.25-3.70 (m, 4H).

10 The following secondary amines were similarly prepared by reductive alkylation of 1,1-dimethylethyl (3*S*)-3-aminopyrrolidine-1-carboxylate with the appropriate aldehyde or ketone:

1,1-Dimethylethyl (3*S*)-3-(cyclopentylamino)pyrrolidine-1-carboxylate

1,1-Dimethylethyl (3*S*)-3-[(cyclohexylmethyl)amino]-pyrrolidine-1-carboxylate

15

1,1-Dimethylethyl (3*S*)-3-([2-(trifluoromethyl)phenyl]-methyl)amino)pyrrolidine-1-carboxylate

Method A

20 a) (3*S*)-*N*-{(E)-[2-(Trifluoromethyl)phenyl]methylidene}-pyrrolidin-3-amine
 3(*S*)-Pyrrolidin-3-amine (0.45g, 5.2mmol) and trifluoromethylbenzaldehyde (0.87g, 5.0mmol), a crystal of 4-toluenesulphonic acid and toluene were refluxed with stirring for one day, using a Dean and Stark apparatus. The solution was evaporated *in vacuo* to give the title compound as a brown oil (M+H = 243).

25

b) 1,1-Dimethylethyl (3*S*)-3-({(E)-[2-(trifluoromethyl)-phenyl]methylidene}amino)pyrrolidine-1-carboxylate

(3*S*)-*N*-{(E)-[2-(Trifluoromethyl)phenyl]methylidene}-pyrrolidin-3-amine (1.21g, 5mmol) was dissolved in dichloromethane (50 mL), and di-*tert*-butyl dicarbonate (1.1g,
 30 5.05mmol) followed by DMAP (60mg, 0.5mmol) was added. After stirring under

nitrogen for 4 hours, the solution was evaporated *in vacuo* to give the title compound as a brown oil (M + H = 343).

- 5 c) 1,1-Dimethylethyl (3*S*)-3-({[2-(trifluoromethyl)-phenyl]methyl}amino)pyrrolidine-1-carboxylate

1,1-Dimethylethyl (3*S*)-3-({(*E*)-[2-(trifluoromethyl)-phenyl]methylidene}amino)pyrrolidine-1-carboxylate (1.71 g, 5mmol) was hydrogenated in the presence of 5% palladium on carbon (250mg) at 65psi in ethanol (60mL). After 3.5
10 hours, the catalyst was filtered off and the filtrate evaporated *in vacuo* to give an oil. The oil was purified by automated flash chromatography over silica, eluting with 10% ethyl acetate in cyclohexane (10:90 to 50:50), to give the title compound as a colourless oil (1.0g, 58%; M + H = 345).

15 Method B

- a) (3*S*)-*N*-{[2-(Trifluoromethyl)phenyl]methyl}pyrrolidin-3-amine

A mixture of 3(*S*)-pyrrolidin-3-amine (4g, 46.5mmol), 2-trifluoromethylbenzaldehyde (9.1g, 46.5mmol), 5% palladium on carbon (0.4g) and ethanol (150mL) was
20 hydrogenated at 60psi for 3 hours using a Parr hydrogenator. The catalyst was filtered off and the filtrate evaporated *in vacuo* to give the title compound as an oil. MS: [M+H] = 245.

- 25 b) 1,1-Dimethylethyl (3*S*)-3-({[2-(trifluoromethyl)-phenyl]methyl}amino)pyrrolidine-1-carboxylate

(3*S*)-*N*-{[2-(Trifluoromethyl)phenyl]methyl}pyrrolidin-3-amine (12g, 49.2mmol) was dissolved in dichloromethane (120 mL), then di-*tert*-butyl dicarbonate (10.7g, 49.2mmol) and DMAP (40mg, 0.33mmol) were added. After stirring under nitrogen for 1 day, the solution was evaporated *in vacuo* to give an oil. The oil was purified by automated flash

chromatography over silica, eluting with ethyl acetate in cyclohexane (0:100 to 40:60), to give the title compound as a colourless oil.

MS: $[M+H] = 345$.

5 **1,1-Dimethylethyl (3S)-3-({[4-fluoro-2-(trifluoromethyl)-phenyl]methyl}amino)pyrrolidine-1-carboxylate**

1,1-Dimethylethyl (3S)-3-aminopiperidine-1-carboxylate (5g) and 4-fluoro-2-(trifluoromethyl)benzaldehyde (5.15g, 26.8mmol) were allowed to stir in methanol for 16h
10 at room temperature. Sodium borohydride (1.62g, 26.8mmol) was then added portionwise. The resulting solution was further stirred for 2 h at room temperature. The solvent was evaporated *in vacuo*, water was added, and the solution extracted with dichloromethane. The organic extracts were absorbed onto a methanol washed cationic ion exchange resin (Isolute™ SCX-2). The basic components were recovered from the
15 column by elution with 7N ammonia in methanol. The resultant solution was concentrated *in vacuo* to yield the desired compound as an oil. This was further purified by column chromatography on silica gel, eluting with ethyl acetate/iso-hexane (0:100 to 40:60). The title compound was used in subsequent reactions without further purification.

^1H NMR (300 MHz, CDCl_3) δ_{H} : 7.37-7.28 (m, 2H), 7.24-7.20 (m, 1H), 3.80 (s, 2H),
20 3.52-3.48 (m, 2H), 3.32 (m, 3H), 3.12 (m, 1H), 2.08-2.0 (m, 1H), 1.75 (m, 1H), 1.45 (s, 9H).

The following secondary amines were similarly prepared by reductive alkylation of 1,1-dimethylethyl (3S)-3-aminopiperidine-1-carboxylate with the appropriate
25 benzaldehyde:

1,1-Dimethylethyl (3S)-3-{{[3,5-dichloro-phenyl]methyl}-amino}pyrrolidine-1-carboxylate.

1,1-Dimethylethyl (3S)-3-{{[5-fluoro-2-(trifluoromethyl)-phenyl]methyl}amino}pyrrolidine-1-carboxylate.

1,1-Dimethylethyl (3*S*)-3-[[2-chloro-4-fluoro-phenyl]-methyl]amino}pyrrolidine-1-carboxylate.

Example 1E: (3*S*)-*N*-(1-Methylethyl)-*N*-{[3,5-dichlorophenyl]-methyl}pyrrolidin-3-amine *D*-tartrate

a) 1,1-Dimethylethyl (3*S*)-3-((1-methylethyl)-{[3,5-dichlorophenyl]methyl}amino)-pyrrolidine-1-carboxylate

To a solution of 1,1-dimethylethyl (3*S*)-3-[(1-methylethyl)amino]-pyrrolidine-1-carboxylate (1g, 4.4 mmol) and 3,5-dichlorobenzaldehyde (1.53g, 8.77 mmol) in trimethylorthoformate (10 mL) at room temperature under a nitrogen atmosphere was added portionwise sodium triacetoxyborohydride (1.3g, 6.1 mmol). The reaction was stirred at room temperature for 72 hours, then evaporated to dryness *in vacuo*. The residue was taken up in aqueous saturated sodium hydrogen carbonate/dichloromethane mixture. The aqueous layer was further extracted with dichloromethane (3X), and the combined organic layers dried (MgSO₄) and evaporated to dryness *in vacuo*. The resulting residue was dissolved in methanol and filtered through a cationic ion exchange resin (Isolute™ SCX-2). The basic components were recovered from the column by elution with 2N ammonia in methanol. This solution was concentrated *in vacuo* to yield the desired compound as a yellow oil that was used in the next step without further purification. ¹H NMR (300 MHz, CDCl₃) δ_H: 0.95-1.04 (m, 6H), 1.45 (s, 9H), 1.56-1.77 (m, 1H), 1.8-1.94 (m, 1H), 2.9-3.09 (m, 2H), 3.11-3.25 (m, 1H), 3.32-3.56 (m, 3H), 3.59 (s, 2H), 7.15-7.27 (m, 3H). MS: [M+H] = 387/389/391.

b)(3*S*)-*N*-(1-Methylethyl)-*N*-{[3,5-dichlorophenyl]methyl}-pyrrolidin-3-amine *D*-tartrate
1,1-Dimethylethyl (3*S*)-3-((1-methylethyl)-{[3,5-dichlorophenyl]methyl}amino)pyrrolidine-1-carboxylate (1.36g, 3.51 mmol) was dissolved in a mixture of dichloromethane and trifluoroacetic acid (10 mL, 2:1) and stirred at room temperature for 30 minutes. The reaction solution was concentrated *in vacuo* and redissolved in MeOH. This solution was filtered through a cationic ion exchange resin (Isolute™ SCX-2). The basic components were isolated by elution with

2N ammonia in methanol and further purified by UV guided prep-LC. The desired compound was isolated from the acidic prep-LC mobile phase *via* a cationic ion exchange resin as described above. After evaporation *in vacuo* the residue was dissolved in hot cyclohexane (5 mL) and to this was added an equimolar amount of D-tartaric acid (450 mg), dissolved in a minimal amount of hot isopropanol. The solution was evaporated *in vacuo* to yield the title compound as a solid. ¹H NMR (300 MHz, d6-DMSO) δ_H: 0.95-0.99 (m, 6H), 1.58-1.71 (m, 1H), 1.91-2.00 (m, 1H), 2.76-2.91 (m, 2H), 2.97-3.07 (m, 1H), 3.18-3.25 (m, 2H), 3.55-3.67 (m, 4H), 3.95 (s, 2H), 7.37-7.38 (m, 2H), 7.43-7.45 (m, 1H). MS: [M+H] = 287/289/291.

10

The following Examples were similarly prepared as described above for Example 1E, by reductive alkylation of 1,1-dimethylethyl (3*S*)-3-[(1-methylethyl)amino]-pyrrolidine-1-carboxylate with the appropriate substituted benzaldehyde:

15 **Example 2E: (3*S*)-*N*-(1-Methylethyl)-*N*-{[2-(methylthio)phenyl]methyl}-pyrrolidin-3-amine fumarate**

¹H NMR (300 MHz, CD₃OD) δ_H: 0.99 (s, 6H), 2.06 (m, 1H), 2.37 (s, 3H), 3.01-2.85 (m, 1H), 3.18-3.06 (m, 1H), 3.46-3.19 (m, 4H), 3.67 (dd, 2H), 6.60 (s, 2H), 7.10-7.02 (m, 20 1H), 7.20-7.11 (m, 2H), 7.40 (dd, 1H); MS: [M+H] = 265.

The following Examples were similarly prepared as described above for Example 1E, by reductive alkylation of 1,1-dimethylethyl (3*S*)-3-[(cyclohexylmethyl)amino]-pyrrolidine-1-carboxylate with the appropriate substituted benzaldehyde:

25

Example 3E: (3*S*)-*N*-(Cyclohexylmethyl)-*N*-{[2-(methylthio)phenyl]methyl}pyrrolidin-3-amine fumarate

¹H NMR (300 MHz, CD₃OD) δ_H: 0.86-0.69 (s, 3H), 1.22-1.12 (m, 3H), 1.41-1.29 (m, 30 1H), 1.84-1.67 (m, 5H), 2.16-1.95 (m, 2H), 2.34 (d, 2H), 2.38 (s, 3H), 3.23-3.05 (m, 1H),

3.44-3.28 (m, 4H), 3.78-3.55 (m, 2H), 6.70 (s, 2H), 7.16 (s, 2H), 7.35-7.32 (m, 1H); MS: [M+H] = 319.

5 **Example 4E: (3S)-N-(Cyclohexylmethyl)-N-[(2-fluorophenyl)methyl]-pyrrolidin-3-amine fumarate**

¹H NMR (300 MHz, CD₃OD) δ_H: 0.83-0.75 (s, 6H), 1.24-1.17 (m, 3H), 1.48-1.42 (m, 1H), 1.85-1.68 (m, 5H), 2.03-1.92 (m, 1H), 2.17-2.10 (m, 1H), 2.35 (d, 2H), 3.25-3.05 (m, 1H), 3.44-3.32 (m, 4H), 3.81-3.62 (m, 2H), 6.71 (s, 2H), 7.20-7.05 (m, 2H), 7.33-7.27 (m, 1H), 7.47-7.42 (m, 1H); MS: [M+H] = 291.

Example 5E: (3S)-N-[(2-Chlorophenyl)methyl]-N-(cyclohexylmethyl)-pyrrolidin-3-amine fumarate

15 ¹H NMR (300 MHz, CD₃OD) δ_H: 0.89-0.77 (m, 2H), 1.24-1.13 (m, 3H), 1.36 (d, 6H), 1.49-1.42 (m, 1H), 1.83-1.68 (m, 5H), 2.15-1.93 (m, 2H), 2.35 (d, 2H), 3.20-3.06 (m, 1H), 3.33-3.23 (m, 4H), 3.75-3.42 (m, 2H), 4.69-4.61 (m, 1H), 6.70 (s, 2H), 6.98-6.88 (m, 2H), 7.35 (d, 1H), 7.50-7.19 (m, 1H); MS: [M+H] = 307.

20 **Example 6E: (3S)-N-(Cyclohexylmethyl)-N-([2-[l-(methylethyl)oxy]-phenyl]methyl)pyrrolidin-3-amine fumarate**

¹H NMR (300 MHz, CD₃OD) δ_H: 0.89-0.77 (m, 2H), 1.24-1.13 (m, 3H), 1.36-1.34 (dd, 6H), 1.49-1.42 (m, 1H), 1.83-1.68 (m, 5H), 1.93 (m, 2H, m), 2.35 (d, 2H), 3.20-3.06 (m, 1H), 3.33-3.23 (m, 4H), 3.75-3.42 (m, 2H), 4.69-4.61 (m, 1H), 6.70 (s, 2H), 6.98-6.88 (m, 2H), 7.35 (d, 1H), 7.50-7.19 (m, 1H); MS: [M+H] = 331.

Example 7E: (3S)-N-([5-Fluoro-2-(trifluoromethyl)phenyl]methyl)-N-(tetrahydro-2H-pyran-4-yl)pyrrolidin-3-amine D-tartrate

a) 1,1-Dimethylethyl (3*S*)-3-[(tetrahydro-2*H*-pyran-4-yl)amino]pyrrolidine-1-carboxylate

Neat tetrahydro-4*H*-pyran-4-one (18.7g, 100mmol) and 1,1-dimethylethyl (3*S*)-3-aminopyrrolidine-1-carboxylate (26.1g, 140.1 mmol) were stirred together for 20 minutes prior to addition of anhydrous dichloroethane (140mL). The solution was then cooled to 0°C under nitrogen and stirred as sodium triacetoxyborohydride (59.2g, 281mmol) was added portionwise. The reaction was allowed to warm to room temperature and stirred for 5 days, after which the reaction solution was carefully poured onto ice-cold aqueous sodium hydrogen carbonate solution. The phases were separated and the aqueous phase washed with dichloromethane. The combined organic phases were dried (MgSO₄) and concentrated *in vacuo*. The crude product was purified by automated flash chromatography on silica, eluting with methanol in ethyl acetate (0:100 to 30:70), to provide the title compound as an off-white solid. ¹H NMR (300 MHz, d₆-DMSO) δ_H: 1.13-1.29 (m, 2H), 1.39 (s, 9H), 1.55-1.65 (m, 1H), 1.68-1.81 (m, 2H), 1.87-2.00 (m, 1H), 2.64 (sep, 1H), 2.91 (sex, 1H), 3.10-3.45 (m, 6H), 3.81 (dt, 2H). MS: [M+H] = 271, [M+H-tBu] = 215.

b) (3*S*)-*N*-{[5-Fluoro-2-(trifluoromethyl)phenyl]methyl}-*N*-(tetrahydro-2*H*-pyran-4-yl)pyrrolidin-3-amine *D*-tartrate

To a stirred solution of 1,1-dimethylethyl (3*S*)-3-[(tetrahydro-2*H*-pyran-4-yl)amino]pyrrolidine-1-carboxylate (1.12g, 4.2mmol) and 5-fluoro-2-(trifluoromethyl)benzaldehyde (4.56g, 23.8mmol) in anhydrous dichloroethane (50mL) was added portionwise sodium triacetoxyborohydride (3.86g, 18.3mmol). The reaction mixture was stirred at room temperature under nitrogen and the reaction progress was followed by MS. After 2 days more reagents were added: 5-fluoro-2-(trifluoromethyl)benzaldehyde (0.98g, 5.1mmol) and sodium triacetoxyborohydride (3.00g, 14.2mmol), and after a further 2 days the reaction was found to be complete. The reaction solution was carefully poured onto ice-cold saturated aqueous sodium hydrogen carbonate solution and filtered through a PTFE hydrophobic frit. The organic phase was concentrated *in vacuo* and the residue redissolved in methanol. The methanolic solution was filtered through a cationic ion exchange resin (Isolute™ SCX-2) and the basic components isolated by elution with 2N ammonia in methanol. After concentrating *in*

vacuo, the residue was redissolved in dichloromethane /trifluoro-acetic acid (2:1) and allowed to stir at room temperature for 4 hours. The reaction mixture was concentrated *in vacuo* and redissolved in methanol. The methanolic solution was filtered through a cationic ion exchange resin (Isolute TM SCX-2) and the basic components isolated by
5 elution with 2N ammonia in methanol. The crude product was purified by UV guided prep-LC, and the desired compound collected from the acidic prep-LC mobile phase *via* a cationic ion exchange resin, as described above. The basic product was dissolved in hot cyclohexane and to this was added an equimolar amount of *D*-tartaric acid dissolved in a minimal amount of hot isopropanol. The solution was allowed to cool overnight, and the
10 next day the resultant solid was filtered off and dried *in vacuo*, to yield the title compound as a white crystalline solid. ¹H NMR (300 MHz, d6-DMSO) δ_H : 1.40-1.80 (m, 5H), 1.91-2.06 (m, 1H), 2.61-2.74 (m, 1H), 2.81-2.93 (dd, 1H), 2.97-3.11 (dt, 1H), 3.12-3.31 (m, 4H), 3.69-3.96 (m, 7H), 7.49-7.61 (m, 2H), 7.90-7.99 (m, 1H). MS: [M+H] = 347.

15 The following Examples were similarly prepared from 1,1-dimethylethyl (3*S*)-3-[(tetrahydro-2*H*-pyran-4-yl)amino]pyrrolidine-1-carboxylate and the appropriate benzaldehyde, as described above for Example 7E:

20 **Example 8E: (3*S*)-*N*-{[2-(Trifluoromethyl)phenyl]methyl}-*N*-(tetrahydro-2*H*-pyran-4-yl)pyrrolidin-3-amine hemi-*D*-tartrate**

¹H NMR (300 MHz, d6-DMSO) δ_H : 1.35-1.75 (m, 5H), 1.90-2.04 (m, 1H), 2.63-2.75 (m, 1H), 2.76-2.86 (m, 1H), 2.94-3.03 (m, 1H), 3.10-3.25 (m, 4H), 3.67-3.90 (m, 6H), 7.43 (t, 1H), 7.66 (t, 2H), 7.92 (d, 1H); MS: [M+H] = 329.

25

Example 9E: (3*S*)-*N*-(1-Methylethyl)-*N*-{[2-(trifluoromethyl)-5-fluorophenyl]methyl}pyrrolidin-3-amine fumarate

a) 1,1-Dimethylethyl (3*S*)-3-((1-methylethyl)-{[2-(trifluoromethyl)-5-fluorophenyl]methyl}amino)-pyrrolidine-1-carboxylate
30

A solution of 1,1-dimethylethyl (3*S*)-3-[(1-methylethyl)amino]pyrrolidine-1-carboxylate (0.34g, 1.5mmol) and 2-(trifluoromethyl)-5-fluorobenzyl bromide (0.58g, 2.25mmol) in acetonitrile (5mL) was heated at reflux with anhydrous potassium carbonate (0.41g, 3mmol) for 24 hours. The reaction mixture was cooled, diluted with ethyl acetate and washed with water. The organic extracts were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo* to give an oil. This was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (0:100 to 10:90), to give the title compound as an oil.

b) (3*S*)-*N*-(1-Methylethyl)-*N*-{[2-(trifluoromethyl)-5-fluorophenyl]methyl}pyrrolidin-3-amine fumarate

A solution of 1,1-dimethylethyl (3*S*)-3-((1-methylethyl)-{[2-(trifluoromethyl)-5-fluorophenyl]-methyl}amino)-pyrrolidine-1-carboxylate (0.26g) in a mixture of trifluoroacetic acid (2mL), dichloromethane (8mL) and water (0.2mL) was stirred at room temperature for 3 hours. The reaction mixture was evaporated *in vacuo*. The crude mixture was taken up in methanol and absorbed onto an SCX-2 ion exchange cartridge. After initially washing with methanol, the product was eluted with 2M methanolic ammonia and the collected fractions evaporated *in vacuo*. The crude product was taken up in methanol and fumaric acid (1 equiv.) in methanol added. The solvent was removed *in vacuo* and the resultant gum triturated with diethyl ether. The solid formed was filtered off and dried *in vacuo* at 50°C to yield the title compound as an off-white microcrystalline solid. ¹H NMR (300 MHz, CD₃OD) δ_H: 1.09 (d, 3H), 1.10 (d, 3H), 1.87 (m, 1H), 2.15 (m, 1H), 3.01 (m, 2H), 3.23 (m, 1H), 3.38 (m, 2H), 3.81 (m, 1H), 3.91 (s, 2H), 6.70 (s, 2H), 7.15 (dt, 1H), 7.73 (m, 2H); MS: [M+H] = 305.

The following Examples were similarly prepared as described for Example 9E, using the appropriate substituted benzyl bromide in step b) above:

Example 10E: (3S)-N-([1,1'-Biphenyl]-2-ylmethyl)-N-(1-methylethyl)-pyrrolidin-3-amine fumarate

¹H NMR (300 MHz, CD₃OD) δ_H: 0.95 (d, 6H), 1.75 (m, 1H), 1.91 (m, 1H), 2.75 (dd, 1H), 2.93 (sept, 1H), 3.10 (m, 2H), 3.25 (m, 1H), 3.60 (m, 3H), 6.70 (s, 2H), 7.17 (dd, 1H), 7.25-7.48 (m, 7H), 7.67 (d, 1H); MS: [M+H] = 295.

Example 11E: Methyl ((3S)-pyrrolidin-3-yl){2-(trifluoromethyl)phenyl}-methyl}amino)acetate D-tartrate

10

60% Sodium hydride oil dispersion (39mg, 0.95mmol) was added to 1,1-dimethylethyl (3S)-3-({2-(trifluoromethyl)-phenyl}methyl}amino)pyrrolidine-1-carboxylate (250mg, 0.73mmol) in DMF (5mL). After heating at 50°C for 1 hour under nitrogen, methyl bromoacetate (123mg, 0.73mmol) was added. After heating overnight at 50°C overnight, excess water was added and the product was extracted into ether. The ether was washed with water, dried (MgSO₄) and evaporated *in vacuo* to give an oil (460mg). The oil was dissolved in dichloromethane (5mL) and trifluoroacetic acid (0.5mL) was added. After stirring for 1 day, the solution was evaporated *in vacuo* to give an oil. The oil was purified using preparative LCMS to give the product as the acetate salt, which was converted to the free base by absorption onto a cationic ion exchange resin (Isolute™ SCX-2) and eluting the basic fractions with 2N ammonia in methanol. The resultant oil was converted to the D-tartaric acid salt (crystallised from ethanol/diethyl ether) to give the title compound as a white solid. ¹H NMR(300 MHz, CD₃OD) δ_H: 1.84-1.96 (m, 1H), 2.06-2.14 (m, 1H), 3.06-3.37 (2 x m, 6H), 3.57 (s, 3H), 3.77-3.86 (quin, 1H), 3.91-4.06 (q, 2H), 4.29 (s, 2H), 7.32-7.36 (t, 1H), 7.49-7.54 (t, 1H), 7.56-7.59 (d, 1H), 7.76-7.89 (d, 1H); MS: [M+H] = 317.

The following Examples were prepared from 1,1-dimethylethyl (3S)-3-aminopyrrolidine-1-carboxylate by initial reductive alkylation with 2-methylpropanaldehyde, followed by a second reductive alkylation with the appropriate benzaldehyde and subsequent deprotection.

30

Example 12E: (3S)-N-{[2-(Methoxy)phenyl]methyl}-N-(2-methylpropyl)pyrrolidin-3-amine fumarate

5 ¹H NMR (300 MHz, CD₃OD) δ_H: 0.82 (dd, 6H), 1.66 (sept, 1H), 1.79-1.92 (m, 1H), 1.92-2.06 (m, 1H), 2.19-2.22 (m, 2H), 2.96-3.13 (m, 2H), 3.18-3.31 (m, 2H), 3.59-3.67 (m, 2H), 3.74 (s, 3H), 6.59 (s, 2H), 6.80-6.87 (m, 2H), 7.11-7.18 (m, 1H), 7.25 (dd, 1H); MS: [M+H] = 263.

10 The following Examples were prepared from 1,1-dimethylethyl (3S)-3-({[2-(trifluoromethyl)phenyl]-methyl}amino)pyrrolidine-1-carboxylate by reductive alkylation with the appropriate aldehyde or ketone and subsequent deprotection.

Example 13E: (3S)-N-(1-Methylethyl)-N-{[2-(trifluoromethyl)-phenyl]methyl}pyrrolidin-3-amine fumarate

15 ¹H NMR (300 MHz, CD₃OD) δ_H: 7.98-8.00 (d, 1H), 7.60-7.68 (d+t, 2H), 7.38-7.43 (t, 1H), 6.70 (s, 2H), 3.91 (bs, 2H), 3.74-3.85 (m, 1H), 3.17-3.40 (M, 5H), 2.96-3.10 (m, 3H), 2.08-2.18 (m, 1H), 1.82-1.96 (m, 1H), 1.08-1.11 (dd, 6H); MS: [M+H] = 287.

Example 14E: (3S)-N-Ethyl-N-{[2-(trifluoromethyl)phenyl]methyl}-pyrrolidin-3-amine fumarate

20 ¹H NMR (300 MHz, CD₃OD) δ_H: 8.00-8.03 (d, 1H), 7.67-7.76 (d+t, 2H), 7.47-7.52 (t, 1H), 6.77 (s, 2H), 3.89-4.03 (q, 2H), 3.65-3.75 (quin, 2H), 3.43-3.53 (m, 2H), 3.28-3.41 (m, 1H), 3.17-3.23 (m, 1H), 2.73-2.84 (q, 2H), 2.19-2.30 (m, 2H), 2.19-2.30 (m, 1H),
25 1.98-2.14 (m, 1H), 1.10-1.15 (t, 3H); MS: [M+H] = 273.

Example 15E: (3S)-N-Propyl-N-{[2-(trifluoromethyl)phenyl]methyl}-pyrrolidin-3-amine fumarate

¹H NMR (300 MHz, CD₃OD) δ_H: 7.92-7.94 (d, 1H), 7.60-7.69 (d+t, 2H), 7.40-7.45 (t, 1H), 6.69-6.73 (s, 2H), 3.82-3.98 (q, 2H), 5.59-3.69 (quin, 1H), 3.35-3.45 (m, 2H), 2.80-3.21 (m, 1H), 3.08-3.15 (m, 1H), 2.54-2.59 (q, 2H), 2.10-2.21 (m, 1H), 1.90-2.06 (m, 1H), 1.44-1.56 (quin, 2H), 0.86-0.91 (T, 3H); MS: [M+H] = 287.

Example 16E: (3S)-N-(Cyclohexylmethyl)-N-{[2-(trifluoromethyl)phenyl]methyl}pyrrolidin-3-amine fumarate

¹H NMR (300 MHz, CD₃OD) δ_H: 7.78-7.92 (d, 1H), 7.61-7.70 (d+t, 2H), 7.41-7.49 (t, 1H), 6.70 (s, 2H), 3.81-3.95 (q, 2H), 3.56-3.67 (quin, 1H), 3.31-3.43 (m, 2H), 3.14-3.23 (m, 1H), 3.04-3.11 (m, 1H), 2.39-2.41 (d, 2H), 2.06-2.13 (m, 1H), 1.70-2.01 (m, 6H), 1.34-1.46 (m, 1H), 1.12-1.23 (m, 1H), 0.83-0.89 (m, 2H); MS: [M+H] = 341.

Example 17E: (3S)-N-Butyl-N-{[2-(trifluoromethyl)phenyl]methyl}-pyrrolidin-3-amine fumarate

¹H NMR (300 MHz, CD₃OD) δ_H: 7.91-7.94 (d, 1H), 7.60-7.69 (m, 2H), 7.40-7.45 (t, 1H), 6.70 (s, 2H), 3.82-3.96 (q, 2H), 3.59-3.69 (quin, 1H), 3.32-3.50 (m, 2H), 3.22-3.29 (m, 1H), 3.09-3.15 (q, 1H), 2.58-2.63 (t, 2H), 2.10-2.21 (m, 1H), 1.90-2.04 (m, 1H), 1.42-1.51 (m, 2H), 1.17-1.37 (m, 2H), 0.87-0.91 (t, 3H); MS: [M+H] = 301.

Example 18E: (3S)-N-(2-Ethylbutyl)-N-{[2-(trifluoromethyl)phenyl]methyl}pyrrolidin-3-amine sesquifumarate

¹H NMR (300 MHz, CD₃OD) δ_H: 7.77-7.80 (d, 1H), 7.49-7.60 (m, 2H), 7.29-7.34 (t, 1H), 6.60 (s, 1.5H), 3.70-3.81 (q, 2H), 3.46-3.57 (quin, 1H), 3.20-3.33 (m, 2H), 2.94-3.13 (m, 2H), 2.32-2.34 (d, 2H), 1.97-2.07 (m, 1H), 1.78-1.91 (m, 1H), 1.05-1.40 (m, 5H), 0.69-0.76 (m, 6H). MS: [M+H] = 329.

Example 19E: (3S)-N-{{2-(Trifluoromethyl)phenyl}methyl}-N-(3,3,3-trifluoropropyl)pyrrolidin-3-amine fumarate

¹H NMR (300 MHz, CD₃OD) δ_H: 7.76-7.78 (d, 1H), 7.50-7.60 (d+t, 2H), 7.32-7.37 (t, 1H), 6.58 (s, 2H), 3.75-3.89 (q, 2H), 3.48-3.59 (quin, 1H), 3.126-3.22 (m, 1H), 2.98-3.05 (dd, 1H), 2.75-2.80 (t, 2H), 2.18-2.34 (m, 2H), 2.02-2.13 (m, 1H), 1.80-1.93 (m, 1H); MS: [M+H] = 341.

Example 20E: (3S)-N-(Furan-2-ylmethyl)-N-{{2-(trifluoromethyl)phenyl}methyl}pyrrolidin-3-amine D-tartrate

10

¹H NMR (300 MHz, CD₃OD) δ_H: 7.83-7.86 (d, 1H), 7.49-7.58 (t+s, 2H), 7.29-7.38 (m, 2H), 6.23-6.26 (m, 1H), 6.14-6.15 (m, 1H), 4.30 (s, 2H), 3.78-3.91 (q, 2H), 3.66-3.67 (m, 2H), 3.25-3.55 (m, 3H), 2.30-3.17 (m, 2H), 2.05-2.16 (m, 1H), 1.83-1.96 (m, 1H); MS: [M+H] = 325.

15

Example 21E: (3S)-N-[3-(Methylthio)propyl]-N-{{2-(trifluoromethyl)phenyl}methyl}pyrrolidin-3-amine D-tartrate

¹H NMR (300 MHz, CD₃OD) δ_H: 7.90-7.92 (d, 1H), 7.61-7.70 (d+t, 2H), 7.41-7.46 (t, 1H), 4.42 (s, 2H), 3.84-3.97 (q, 2H), 3.59-3.69 (quin, 1H), 3.38-3.47 (m, 2H), 3.19-3.29 (m, 1H), 3.09-3.16 (m, 1H), 2.70-2.77 (dt, 2H), 2.48-2.52 (t, 2H), 2.08-2.21 (m, 1H), 1.89-2.08 (s+m, 4H), 1.69-1.79 (quin, 2H); MS: [M+H] = 333.

20

Example 22E: N-(Phenylmethyl)-N-[(3S)-pyrrolidin-3-yl]-N-{{2-(trifluoromethyl)phenyl}methyl}amine fumarate

25

¹H NMR (300 MHz, CD₃OD) δ_H: 7.93-7.96 (d, 1H), 7.60-7.68 (q, 2H), 7.23-7.44 (m, 6H), 6.69 (s, 2H), 3.83-3.94 (s, 2H), 3.61-3.80 (m, 3H), 3.32-3.44 (m, 2H), 3.08-3.25 (m, 2H), 1.99-2.22 (m, 2H); MS: [M+H] = 335.

Example 23E: (3S)-N-{{2-(Methyloxy)phenyl}methyl}-N-{{2-(trifluoromethyl)phenyl}methyl}pyrrolidin-3-amine fumarate

¹H NMR (300 MHz, CD₃OD) δ_H: 7.85-7.87 (d, 1H), 7.61-7.64 (d, 1H), 7.52-7.58 (t, 1H), 7.21-7.40 (m, 3H), 6.81-6.97 (m, 2H), 6.69 (s, 2H), 3.61-3.97 (m, 8H), 3.16-3.44 (m, 4H), 1.20-2.21 (m, 2H); MS: [M+H] = 365.

Example 24E: (3S)-N,N-bis{{2-(Trifluoromethyl)phenyl}methyl}-pyrrolidin-3-amine fumarate

¹H NMR (300 MHz, CD₃OD) δ_H: 7.90-7.92 (d, 2H), 7.66-7.69 (d, 2H), 7.59-7.64 (t, 2H), 7.40-7.45 (t, 2H), 6.69 (s, 2H), 3.91 (s, 4H), 3.62-3.74 (quin, 1H), 3.36-3.46 (m, 2H), 3.16-3.26 (m, 2H), 2.02-2.24 (m, 2H); MS: [M+H] = 403.

The following examples illustrate compounds of of Formulae (IF) above and methods for their preparation.

Preparation of Intermediates

1,1-Dimethylethyl (3S)-3-aminopiperidine-1-carboxylate

e) 1,1-Dimethylethyl (3R)-3-hydroxypiperidine-1-carboxylate

Solid di-*tert*-butyldicarbonate (26.6g, 122mmol) was added in portions over 15 minutes to a stirred solution of (3R)-piperidin-3-ol hydrochloride (15.25g, 111mmol), triethylamine (30.9mL, 222mmol) and 4-(dimethylamino)-pyridine (50mg) in dry dichloromethane (300mL). After stirring for 18 hours at room temperature, the mixture was washed with aqueous citric acid, then brine. The organic extracts were dried (MgSO₄), filtered and evaporated *in vacuo* to give an oil. This was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (20:80 to 80:20), to give the title compound as a solid.

f) 1,1-Dimethylethyl (3R)-3-[(methylsulfonyl)oxy]-piperidine-1-carboxylate

Methanesulfonyl chloride (9.56mL, 124mmol) was added dropwise over 10 minutes to a stirred solution of 1,1-dimethylethyl (3*R*)-3-hydroxypiperidine-1-carboxylate (20.7g, 103mmol) and triethylamine (21.5mL, 154mmol) in dichloromethane (300mL) at 0°C. After stirring for 3 hour at 0°C, the reaction was quenched by addition of water. The organic phase was washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo* to give an oil. This was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (20:80 to 50:50), to give the title compound as an oil.

g) 1,1-Dimethylethyl (3*S*)-3-azidopiperidine-1-carboxylate

Sodium azide (7.65g, 118mmol) was added to a solution of 1,1-dimethylethyl (3*R*)-3-[(methylsulfonyl)oxy]-piperidine-1-carboxylate (21.9g, 78.5mmol) in dry dimethylformamide (120mL) and the resultant suspension heated at 70°C for 48 hours. After cooling to room temperature, the reaction mixture was diluted with water and extracted into ethyl acetate. The organic phase was washed two further times with water, then brine. The organic extracts were dried (MgSO₄), filtered and evaporated *in vacuo* to give an oil. This was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (10:90 to 50:50), to give the title compound as an oil.

h) 1,1-Dimethylethyl (3*S*)-3-aminopiperidine-1-carboxylate

A mixture of 1,1-dimethylethyl (3*S*)-3-azidopiperidine-1-carboxylate (7.5g) and 10% palladium-on-carbon (0.75g) in methanol (100mL) was hydrogenated in a Parr apparatus at 70 p.s.i. for 16 hours. The catalyst was removed by filtration through Celite and the solvent evaporated *in vacuo* to give an oil. The resultant title compound was used in subsequent reactions without further purification.

2-(Bromomethyl)-4-fluoro-1,1'-biphenyl

a) Methyl 5-fluoro-2-{[(trifluoromethyl)sulfonyl]-oxy} benzoate

5-Fluorosalicylic acid methyl ester (28.2g, 166mmol) was dissolved in dry dimethylformamide (165mL) and stirred as sodium hydride (60% in oil) (7.30g, 1.1eq) was added portionwise over 30 mins at 0°C. The reaction mixture was stirred for a further

30 mins at room temperature, then *N*-phenyl trifluoromethanesulfonimide (62.8g, 1.05eq) was added in portions over 30 mins, then left to stir for 3 hours. The mixture was diluted with diethyl ether and washed successively with water, then brine. The organic layers were combined, dried (MgSO₄), filtered and the solvent removed *in vacuo*. The resulting
5 oil was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (10:90 to 40:60), to give the title compound as an oil.

b) Methyl 4-fluoro-[1,1'-biphenyl]-2-carboxylate

Palladium acetate (635mg, 0.05eq), tricyclohexyl-phosphine (952mg, 0.06eq),
10 potassium fluoride (10.85g, 3.3eq) and phenyl boronic acid (7.6g, 1.1eq) were taken up in dry THF (150mL) and the reaction mixture flushed with nitrogen for 5 mins. A solution of methyl 5-fluoro-2-[[trifluoromethyl)sulfonyl]oxy]benzoate (17.12g, 56.7 mmol) in THF (20mL) was added in one portion and the reaction mixture stirred at reflux under nitrogen for 5 hours. The reaction mixture was cooled to room temperature, diluted with
15 ethyl acetate, then washed with water, dried (MgSO₄), filtered and the solvent removed *in vacuo*. The resulting oil was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (3:97 to 10:90), to give the title compound as an oil.

c) (4-Fluoro-[1,1'-biphenyl]-2-yl)methanol

20 A solution of methyl 4-fluoro-[1,1'-biphenyl]-2-carboxylate (3g, 13.1mmol) in THF (20mL) was added at 0°C to a suspension of lithium aluminium hydride pellets (1g, 26mmol) in THF (30mL). Upon addition the reaction mixture was heated at 60°C under nitrogen for 2 h. The reaction was then cooled to 0°C and the excess lithium aluminium hydride destroyed by adding water, then 1N sodium hydroxide (2mL). The mixture was
25 extracted into diethyl ether and the organic phase was dried (MgSO₄), filtered and the solvent removed *in vacuo*. The resulting oil was purified by flash chromatography on silica, eluting with ethyl acetate/heptane (2:98 to 25:75), to give the title compound as an oil.

30 d) 2-(Bromomethyl)-4-fluoro-1,1'-biphenyl

Triphenylphosphine dibromide (35.5g, 2eq) was added in one portion to a solution of (4-fluoro-[1,1'-biphenyl]-2-yl)methanol (8.5g, 42mmol) in chloroform (250mL). The reaction mixture was heated at 60°C and left to stir overnight. The solid was filtered off and the solvent removed *in vacuo*. The resulting oil was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (0:100 to 30:70), to give the title compound as an oil.

Example 1F: (3*S*)-*N*-(2-Methylpropyl)-*N*-{[2-(trifluoromethyl)-phenyl]methyl}piperidin-3-amine, fumarate

10

a) 1,1-Dimethylethyl (3*S*)-3-([2-(trifluoromethyl)-phenyl]methyl)amino)piperidine-1-carboxylate

1,1-Dimethylethyl (3*S*)-3-aminopiperidine-1-carboxylate (1.0g, 5mmol), 2-trifluoromethylbenzaldehyde (0.87g, 5mmol), 5% palladium on carbon (0.35g) and ethanol (40mL) were hydrogenated at 60psi for 2.5 h. using a Parr hydrogenator. The catalyst was filtered off and the filtrate evaporated *in vacuo*. The resultant oil was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (0:100 to 75:25), to give the title compound as an oil.

20 b) 1,1-Dimethylethyl (3*S*)-3-((2-methylpropyl){[2-(trifluoromethyl)phenyl]methyl}amino)piperidine-1-carboxylate

Sodium triacetoxyborohydride (0.23g, 1.08mmol) was added to a stirred solution of 1,1-dimethylethyl (3*S*)-3-([2-(trifluoromethyl)phenyl]methyl)amino)piperidine-1-carboxylate (0.19g, 0.53mmol), isobutyraldehyde (0.12g, 1.6mmol) and 1,2-dichloroethane (5mL). After stirring under nitrogen at room temperature for 1 day, the reaction mixture was diluted with methanol (6mL) and absorbed onto a cationic ion exchange resin (Isolute™ SCX-2). After washing the cartridge with methanol (25mL), the basic components were isolated by elution with 2N ammonia in methanol and the eluate evaporated to give an oil.

30

c) (3*S*)-*N*-(2-Methylpropyl)-*N*-{[2-(trifluoromethyl)-phenyl]methyl}piperidin-3-amine, fumarate

1,1-Dimethylethyl (3*S*)-3-((2-methylpropyl){[2-(trifluoromethyl)phenyl]methyl}amino)piperidine-1-carboxylate (0.139mg, 0.335mmol),
5 trifluoroacetic acid (4mL) and dichloromethane (10mL) were stirred at room temperature for 1 day. The solution was evaporated *in vacuo* to give an oil, which was redissolved in methanol and filtered through a cationic ion exchange resin (Isolute™ SCX-2). The basic components were isolated by elution with 2N ammonia in methanol. The eluate was evaporated *in vacuo* and the resultant oil converted to the fumaric acid salt (crystallisation
10 from ethanol/ether), to give the title compound as a white solid. ¹H NMR (300MHz, CD₃OD): δ_H 7.77-7.74 (d, H), 7.51-7.43 (m, 2H), 7.25-7.22 (t, 1H), 4.23 (s, 2H), 3.79-3.66 (q, 2H), 3.21-3.08 (m, 4H), 2.83-2.61 (m, 3H), 2.28-2.10 (m, 2H), 1.90-1.82 (m, 2H), 1.59-1.37 (m, 3H), 0.77-0.72 (t, 6H); MS: (M+H) = 315.

15 The following Examples were similarly prepared as described above for Example 1F, by reductive alkylation of 1,1-dimethylethyl (3*S*)-3-({[2-(trifluoromethyl)-phenyl]methyl}amino)piperidine-1-carboxylate with the appropriate aldehyde or ketone, and subsequent deprotection:

20 **Example 2F: (3*S*)-*N*-(3,3-Dimethylbutyl)-*N*-{[2-(trifluoromethyl)-phenyl]methyl}piperidin-3-amine, *D*-tartrate**

¹HNMR (300MHz, CD₃OD): δ_H 7.79-7.86 (d, 1H), 7.47-7.56 (m, 2H), 7.27-7.32 (t, 2H), 4.30 (s, 2H), 3.73-3.84 (t, 2H), 3.16-3.28 (m, 2H), 2.71-2.89 (m, 3H), 2.47-2.52 (t, 2H), 1.84-1.97 (m, 2H), 1.47-1.63 (m, 2H), 1.22-1.33 (m, 2H), 0.75 (s, 9H); MS: [M+H]
25 = 343.

Example 3F: (3S)-N-Cyclohexyl-N-{[2-(trifluoromethyl)phenyl]-methyl}piperidin-3-amine, D-tartrate

¹HNMR (300MHz, CD₃OD): δ_H 7.88-7.91 (d, 1H), 7.51-7.58 (m, 2H), 7.29-7.34 (t, 1H), 4.29 (s, 2H), 3.68-3.83 (q, 2H), 3.43-3.50 (m, 1H), 3.08-3.27 (m, 1H), 2.87-3.00 (m, 2H), 2.39-2.45 (dd, 1H), 2.22-2.29 (dd, 1H), 2.22-2.16 (m, 2H), 1.76-1.90 (m, 2H), 1.58-1.62 (m, 1H), 1.27-1.41 (m, 2H), 1.08-1.22 (m, 2H), 0.97-1.03 (1H), 0.63-0.74 (m, 4H); MS: [M+H] = 341.

Example 4F: (3S)-N-{[5-Fluoro-2-(trifluoromethyl)phenyl]methyl}-N-tetrahydro-2H-pyran-4-ylpiperidin-3-amine, L-tartrate

a) 1,1-Dimethylethyl (3S)-3-(tetrahydro-2H-pyran-4-ylamino)piperidine-1-carboxylate 1,1-Dimethylethyl-(3S)-3-aminopiperidine-1-carboxylate (2g, 11mmol), 4H-tetrahydropyran-4-one (1.1g, 11mmol) and dichloroethane (40mL) were stirred under nitrogen at room temperature for 15 min. Sodium triacetoxyborohydride (2.9g, 14mmol) was added in 3 lots over 30 minutes and stirred overnight. The reaction was diluted with water (50mL) and made basic by addition of 2N NaOH solution. After stirring for 1h, the mixture was extracted into dichloromethane, and the combined organic extracts washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo* to give the title compound as an oil.

b) (3S)-N-{[5-Fluoro-2-(trifluoromethyl)phenyl]methyl}-N-tetrahydro-2H-pyran-4-ylpiperidin-3-amine, L-tartrate

1,1-Dimethylethyl (3S)-3-(tetrahydro-2H-pyran-4-ylamino)piperidine-1-carboxylate was reductively alkylated with 5-fluoro-2-(trifluoromethyl)benzaldehyde, then deprotected and crystallised as its L-tartrate salt as described above for Example 1 b) and c), to give the title compound. ¹HNMR (300MHz, CD₃OD): δ_H 7.74-7.75 (m, 2H), 7.05-6.98 (t, 1H), 4.50 (s, 2H), 3.99-3.85 (m, 4H), 3.43-2.58 (m, 8H), 2.02-1.42 (m, 8H); MS: [M+H] = 361.

The following Examples were similarly prepared as described above for Example 4F, by reductive alkylation of 1,1-dimethylethyl (3*S*)-3-(tetrahydro-2*H*-pyran-4-ylamino)-piperidine-1-carboxylate with the appropriate benzaldehyde, and subsequent deprotection:

5 **Example 5F: (3*S*)-*N*-[(2-Chloro-5-fluorophenyl)methyl]-*N*-tetrahydro-2*H*-pyran-4-ylpiperidin-3-amine, *L*-tartrate**

¹HNMR (300MHz, CD₃OD): δ_H 7.32-7.24 (m, 2H), 6.92-6.85 (t, 1H), 4.30 (s, 2H), 3.90-3.84 (m, 4H), 3.32-3.17 (m, 4H), 3.08-2.97 (m, 1H), 2.85-2.67 (m, 3H), 1.98-1.82 (m, 2H), 1.73-1.82 (m, 2H), 1.73-1.46 (m, 6H); MS: [M+H] = 327/329.

10

Example 6F: (3*S*)-*N*-([1,1'-Biphenyl]-2-ylmethyl)-*N*-tetrahydro-2*H*-pyran-4-ylpiperidin-3-amine, sesqui-*L*-tartrate

¹HNMR (300MHz, CD₃OD): δ_H 7.51-7.48 (d, 1H), 7.35-7.17 (m, 7H), 7.08-7.05 (d, 1H), 3.30 (s, 1.5H), 3.79-3.74 (dd, 2H), 3.69 (s, 2H), 3.25-3.10 (m, 9H), 3.20-3.09 (m, 15 2H), 2.91-2.77 (m, 2H), 2.66-2.51 (m, 3H); MS: [M+H] = 351.

Example 7F: (3*S*)-*N*-[(2-Chlorophenyl)methyl]-*N*-tetrahydro-2*H*-pyran-4-ylpiperidin-3-amine, *D*-tartrate

¹HNMR (300MHz, CD₃OD): δ_H 7.52-7.49 (d, 1H), 7.26-7.87 (m, 3H), 4.30 (s, 2H), 20 3.92-3.80 (m, 4H), 3.16-2.34 (m, 4H), 2.92-2.05 (m, 1H), 2.90-2.66 (m, 3H), 1.93-1.87 (m, 2H), 1.68-1.39 (m, 6H); MS: [M+H] = 309/311.

Example 8F: (3S)-N-Tetrahydro-2H-pyran-4-yl-N-{{2-(trifluoromethyl)phenyl}methyl}piperidin-3-amine, D-tartrate

¹HNMR (300MHz, CD₃OD): δ_H 7.98-7.95 (d, 1H), 7.71-7.62 (q, 2H), 7.47-7.42 (t, 1H), 4.44 (s, 2H), 4.14-3.98 (m, 4H), 3.43-3.29 (m, 4H), 3.11-2.82 (m, 4H), 2.06-2.03 (m, 2H), 1.82-1.66 (m, 6H); MS: [M+H] = 343.

Example 9F: (3S)-N-Cyclopentyl-N-{{2-(trifluoromethyl)phenyl}-methyl}piperidin-3-amine, L-tartrate

a) 1,1-Dimethylethyl (3S)-3-(cyclopentylamino)-piperidine-1-carboxylate
10 1,1-Dimethylethyl (3S)-3-aminopiperidine-1-carboxylate (2.1 g, 10.5mmol), cyclopentanone (4.65mL, 52.5mmol), and 10% palladium on carbon (0.2g) in methanol (80mL) were hydrogenated at 60psi overnight in a Parr hydrogenator. The catalyst was filtered off and the filtrate evaporated *in vacuo*. The resultant oil was purified by flash chromatography on silica, eluting with ethyl acetate/cyclohexane (15:85 to 30:70), to give
15 the title compound as an oil.

b) 1,1-Dimethylethyl (3S)-3-(cyclopentyl{{2-(trifluoromethyl)phenyl}methyl}amino)piperidine-1-carboxylate
1,1-Dimethylethyl (3S)-3-(cyclopentylamino)-piperidine-1-carboxylate (155mg,
20 0.577mmol), 2-(trifluoromethyl)benzyl bromide (0.105mL, 1.2eq) and anhydrous potassium carbonate (128mg, 1.6eq) in acetonitrile (3mL) were heated at reflux under nitrogen for 2 days. The reaction mixture was cooled to room temperature, diluted with ethyl acetate and washed with water, then brine. The organic extracts were dried (MgSO₄), filtered and evaporated *in vacuo*. The resulting oil was purified by flash
25 chromatography on silica eluting with ethyl acetate/cyclohexane (0:100 to 30:70), to give the title compound as an oil.

c) (3S)-N-Cyclopentyl-N-{{2-(trifluoromethyl)phenyl}-methyl}piperidin-3-amine, L-tartrate

1,1-Dimethylethyl (3*S*)-3-(cyclopentyl {[2-(trifluoromethyl)phenyl]methyl} amino)piperidine-1-carboxylate (160mg, 0.38mmol), trifluoroacetic acid (0.5mL) and dichloromethane (2mL) were stirred at room temperature overnight. The solution was evaporated *in vacuo* to give an oil, which was redissolved in methanol and filtered
5 through a cationic ion exchange resin (Isolute™ SCX-2). The basic components were isolated by elution with 2N ammonia in methanol. The eluate was evaporated *in vacuo* and the resultant oil converted to the *L*-tartaric acid salt (freeze drying from acetonitrile/water 1:1), to give the title compound as a white solid. ¹H NMR (300MHz, CD₃OD): δ_H 7.89-7.86 (d, 1H), 7.54-7.46 (m, 2H), 7.30-7.25 (t, 1H), 4.34 (s, 2H), 3.90-
10 3.78 (q, 2H), 3.30-3.18 (m, 4H), 3.05-2.87 (m, 1H), 2.81-2.59 (m, 2H), 1.95-1.79 (m, 2H), 1.68-1.30 (m, 9H); MS: [M+H] = 327.

The following Examples were similarly prepared as described above for Example 9F, by reaction of 1,1-dimethylethyl (3*R*)-3-(cyclopentylamino)piperidine-1-carboxylate with
15 the appropriate benzyl bromide and subsequent deprotection:

Example 10F: (3*S*)-*N*-([1,1'-Biphenyl]-2-ylmethyl)-*N*-cyclopentyl-piperidin-3-amine,

***L*-tartrate**

¹H NMR (300MHz, CD₃OD): δ_H 7.57-7.55 (d, 1H), 7.35-7.13 (m, 7H), 7.06-7.03
20 (d, 1H), 4.30 (s, 2H), 3.58 (s, 2H), 3.12-2.98 (m, 3H), 2.82-2.73 (m, 1H), 2.65-2.42 (m, 2H), 1.79-1.75(m, 1H), 1.69-1.65 (m, 1H), 1.53-1.19(m, 10H); MS: [M+H] = 335.

Example 11F: (3*S*)-*N*-Cyclopentyl-*N*-([5-fluoro-1,1'-biphenyl]-2-ylmethyl)-piperidin-3-amine, *L*-tartrate

¹H NMR (300MHz, CD₃OD): δ_H 7.35-7.24 (m, 4H), 7.18-7.15 (m, 2H), 7.09-7.04
25 (m, 1H), 6.92-6.85 (m, 1H), 4.28 (s, 2H), 3.55 (m, 2H), 3.22-3.06 (m, 3H), 2.82-2.77 (m, 1H), 2.68-2.58 (m, 2H), 1.88-1.68 (m, 2H), 1.57-1.19 (m, 10H); MS: [M+H] = 353.

Example 12 F: (3*S*)-*N*-(Tetrahydrofuran-3-ylmethyl)-*N*-{[2-(trifluoromethyl)phenyl]methyl}piperidin-3-amine, *L*-tartrate

a) 1,1-Dimethylethyl (3*S*)-3-[(tetrahydrofuran-3-ylmethyl)amino]piperidine-1-carboxylate

- 5 To 5% palladium on carbon (0.05g) under nitrogen was added a solution of 1,1-dimethylethyl-(3*S*)-3-aminopiperidine-1-carboxylate (0.50g, 2.5mmol) and tetrahydrofuran-3-carboxaldehyde (50%^w/_w in water) (0.50g, 2.5mmol) in ethanol (20mL). The reaction mixture was hydrogenated overnight at 60psi in a Parr hydrogenator. The catalyst was removed by filtration through Celite and the solvent
10 removed *in vacuo* to give 1,1-dimethylethyl (3*S*)-3-[(tetrahydrofuran-3-ylmethyl)amino]piperidine-1-carboxylate as a colourless, slightly cloudy oil.

b) (3*S*)-*N*-(Tetrahydrofuran-3-ylmethyl)-*N*-{[2-(trifluoromethyl)phenyl]methyl}piperidin-3-amine, *L*-tartrate

- 15 To a solution of 1,1-dimethylethyl (3*S*)-3-[(tetrahydrofuran-3-ylmethyl)amino]piperidine-1-carboxylate (0.67g, 2.36 mmol) in 1,2-dichloroethane (15 mL) was added 2-(trifluoromethyl)benzaldehyde (0.93mL, 7.07mmol). To this mixture was added a solution of sodium triacetoxymethylborohydride (1.50g, 7.07mmol) in dimethylformamide (3 mL) and left to stir under nitrogen, at room temperature, over the
20 weekend. To the reaction mixture was added water (10 mL) and the solution stirred vigorously for several minutes. The chlorinated organic layer was absorbed directly onto a silica column and the product eluted with methanol/ethyl acetate (0:100 to 30:70). The resultant pale yellow oil was taken up in methanol and absorbed onto a cationic ion
25 exchange resin (IsoluteTM SCX-2). After washing the cartridge with methanol (25mL), the basic components were isolated by elution with 2*N* ammonia in methanol and the eluate evaporated to give 1,1-dimethylethyl (3*S*)-3-{(tetrahydrofuran-3-ylmethyl) {[2-(trifluoromethyl)-phenyl]methyl}amino}piperidine-1-carboxylate as a colourless oil.

- 30 To a solution of this oil (0.82g, 1.85mmol) in dichloromethane (10 mL) was added trifluoroacetic acid (2.06mL, 27.8mmol). The reaction mixture was stirred overnight at room temperature, then the solvent removed *in vacuo*. The resulting oil was taken up in methanol and absorbed onto a cationic ion exchange resin (IsoluteTM SCX-2). After

washing the cartridge with methanol (50mL), the basic components were isolated by elution with 2N ammonia in methanol. The eluate was evaporated *in vacuo* to give a colourless oil. The diastereomers were separated by hplc (Chiralpak AD-H column; 98% heptane, 2% ethanol and 0.2% diethylamine). The faster eluting isomer was taken up in methanol and to this was added a solution of *L*-tartaric acid (0.046g, 0.31 mmol) in methanol. Solvent was removed *in vacuo* and the resulting oil triturated with diethyl ether. Filtration of the resultant suspension gave the title compound as a white solid.

¹HNMR (300MHz, CD₃OD): δ_H 7.75 (1H, d), 7.58-7.50 (2H, m), 7.34-7.29 (1H, m), 4.30 (3H, s), 3.83 (2H, s), 3.70-3.53 (3H, m), 3.42-3.31 (2H, m), 3.16 (1H, m), 2.90-2.67 (3H, m), 2.54-2.34 (2H, m), 2.34-2.20 (1H, m), 1.95-1.84 (3H, m), 1.63-1.45 (3H, m); MS: [M+H] = 343.

The following Examples were prepared from racemic 1,1-dimethylethyl 3-aminopiperidine-1-carboxylate, as described above in Example 1F:

Example 13F: *N*-{[2-(Methyloxy)phenyl]methyl}-*N*-{[2-(trifluoromethyl)phenyl]methyl}piperidin-3-amine

¹HNMR (300MHz, CDCl₃) δ_H 8.04-7.95 (d, 1H), 7.57-7.54 (d, 1H), 7.48-7.44 (m, 2H), 7.28-7.11 (m, 2H), 6.93-6.88 (t, 1H), 6.83-6.80 (d, 1H), 3.94-3.86 (d, 2H), 3.20-3.18 (d, 1H), 2.94-2.90 (d, 1H), 2.68-2.55 (m, 2H), 2.49-2.40 (dt, 1H), 2.08-2.04 (d, 1H), 1.76-1.72 (d, 1H), 1.52-1.25 (m, 4H); MS: [M+H] = 379.

Example 14F: *N*-Cyclohexyl-*N*-{[2-(trifluoromethyl)phenyl]methyl}-piperidin-3-amine

¹HNMR (300MHz, CDCl₃) δ_H 8.01-7.93 (d, 1H), 7.59-7.56 (d, 1H), 7.51-7.46 (t, 1H), 7.30-7.19 (m, 1H), 3.91 (s, 2H), 3.15-3.11 (d, 1H), 3.02-2.98 (d, 1H), 2.88-2.80 (d, 1H), 2.55-2.41 (m, 3H), 1.93-1.01 (m, 14); MS: [M+H] = 341.

Example 15F: *N*-(Phenylmethyl)-*N*-{[2-(trifluoromethyl)phenyl]-methyl}piperidin-3-amine

¹HNMR (300MHz, CDCl₃) δ_H 7.93-7.96 (d, 1H), 7.55-7.61 (d, 1H), 7.47-7.51 (t, 1H), 7.18-7.35 (m, 6H), 3.77-3.90 (q, 2H), 3.64-3.74 (q, 2H), 3.17-3.20 (d, 1H), 2.91-2.95 (d, 1H), 2.53-2.67 (m, 2H), 2.39-2.48 (dt, 1H), 1.97-2.06 (d, 1H), 1.22-1.82 (m, 3H); MS: [M+H] = 349.

Example 16F: (3*S*)-*N*-(2-Methylpropyl)-*N*-{[2-(trifluoromethyl)phenyl]-methyl}-1-azabicyclo[2.2.2]octan-3-amine, sesquifumarate

10

a) (3*S*)-*N*-{[2-(Trifluoromethyl)phenyl]methyl}-1-azabicyclo[2.2.2]octan-3-amine

Sodium triacetoxyborohydride (18.7g, 88.3mmol) was added portionwise over 20 min. to a stirred solution of (3*S*)-1-azabicyclo[2.2.2]octan-3-amine dihydrochloride (5g, 25.1mmol) and 2-trifluoromethylbenzaldehyde (4.81g, 27.6mmol) in DMF (100mL).

15 After stirring under nitrogen for 4 days, the mixture was diluted with excess water, basified with 2N sodium hydroxide and stirred for 1h. The product was extracted into dichloromethane and evaporated *in vacuo* to give an oil, which was dissolved in 2N hydrochloric acid. After washing with ether, the aqueous phase was basified with 2N sodium hydroxide and extracted with dichloromethane. The organic phase was dried
20 (MgSO₄) and evaporated *in vacuo* to give an oil. ¹HNMR (300 MHz, CD₃OD) δ_H: 7.62-7.69 (t, 2H), 7.50-7.55 (t, 1H), 7.32-7.37 (t, 1H), 3.83-3.96 (q, 2H), 3.1-3.19 (m, 1H), 2.72-2.93 (m, 5H), 2.42-2.49 (m, 1H), 1.85-1.95 (m, 1H), 1.63-1.73 (m, 1H), 1.32-1.53); MS: [M+H]= 285.

25 b) (3*S*)-*N*-(2-Methylpropyl)-*N*-{[2-(trifluoromethyl)-phenyl]methyl}-1-azabicyclo[2.2.2]octan-3-amine, sesquifumarate

(3*S*)-*N*-{[2-(Trifluoromethyl)phenyl]methyl}-1-azabicyclo[2.2.2]octan-3-amine (0.30g, 1.06mmol), isobutyraldehyde (0.152g, 2.1mmol) and 1,2-dichloroethane (6mL) were stirred under nitrogen at room temperature for 15 min. Sodium
30 triacetoxyborohydride (0.492g, 2.32mmol) was added in two lots over 5 min. TLC after 1 day showed the reaction to be incomplete, so additional sodium triacetoxyborohydride

(0.24g, 1.15mmol) was added and the mixture heated at 50°C for 5 days. After cooling to room temperature, methanol was added and the mixture was stirred for 1h. This solution was filtered through a cationic ion exchange resin (Isolute™ SCX-2) and the basic fractions isolated by elution with 2N ammonia in methanol to give, after evaporation *in vacuo*, an oil. The crude product was purified using preparative LCMS to give the product as an acetate salt, which was converted to the free base using cationic ion exchange resin as described above. The free base was converted to the fumarate salt, to give the title compound as a white solid from ethanol/diethyl ether. ¹HNMR (300 MHz, CD₃OD) δ_H: 7.88-7.91 (d, 1H), 7.51-7.58 (m, H), 7.30-7.35 (t, 1H), 6.60 (s, 3H), 3.71-3.85 (q, 2H), 3.42-4.50 (m, 1H), 2.88-3.26 (m, 6H), 2.25-2.39 (m, 1H), 2.09-2.23 (m, 3H), 1.74-1.91 (m, 2H), 1.42-1.63 (m, 2H), 0.78-0.83 (t, 6H); MS: [M+H] = 341.

The following Examples were similarly prepared as described above for Example 16F, from (3*S*)-*N*-{[2-(trifluoromethyl)phenyl]methyl}-1-azabicyclo-[2.2.2]octan-3-amine and the appropriate substituted benzaldehyde:

Example 17F: (3*S*)-*N*-([1,1'-Biphenyl]-2-ylmethyl)-*N*-(2-methylpropyl)-1-azabicyclo[2.2.2]octan-3-amine, *D*-tartrate

¹HNMR (300 MHz, CD₃OD) δ_H: 7.50-7.47 (d, 1H), 7.38-7.18 (m, 7H), 7.09-7.06 (dd, 1H), 4.29 (s, 2H), 3.58-3.54 (d, 1H), 3.43-3.39 (d, 1H), 3.25-3.18 (m, 1H), 3.09-3.90 (4H), 2.68-2.63 (t, 1H), 2.45-2.39 (dq, 1H), 2.16-1.98 (m, 3H), 1.83-1.74 (m, 2H), 1.65-1.61 (m, 1H), 1.45-1.42 (m, 1H), 1.31-1.22 (quin, 1H), 0.65-0.61 (t, 6H); MS: [M+H] = 349.

Example 18F: (3*S*)-*N*-{[4-Fluoro-2-(trifluoromethyl)phenyl]methyl}-*N*-(2-methylpropyl)-1-azabicyclo[2.2.2]octan-3-amine, *L*-tartrate

¹HNMR (300 MHz, CD₃OD) δ_H: 7.94-7.89 (t, 1H), 7.34-7.27 (m, 2H), 4.29 (s, 4.29), 3.81-3.66 (q, 2H), 3.51-3.44 (t, 1H), 3.40-2.89 (m, 6H), 2.37-2.04 (m, 4H), 1.93-1.38 (m, 4H), 0.82-0.76 (dd, 6H); MS: [M+H] = 359.


Example 19F: (3S)-N-[(4-Fluoro[1,1'-biphenyl]-2-yl)methyl]-N-(2-methylpropyl)-1-azabicyclo[2.2.2]octan-3-amine, L-tartrate

¹HNMR (300 MHz, CD₃OD) δ_{H} : 7.40-7.08 (m, 7H), 6.68-6.91 (dt, 1H), 4.29 (s, 2H), 3.56-4.0 (q, 2H), 3.31-2.96 (m, 5H), 2.72-2.67 (t, 1H), 2.58-2.52 (dq, 1H), 2.18-1.30 (m, 8H), 0.70-0.68 (dd, 6H); MS: [M+H] = 367.

The following examples illustrate compounds of of Formulae (IG) above and methods for their preparation.


Preparation of Intermediates

(2S)-(4-Benzyl-morpholin-2-yl)-phenyl-methanone



Described above in section entitled "Preparation of intermediates for the synthesis of Examples 1C-17C".


(S)-Phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methanol (2)



Described above in section entitled "Preparation of intermediates for the synthesis of Examples 1C-17C".


(2S)-2-[(R)-bromo(phenyl)methyl]-4-(phenylmethyl)morpholine (3)

251




To a solution of (S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methanol (2) (4.71 g, 16.63 mmole) in chloroform (200 ml) is added the triphenylphosphine dibromide (14.04 g, 33.26 mmole). The mixture is heated at 60°C overnight. The mixture is allowed to cool to room temperature then washed with saturated sodium carbonate solution (aqueous, ~100 ml), dried (Na_2SO_4) and concentrated *in vacuo*. The resulting residue is purified by automated flash chromatography (ISCO system: 120 g column, 10-30% EtOAc in isohexane) to give (2S)-2-[(R)-bromo(phenyl)methyl]-4-(phenylmethyl)morpholine (3) as a white solid (4.63 g, 80%). LCMS 6 min gradient method, $R_t = 2.5$ min, $(\text{M}+\text{H}^+) = 346/348$

S-{(S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methyl} ethanethioate (5)




A solution of (2S)-2-[(R)-bromo(phenyl)methyl]-4-(phenylmethyl)morpholine (3) (1.76 g, 5.08 mmole) and potassium thioacetate (1.16 g, 10.16 mmole) in 1:1 anhydrous THF:DMF (30 ml), is stirred at 40 °C under nitrogen overnight. The mixture is then taken up in acetonitrile and loaded onto an SC10-2 column (4 x 10 g). The SC10-2 columns are washed with further acetonitrile. The target compound is eluted with 4:1 acetonitrile : Et_3N . This is concentrated *in vacuo* to give an orange oil which is purified by automated flash chromatography (ISCO system: 35 g SiO_2 Redisep column, 10-30% EtOAc in isohexane over 40 minutes) to give S-{(S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methyl} ethanethioate (5) as an amber coloured crystalline solid (1.54 g, 89%). LCMS 6 min gradient method, $R_t = 2.5$ min, $(\text{M}+\text{H}^+) = 342$

(S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6)

- The *S*-{(S)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methyl} ethanethioate (5) (11.02 g, 32.3 mmole) is taken up in methanol (100 ml, dry, degassed), under nitrogen. To this is added the sodium thiomethoxide (2.26 g, 32.3 mmole) in one portion (as solid). The reaction mixture is left to stir at room temperature for 2 hours. The solution is then added to an aqueous solution of HCl (0.1 M). This is extracted with DCM (3 x). The extracts are dried (Na₂SO₄) and concentrated *in vacuo* to give (S)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) as a yellow solid (9.59 g, 99%). LCMS 6 min gradient method, Rt = 2.7 min, (M+H⁺) = 300

Examples

15 **Example 1G: (2S)-2-{(S)-phenyl[(3-phenylpyridin-2-yl)thio]methyl}morpholine hemifumarate**



- i) To palladium acetate (0.026 g, 0.12 mmole) in acetonitrile (3 ml), is added triphenylphosphine (0.122 g, 0.46 mmole), under nitrogen, at room temperature. The mixture is left to stir for 15 minutes. To this mixture is added water (distilled, 1 ml), phenylboronic acid (0.846 g, 6.94 mmole), 3-bromo-2-fluoropyridine (1.02 g, 5.78 mmole) and potassium carbonate (4.80 g, 34.70 mmole). The reaction mixture is heated at 70 °C overnight. After cooling to room temperature, the reaction mixture is loaded

directly onto a 40 g Redisep SiO₂ column and components isolated by automated flash chromatography (ISCO System, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes). This gave 2-fluoro-3-phenylpyridine as a very pale yellow oil (1.00 g, 100 %). LCMS 6 min gradient method, Rt = 3.7 min, (M+H⁺) = 174.


5

ii) To a solution of (S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) (1.50 g, 5.01 mmole) and 2-fluoro-3-phenylpyridine (2.44 g, 14.09 mmole) in dry, degassed DMF (10 ml) is added, under nitrogen, sodium hydride (60 % dispersion in oil, 0.24 g, 6.01 mmole). The mixture is left to stir overnight at room temperature. The reaction mixture is loaded neat onto a 120 g SiO₂ Redisep column (preconditioned with cyclohexane). Automated flash chromatography (ISCO System, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes at 40 ml/minute flow rate) yielded an orange oil (2.26 g). Chromatography is repeated using chromatography (ISCO System, 40 g column, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes at 30 ml/minute flow rate) to give (2S)-2-{(S)-phenyl[(3-phenylpyridin-2-yl)thio]methyl}-4-(phenylmethyl)morpholine as a pale orange oil (1.65 g, 73 %). LCMS 6 min gradient method, Rt = 4.0 min, (M+H⁺) = 453.

iii) To a suspension of polymer supported diisopropylamine (3.78 mmol/g, 0.54 g, 2.03 mmole) and (2S)-2-{(S)-phenyl[(3-phenylpyridin-2-yl)thio]methyl}-4-(phenylmethyl)morpholine (0.184 g, 0.41 mmole) in dry DCM (5 ml) is added 1-chloroethyl chloroformate (0.22 ml, 2.03 mmole) at room temperature and under nitrogen. The mixture is heated at 40°C for 3.75 hours. The reaction mixture is filtered, concentrated *in vacuo* then taken up in methanol (5 ml). The solution is left to stir at room temperature overnight. After this time, the reaction mixture is loaded directly onto an SC10-2 column. The SC10-2 column is washed with methanol. The title compound is eluted with 2 N NH₃/methanol. This is concentrated *in vacuo* to give (2S)-2-{(S)-phenyl[(3-phenylpyridin-2-yl)thio]methyl}morpholine as white foam (0.148 g, 100 %). The foam is taken up in ethyl acetate. To this is added a solution of fumaric acid (1.1 equiv, 0.052 g) in methanol. The resulting solution is filtered then concentrated *in vacuo*. To the resulting white solid is added methanol (1.5 ml). This is stirred for a couple of

minutes, then the remaining solid collected by filtration to give the hemi-fumarate salt of (2*S*)-2-[(*S*)-phenyl[(3-phenylpyridin-2-yl)thio]methyl]morpholine as a white solid (0.127 g). LCMS 12 min gradient method, $R_t = 5.5$ min, $(M+H^+) = 363$

5 **Example 2G: (2*S*)-2-[(*S*)-{3-(4-fluorophenyl)pyridin-2-yl}thio}(phenyl)methyl]morpholine fumarate**




- i) To bis(benzonitrile)palladium(II)dichloride (0.054 g, 0.14 mmole) and 1,4-bis(diphenylphosphine)butane (0.091 g, 0.21 mmole) is added dry toluene (6 ml), under nitrogen, and the mixture stirred for half an hour. To this is added 3-bromo-2-fluoropyridine (0.50 g, 2.83 mmole) in ethanol (1.4 ml) followed by a solution of 4-fluorophenylboronic acid (0.793 g, 5.67 mmole) in ethanol (2.4 ml). To this is added an aqueous solution of sodium carbonate (1 M, 2.83 ml, 2.83 mmole). The mixture is heated at 60°C for 24 hours, then at 75°C for a further 16 hours. The organic layer is loaded directly onto a 40 g Redisep SiO₂ column and components isolated by automated flash chromatography (ISCO System, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes). This gave 3-(4-fluorophenyl)-2-fluoropyridine as a white solid (0.387 g, 71 %). LCMS 6 min gradient method, $R_t = 3.6$ min, $(M+H^+) = 192$
- ii) To a solution of (*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) (0.505 g, 1.69 mmole) and 3-(4-fluorophenyl)-2-fluoropyridine (0.387 g, 2.02 mmole) in dry, degassed DMF (3 ml) is added, under nitrogen, cesium fluoride (0.385 g, 2.54 mmole). The mixture is heated at 65°C over the weekend. After this time, the reaction mixture is allowed to cool and loaded directly onto an SC10-2 column. The SC10-2 column is washed with methanol. The (2*S*)-2-[(*S*)-{3-(4-fluorophenyl)pyridin-2-yl}thio}(phenyl)methyl]-4-(phenylmethyl)morpholine is eluted with 2 N NH₃/methanol. This is concentrated *in vacuo* to give an orange solid (0.649 g). This is purified by

automated flash chromatography (ISCO System, 40 g SiO₂ Redisep column, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes at 30 ml/minute flow rate) to give (2*S*)-2-[(*S*)-{[3-(4-fluorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-4-(phenylmethyl)morpholine as a off-white foam (0.395 g, 50 %). LCMS 6 min gradient
 5 method, Rt = 3.3 min, (M+H⁺) = 471.

iii) Deprotection of the morpholine nitrogen is carried out using the method and work up as described in Example 1G, using polymer supported diisopropylamine (3.78 mmole/g, 1.09 g, 4.14 mmole), (2*S*)-2-[(*S*)-{[3-(4-fluorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-
 10 4-(phenylmethyl)morpholine (0.390 g, 0.83 mmole), dry DCM (20 ml), 1-chloroethyl chloroformate (0.45 ml, 4.14 mmole) and methanol (20 ml). This gave (2*S*)-2-[(*S*)-{[3-(4-fluorophenyl)pyridin-2-yl]thio}(phenyl)methyl]morpholine as a pale yellow oil (0.232 g, 74 %). This oil is taken up in ethyl acetate. To this is added a solution of fumaric acid (1.1 equiv, 0.071 g) in methanol. The resulting solid is collected by filtration to give a white
 15 solid (0.115 g). This is recrystallised from MeOH/CHCl₃/Et₂O to give a white solid (0.061 g). LCMS 12 min gradient method, Rt = 5.4 min, (M+H⁺) = 381

Example 3G: (2*S*)-2-[(*S*)-{[3-(3-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]morpholine fumarate



20

i) To bis(benzonitrile)palladium(II)dichloride (0.054 g, 0.14 mmole) and 1,4-bis(diphenylphosphine)butane (0.091 g, 0.21 mmole) is added dry toluene (6 ml), under nitrogen, and the mixture stirred for half an hour. To this is added 3-bromo-2-fluoropyridine (0.50 g, 2.83 mmole) in ethanol (1.4 ml) followed by a solution of 3-chlorophenylboronic acid (0.887 g, 5.67 mmole) in ethanol (2.4 ml). To this is added an
 25 aqueous solution of sodium carbonate (1 M, 2.83 ml, 2.83 mmole). The mixture is heated at 60°C for 24 hours, then at 75°C for a further 16 hours. The organic layer is loaded

directly onto a 40 g Redisep SiO₂ column and components isolated by automated flash chromatography (ISCO System, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes). This gave 3-(3-chlorophenyl)-2-fluoropyridine as an off-white solid (0.333 g, 57 %). LCMS 6 min gradient method, Rt = 4.0 min, (M+H⁺) = 208.

5

ii) To a solution of (S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) (0.400 g, 1.34 mmole) and 3-(3-chlorophenyl)-2-fluoropyridine (0.333 g, 1.60 mmole) in dry, degassed DMF (3 ml) is added, under nitrogen, cesium fluoride (0.305 g, 2.00 mmole). The mixture is heated at 65°C over the weekend. After this time, the reaction

10 mixture allowed to cool. The resulting solid is taken up in MeOH/DCM and loaded directly onto an SC10-2 column. The SC10-2 column is washed with methanol. The (2S)-2-[(S)-{[3-(3-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-4-(phenylmethyl)morpholine is eluted with 2 N NH₃/methanol. This is concentrated *in vacuo* to give a white foam (0.555 g). This is purified by automated flash chromatography (ISCO System, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes at 40 ml/minute flow rate) to yield (2S)-2-[(S)-{[3-(3-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-4-(phenylmethyl)morpholine as a white foam (0.258 g, 40 %). LCMS 6 min gradient method, Rt = 4.2 min, (M+H⁺) = 487.


20 iii) Deprotection of the morpholine nitrogen is carried out using the method and work up as described in Example 1G, using polymer supported diisopropylamine (3.72 mmole/g, 0.70 g, 1.80 mmole), (2S)-2-[(S)-{[3-(3-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-4-(phenylmethyl)morpholine (0.255 g, 0.52 mmole), dry DCM (15 ml), 1-chloroethyl chloroformate (0.29 ml, 2.62 mmole) and methanol (15 ml). This gave a colourless

25 residue (0.211 g). This residue is taken up in ethyl acetate. To this is added a solution of fumaric acid (1.1 equiv, 0.062 g) in methanol. If the resulting solid contains impurities it may be recombined with the mother liquor and purified on a UV Guided PrepHPLC (Flex) System and treated with SC10-2 to give (2S)-2-[(S)-{[3-(3-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]morpholine as a pale yellow oil (0.127 g, 65 %). This oil is taken

30 up in MeOH/DCM. To this is added a solution of fumaric acid (1.1 equiv, 0.0145 g) in methanol, followed by Et₂O. The resulting crystals are collected by filtration to give the

fumarate salt of (2*S*)-2-[(*S*)-{[3-(3-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]morpholine (1:1 fumarate salt) as a white solid (0.047 g). LCMS 12 min gradient method, $R_t = 5.7$ min, $(M+H^+) = 397$

5 **Example 4G: (2*S*)-2-[[3-(2-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]morpholine fumarate**



- i) To palladium acetate (0.0025 g, 0.0011 mmole) in acetonitrile (3 ml), is added triphenylphosphine (0.0119 g, 0.045 mmole), under nitrogen, at room temperature. The mixture is left to stir for 15 minutes. To this mixture is added water (distilled, 1 ml), 2-chlorophenylboronic acid (0.106 g, 0.68 mmole), 3-bromo-2-fluoropyridine (0.10 g, 0.57 mmole) and potassium carbonate (0.470 g, 3.40 mmole). The reaction mixture is heated to 60°C increasing to 75 °C over 5 hours then allowed to cool to room temperature. To the reaction mixture is added MeOH and this is loaded onto an SC10-2 column (10 g)
- 10
- 15 preconditioned with MeOH. The column is washed with MeOH and the resulting solution concentrated *in vacuo* to give an orange oil (0.196 g). The oil is purified by automated flash chromatography (ISCO System, a 10 g Redisep SiO₂ column, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes). This gave 2-fluoro-3-(2-chlorophenyl)pyridine as a colourless oil (0.050 g, 42 %). LCMS 6 min gradient method,
- 20 $R_t = 3.3$ min, $(M+H^+) = 208$


- ii) To a solution of (*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) (0.288g, 0.96 mmole) and 3-(2-chlorophenyl)-2-fluoropyridine (0.40 g, 1.93 mmole) in dry, degassed DMF (2 ml) is added, under nitrogen, sodium hydride (60% dispersion in oil, 0.046 g, 1.15 mmole). The mixture is left to stir at room temperature over the weekend. The reaction mixture is loaded directly onto an a 40 g Redisep SiO₂ column. Components are eluted using automated flash chromatography (ISCO System, 0 – 30 %
- 25

ethyl acetate in cyclohexane gradient elution over 30 minutes at 40 ml/minute flow rate) to give (2*S*)-2-[[3-(2-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-4-(phenylmethyl)morpholine as a white solid (0.021 g, 5 %). LCMS 6 min gradient method, $R_t = 4.3$ min, $(M+H^+) = 487$.

5

iii) Deprotection of the morpholine nitrogen is carried out using the method and work up as described in Example 1G, using polymer supported diisopropylamine (3.78 mmole/g, 0.057 g, 0.216 mmole), (2*S*)-2-[[3-(2-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-4-(phenylmethyl)morpholine (0.021 g, 0.043 mmole), dry DCM (2 ml), 1-chloroethyl
 10 chloroformate (0.024 ml, 0.216 mmole) and methanol (2 ml). This gave a colourless residue (0.017 g, 100 %). This residue is taken up in ethyl acetate. To this is added a solution of fumaric acid (1 equiv, 0.005 g) in methanol. This is reduced in volume and Et₂O added. The resulting solid is collected by filtration to give the fumarate salt of (2*S*)-2-[[3-(2-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl)morpholine (1:1 fumarate salt)
 15 as a pale green solid (0.012 g). LCMS 12 min gradient method, $R_t = 5.4$ min, $(M+H^+) = 397$

Example 5G: (2*S*)-2-((*S*)-phenyl{3-(trifluoromethyl)pyridin-2-yl}thio)methyl)morpholine




i) Potassium fluoride (0.048 g, 0.84 mmole) and copper (I) iodide (0.159 g, 0.84 mmole) are thoroughly mixed and dried under reduced pressure with a hot air gun for 20 minutes. To the resulting yellow solid, at room temperature is added (2*S*)-2-[(*S*)-[(3-iodopyridin-2-yl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine (as prepared in Example 15) (0.190
 25 g, 0.38 mmole) in anhydrous NMP (0.5 ml) followed by anhydrous DMF (0.5 ml) then (trifluoromethyl)trimethylsilane (0.11 ml, 0.76 mmole). After 3 days at room temperature, the temperature is increased to 50 °C. The reaction mixture is heated at 50 °C overnight.

After cooling to room temperature, further (trifluoromethyl)trimethylsilane (0.11 ml, 0.76 mmole) is added to the reaction mixture and the mixture is left to stir overnight at room temperature. To the reaction mixture is added MeOH before loading onto an SC10-2 column (10 g) preconditioned with MeOH. The column is washed with MeOH. Basic material is eluted with 2 N NH₃/methanol. This is concentrated *in vacuo* to give a pale yellow solid (0.199 g). This is purified by automated flash chromatography (ISCO System, 3 x 4 g Redisep SiO₂ columns, in parallel, 0 – 20 % ethyl acetate in cyclohexane gradient elution over 40 minutes) to give the (2*S*)-2-[(*S*)-[3-iodopyridin-2-yl]thio](phenyl)methyl]-4-(phenylmethyl)morpholine as a white foam (0.108 g, 57 % recovery of this starting material) and (2*S*)-2-((*S*)-phenyl{[3-(trifluoromethyl)pyridin-2-yl]thio}methyl)-4-(phenylmethyl)morpholine as a colourless oil (0.033 g, 20 %). LCMS 6 min gradient method, Rt = 4.2 min, (M+H⁺) = 445

ii) To a suspension of polymer supported diisopropylamine (3.72 mmol/g, 0.097 g, 0.36 mmole) and (2*S*)-2-((*S*)-phenyl{[3-(trifluoromethyl)pyridin-2-yl]thio}methyl)-4-(phenylmethyl)morpholine (0.032 g, 0.07 mmole) in dry DCM (0.5 ml) is added 1-chloroethyl chloroformate (0.039 ml, 0.36 mmole) at room temperature and under nitrogen. The mixture is heated at 40 °C for 2 hours. The reaction mixture is filtered and concentrated *in vacuo* then taken up in methanol (0.5 ml). The solution left to stir at room temperature overnight. After this time, the reaction mixture is loaded directly onto an SC10-2 column. The SC10-2 column is washed with methanol. The target compound is eluted with 2 N NH₃/methanol. This is concentrated *in vacuo* to give a pale yellow oil (0.024 g). The pale yellow oil is purified using an automated PrepLCMS system, then liberated as the free base by treatment with SC10-2 and concentrated under vacuum to give (2*S*)-2-((*S*)-phenyl{[3-(trifluoromethyl)pyridin-2-yl]thio}methyl)morpholine as a white solid (0.005 g, 20 %). LCMS 12 min gradient method, Rt = 4.9 min, (M+H⁺) = 354

Example 6G: (2*S*)-2-((*S*)-phenyl{[3-(phenylmethyl)pyridin-2-yl]thio}methyl)morpholine fumarate

260




- i) To a 100 ml round-bottomed flask, under nitrogen, containing dry THF (25 ml) is added *n*-butyllithium (1.6 M solution in hexanes, 3.99 ml, 6.39 mmole) at 0°C followed by lithium diisopropylamide (2 M solution in THF/*n*-heptane, 3.19 ml, 6.39 mmole). The reaction mixture is left to stir for 1 hour at 0°C. The mixture is cooled to -70°C then 2-fluoropyridine added. The solution is stirred at -70°C for 4 hours. To the solution is added benzaldehyde (0.71 ml, 6.97 mmole). This is then left to stir for 1 hour at -70°C, after which time water (100 ml) is added. On warming to room temperature the solution is extracted with chloroform (2 x 100 ml). The combined extracts are dried (Na₂SO₄) and concentrated *in vacuo* to yield a yellow oil (1.58 g). Purification by automated flash chromatography (ISCO System, Redisep 10 g SiO₂ column, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 30 minutes at 20 ml/min flow rate) gave 2-fluoro-3-(phenyl-1-hydroxymethyl)pyridine as a yellow oil (0.71 g, 59 %). FIA (M+H⁺) = 204
- ii) To 5 % Pd/C (0.07 g), under nitrogen, is added a solution of 2-fluoro-3-(1-hydroxy-1-phenylmethyl)pyridine (0.71 g, 3.5 mmole) in ethanol (50 ml). This is then put on a Parr Hydrogenator at 60 psi H₂ and left over the weekend. The reaction mixture is filtered through Celite[®]. Removal of solvent from the resulting solution gave a pale yellow oil. This is purified by automated flash chromatography (ISCO System, 10 g SiO₂ Redisep column, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes at 20 ml/minute flow rate) to give 2-fluoro-3-(phenylmethyl)pyridine as a colourless oil (0.18 g, 27 %).
- iii) To a solution of (*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) (0.27 g, 0.91 mmole) and 2-fluoro-3-(1-hydroxy-1-phenylmethyl)pyridine (0.17 g, 0.91 mmole) in dry, degassed DMF (1.5 ml) is added, under nitrogen, sodium hydride (60 % dispersion in oil, 0.07 g, 1.82 mmole). The mixture is left to stir overnight at room

temperature. A further portion of sodium hydride (605 dispersion in oil, 0.07 g, 1.82 mmole) and DMF (1 ml) is added. After 5 hours at room temperature, the reaction mixture is taken up in MeOH and loaded onto an SC10-2 column. The SC10-2 column is washed with methanol. The (2*S*)-2-((*S*)-phenyl{[3-(phenylmethyl)pyridin-2-yl]thio}methyl)-4-(phenylmethyl)morpholine is eluted with 2 N NH₃/methanol. This is concentrated *in vacuo* to give a yellow residue (0.36 g). The residue is purified by automated flash chromatography (ISCO System, 35 g SiO₂ Redisep column, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes at 40 ml/minute flow rate) which yields (2*S*)-2-((*S*)-phenyl{[3-(phenylmethyl)pyridin-2-yl]thio}methyl)-4-(phenylmethyl)morpholine as a pale yellow oil (0.10 g, 24 %). LCMS 6 min gradient method, Rt = 3.8min, (M+H⁺) = 467

iv) Deprotection of the morpholine nitrogen is carried out using the method and work up as described in Example 1G, using polymer supported diisopropylamine (3.78 mmole/g, 0.28 g, 1.07 mmole), of (2*S*)-2-((*S*)-phenyl{[3-(phenylmethyl)pyridin-2-yl]thio}methyl)-4-(phenylmethyl)morpholine (0.092 g, 0.20 mmole), dry DCM (5 ml), 1-chloroethyl chloroformate (0.12 ml, 1.07 mmole) and methanol (5 ml). This gives (2*S*)-2-((*S*)-phenyl{[3-(phenylmethyl)pyridin-2-yl]thio}methyl)morpholine as a colourless residue (0.076 g, 94 %). This oil is taken up in ethyl acetate. To this is added a solution of fumaric acid (1.1 equiv, 0.026 g) in methanol. The resulting solution is concentrated *in vacuo* and the resulting oil triturated with ethyl acetate. The solid is collected by filtration to give the fumarate salt of (2*S*)-2-((*S*)-phenyl{[3-(phenylmethyl)pyridin-2-yl]thio}methyl)morpholine (1:1 fumarate salt) as a white solid (0.070 g). LCMS 12 min gradient method, Rt = 5.6 min, (M+H⁺) = 377

Example 7G: (2*S*)-2-((*S*)-phenyl{[3-(phenyloxy)pyridin-2-yl]thio}methyl)morpholine fumarate

262




i) To a 100 ml round bottomed flask is added 2-chloro-3-pyridinol (0.50 g, 3.86 mmole), copper (II) acetate (0.70 g, 3.86 mmole), phenylboronic acid (0.94 g, 7.72 mmole) and powdered 4Å molecular sieves. To the mixture is added DCM (39 ml) followed by triethylamine (2.69 ml, 19.30 mmole). This is stirred overnight, under nitrogen, at room temperature. The reaction mixture is poured into water (75 ml) and extracted with ethyl acetate (3 x 75 ml). The combined extracts are concentrated *in vacuo* to give a brown oil (0.65 g). Purification by automated flash chromatography (ISCO System, Redisep 35 g SiO₂ column, 0 – 20 % ethyl acetate in cyclohexane gradient elution over 40 minutes) gives 2-chloro-3-phenoxy-pyridine as a colourless oil (0.32 g, 41%). LCMS 6 min gradient method, Rt = 3.6min, (M+H⁺) = 206

ii) To a solution of (S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) (0.352 g, 1.18 mmole) and 2-chloro-3-phenoxy-pyridine (0.29 g, 1.41 mmole) in dry, degassed DMF (3 ml) is added, under nitrogen, cesium fluoride (0.179 g, 1.18 mmole). The mixture is left to stir for two days at 55°C. A further portion of cesium fluoride (0.063 g, 0.41 mmole) is added and the solution heated for 5 hours at 55°C. The reaction mixture is allowed to cool then loaded neat onto a 35 g SiO₂ Redisep column (preconditioned with cyclohexane). Automated flash chromatography (ISCO System, 0 – 40 % ethyl acetate in cyclohexane gradient elution over 40 minutes at 30 ml/minute flow rate) yields a yellow oil (2.26 g). This is taken up in MeOH and loaded onto an SC10-2 column. The SC10-2 column is washed with methanol. The title compound is eluted with 2 N NH₃/methanol. This is concentrated *in vacuo* to give (2S)-2-[(S)-phenyl[(3-phenoxy-pyridin-2-yl)thio]methyl]-4-(phenylmethyl)morpholine as a pale orange oil (0.092 g, 17 %). LCMS 6 min gradient method, Rt = 3.6 min, (M+H⁺) = 469

iii) Deprotection of the morpholine nitrogen is carried out using the method and work up as described in Example 1G, using polymer supported diisopropylamine (3.78 mmole/g, 0.26 g, 0.98 mmole), (2*S*)-2-[(*S*)-phenyl[(3-phenyloxy)pyridin-2-yl]thio]methyl]-4-(phenylmethyl)morpholine (0.092 g, 0.20 mmole), dry DCM (5 ml), 1-chloroethyl chloroformate (0.11 ml, 0.98 mmole) and methanol (5 ml). This gave (2*S*)-2-[(*S*)-phenyl{[3-(phenyloxy)pyridin-2-yl]thio}methyl)morpholine as a pale yellow oil (0.070 g, 95 %). This oil is taken up in ethyl acetate. To this is added a solution of fumaric acid (1.1 equiv, 0.024 g) in methanol. The resulting solution is concentrated *in vacuo* and the resulting oil triturated with ethyl acetate. The solid is collected by filtration to give the fumarate salt of (2*S*)-2-[(*S*)-phenyl{[3-(phenyloxy)pyridin-2-yl]thio}methyl)morpholine (1:1 fumarate salt) as an off-white solid (0.094 g). LCMS 12 min gradient method, *R*_t = 5.5 min, (*M*+*H*⁺) = 379

Example 8G: (2*S*)-2-[(*S*)-[(3-chloropyridin-2-yl)thio](phenyl)methyl]morpholine


fumarate



i) To a solution of (*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) (0.446 g, 1.49 mmole) and 2,3-dichloropyridine (0.246 g, 1.67 mmole) in dry, degassed DMF (3 ml) is added, under nitrogen, sodium hydride (60 % dispersion in oil, 0.061 g, 1.53 mmole). The mixture is left to stir overnight at room temperature. The reaction mixture is taken up in MeOH and loaded onto an SC10-2 column. The SC10-2 column is washed with methanol. The (2*S*)-2-[(*S*)-[(3-chloropyridin-2-yl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine is eluted with 2 N NH₃/methanol. This is concentrated *in vacuo* to give (2*S*)-2-[(*S*)-[(3-chloropyridin-2-yl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine as a pale yellow oil (0.61 g). LCMS 6 min gradient method, *R*_t = 3.5 min, (*M*+*H*⁺) = 411

ii) Deprotection of the morpholine nitrogen is carried out using the method and work up as described in Example 1G, using polymer supported diisopropylamine (3.78 mmole/g, 0.39g, 1.46 mmole), (2*S*)-2-[(*S*)-[(3-chloropyridin-2-yl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine (0.120 g, 0.292 mmole), dry DCM (15 ml), 1-chloroethyl chloroformate (0.16 ml, 1.46 mmole) and methanol (15 ml). This gives (2*S*)-2-[(*S*)-[(3-chloropyridin-2-yl)thio](phenyl)methyl]morpholine as a pale yellow oil (0.092 g, 98 %). This oil is taken up in ethyl acetate. To this is added a solution of fumaric acid (1 equiv, 0.033 g) in methanol. The resulting solution is concentrated *in vacuo* to give an oil which is crystallised from IPA. The solid is collected by filtration to give the fumarate salt of (2*S*)-2-[(*S*)-[(3-chloropyridin-2-yl)thio](phenyl)methyl]morpholine (1:1 fumarate salt) as a white solid (0.111 g). LCMS 12 min gradient method, *R*_t = 4.8 min, (*M*+*H*⁺) = 321

Example 9C: (2*S*)-2-[(*S*)-[(3-methylpyridin-2-yl)thio](phenyl)methyl]morpholine fumarate




i) To a degassed solution of *S*-{(*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methyl} ethanethioate (5) (0.100 g, 0.293 mmole) and 2-fluoro-3-methylpyridine (0.325 g, 2.93 mmole) in DMF (1 ml) is added sodium methoxide (0.016 g, 0.293 mmole). The reaction mixture is left to stir at room temperature, under nitrogen, overnight. The reaction mixture is diluted with methanol and loaded onto an SC10-2 (5 g) column preconditioned with MeOH. The column is washed with MeOH then basic material is eluted with 2 N NH₃/methanol. This ammonia solution is concentrated *in vacuo* to give an orange oil (0.067 g) which is purified by automated flash chromatography (ISCO System, Redisep SiO₂ column, 0 – 20 % ethyl acetate in cyclohexane gradient elution over 40 minutes) to give (2*S*)-2-[(*S*)-[(3-methylpyridin-2-yl)thio](phenyl)methyl]-4-

(phenylmethyl)morpholine as a colourless oil (0.055 g, 44%). LCMS 6 min gradient method, $R_t = 2.9$ min, $(M+H^+) = 391$

- ii) To a suspension of polymer supported diisopropylamine (3.78 mmol/g, 0.167 g, 0.64 mmole) and (2*S*)-2-[(*S*)-[(3-methylpyridin-2-yl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine (0.050 g, 0.13 mmole) in dry DCM (5 ml) is added 1-chloroethyl chloroformate (0.070 ml, 0.64 mmole) at room temperature and under nitrogen. The mixture is heated at 40°C for 1.5 hours. The reaction mixture is filtered and concentrated *in vacuo* then taken up in methanol (5 ml). The solution is left to stir at room temperature for 2.5 hours. After this time, the reaction mixture is loaded directly onto an SC10-2 column. The SC10-2 column is washed with methanol. The free base of the title compound is eluted with 2 N NH_3 /methanol. This ammonia solution is concentrated *in vacuo* to give (2*S*)-2-[(*S*)-[(3-methylpyridin-2-yl)thio](phenyl)methyl]morpholine as an orange oil (0.037 g, 97 %). This oil is taken up in methanol. To this is added a solution of fumaric acid (1 equiv, 0.014 g) in methanol. This is stirred for a couple of minutes, then EtOAc followed by isohexane added. The resulting precipitate is collected by filtration to yield a white solid (0.048 g). This is recrystallised from ethyl acetate and isohexane to give the fumarate salt of (2*S*)-2-[(*S*)-[(3-methylpyridin-2-yl)thio](phenyl)methyl]morpholine (1:1 fumarate salt) as a white solid (0.013 g) LCMS 12 min gradient method, $R_t = 4.5$ min, $(M+H^+) = 301$

Example 10G: (2*S*)-2-[(*S*)-[3-(4-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]morpholine fumarate




- i) To bis(benzonitrile)palladium(II)dichloride (0.054 g, 0.14 mmole) and 1,4-bis(diphenylphosphine)butane (0.091 g, 0.21 mmole) is added dry toluene (6 ml), under

- nitrogen, and the mixture stirred for half an hour. To this is added 3-bromo-2-fluoropyridine (0.50 g, 2.83 mmole) in ethanol (1.4 ml) followed by a solution of 4-chlorophenylboronic acid (0.887 g, 5.67 mmole) in ethanol (2.4 ml). To this is added an aqueous solution of sodium carbonate (1 M, 2.83 ml, 2.83 mmole). The mixture is heated at 60°C for 24 hours, then at 75°C for a further 16 hours. The organic layer is loaded directly onto a 40 g Redisep SiO₂ column and components isolated by automated flash chromatography (ISCO System, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes). This gave 3-(4-chlorophenyl)-2-fluoropyridine as a white solid (0.323 g, 55 %). LCMS 6 min gradient method, Rt = 4.0 min, (M+H⁺) = 208
- ii) To a solution of (S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) (0.388 g, 1.30 mmole) and 3-(4-chlorophenyl)-2-fluoropyridine (0.323 g, 1.56 mmole) in dry, degassed DMF (3 ml) is added, under nitrogen, cesium fluoride (0.295 g, 1.94 mmole). The mixture is heated at 65°C over the weekend. After this time, the reaction mixture is allowed to cool. The resulting solid is taken up in MeOH/DCM and loaded directly onto an SC10-2 column. The SC10-2 column is washed with methanol followed by 2 N NH₃/methanol. The ammonia solution is concentrated *in vacuo* to give (2S)-2-[(S)-{[3-(4-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-4-(phenylmethyl)morpholine as an orange foam (0.514 g). This is purified by automated flash chromatography (ISCO System, 0 – 30 % ethyl acetate in cyclohexane gradient elution over 40 minutes at 40 ml/minute flow rate) to give (2S)-2-[(S)-{[3-(4-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-4-(phenylmethyl)morpholine as a white foam (0.178 g, 28 %). LCMS 6 min gradient method, Rt = 4.2 min, (M+H⁺) = 487
- iii) Deprotection of the morpholine nitrogen is carried out using the method and work up as described in Example 1G, using polymer supported diisopropylamine (3.78 mole/g, 0.48 g, 1.80 mmole), (2S)-2-[(S)-{[3-(4-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]-4-(phenylmethyl)morpholine (0.175 g, 0.36 mmole), dry DCM (10 ml), 1-chloroethyl chloroformate (0.20 ml, 1.80 mmole) and methanol (10 ml). This gave a colourless residue (0.129 g, 90 %). This residue is taken up in ethyl acetate. To this is added a solution of fumaric acid (1.1 equiv, 0.035 g) in methanol. The resulting solid is

recombined with the mother liquor and purified on a UV Guided PrepHPLC (Flex) System and treated with SC10-2 to give a yellow solid. This is further purified by automated flash chromatography (ISCO System, Redisep 4 g SiO₂ column, 0 – 5 % methanol in dichloromethane gradient elution over 40 minutes, then 10 minutes at 5 % Methanol in dichloromethane with 10 ml/min flow rate) to give (2*S*)-2-[(*S*)-{[3-(4-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]morpholine as a pale yellow oil (0.049 g, 34 %). This oil is taken up in ethyl acetate. To this is added a solution of fumaric acid (1.1 equiv, 0.0145 g) in methanol. The resulting solution is concentrated *in vacuo* and recrystallised from MeOH and Et₂O. The solid is collected by filtration to give the fumarate salt of (2*S*)-2-[(*S*)-{[3-(4-chlorophenyl)pyridin-2-yl]thio}(phenyl)methyl]morpholine (1:1 fumarate salt) as a white solid (0.047 g). LCMS 12 min gradient method, Rt = 5.7 min, (M+H⁺) = 397

Example 11G: (2*S*)-2-[(*S*)-[(5-bromopyridin-2-yl)thio](phenyl)methyl]morpholine fumarate




i) To a solution of *S*-{(*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methyl} ethanethioate (5) (0.25 g, 0.73 mmole) in dry methanol (2 ml) is added sodium methoxide (0.040 g, 0.73 mmole) under nitrogen. This is left to stir at room temperature for 1 hour. Methanol is removed *in vacuo* and replaced with DMF (1 ml). To this is then added the 5-bromo-2-fluoropyridine (0.11 ml, 1.02 mmole). The reaction mixture is left to stir at room temperature, under nitrogen, overnight. The reaction mixture is diluted with DCM and loaded directly onto a 35 g Redisep column. Purification by automated flash chromatography (ISCO System, Redisep 35 g SiO₂ column, 0 – 20 % ethyl acetate in cyclohexane gradient elution over 40 minutes) gave (2*S*)-2-[(*S*)-[(5-bromopyridin-2-

yl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine as a colourless oil (0.186 g, 56%).
LCMS 6 min gradient method, $R_t = 3.6$ min, $(M+H^+) = 455/457$

- ii) To a suspension of polymer supported diisopropylamine (3.78 mmol/g, 0.108 g, 20.4 mmole) and (2*S*)-2-[(*S*)-[(5-bromopyridin-2-yl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine (0.186 g, 0.408 mmole) in dry DCM (10 ml) is added 1-chloroethyl chloroformate (0.22 ml, 2.04 mmole) at room temperature and under nitrogen. The mixture is heated at 40°C for 2.5 hours. The reaction mixture is then filtered and concentrated *in vacuo* then taken up in methanol (10 ml). The solution is left to stir at room temperature overnight. After this time, the reaction mixture is loaded directly onto an SC10-2 column (5 g). The SC10-2 column is washed with methanol. The target compound is eluted with 2 N NH_3 /methanol. This is concentrated *in vacuo* to give (2*S*)-2-[(*S*)-[(5-bromopyridin-2-yl)thio](phenyl)methyl]morpholine as a colourless oil (0.108 g, 72 %). This oil is taken up in ethanol. To this is added a solution of fumaric acid (1.2 equiv, 0.041 g) in ethanol. Solvent is removed in vacuo and the resulting residue triturated with EtOAc. This solid is collected by filtration to give the fumarate salt of (2*S*)-2-[(*S*)-[(5-bromopyridin-2-yl)thio](phenyl)methyl]morpholine (1:1 fumarate salt) as a white solid (0.105 g). LCMS 12 min gradient method, $R_t = 5.0$ min, $(M+H^+) = 365/367$

20 **Example 12G: 2-{[(*S*)-(2*S*)-morpholin-2-yl(phenyl)methyl]thio}pyridine-3-carboxamide fumarate**




- i) To a degassed solution of *S*-{[(*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methyl} ethanethioate (5) (0.100 g, 0.293 mmole) and 2-chloronicotinamide (0.046 g, 0.293 mmole) in ethanol (3 ml) is added a solution of sodium hydroxide in water (2 M, 0.293 ml, 0.586 mmole). The resulting solution is stirred at room temperature overnight. An

- additional portion of 2-chloronicotinamide (0.046 g, 0.293 mmole) is added to the reaction mixture which is then heated at 40 °C overnight. The reaction mixture is diluted with methanol and loaded onto an SC10-2 column preconditioned with MeOH. The column is washed with MeOH then basic material is eluted with 2 N NH₃/methanol. This ammonia solution is concentrated *in vacuo* to give 2-({[(S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methyl}thio]pyridine-3-carboxamide as a pale orange residue (0.124 g, 100%). LCMS 6 min gradient method, Rt = 2.1 min, (M+H⁺) = 420
- 5
- ii) To a suspension of polymer supported diisopropylamine (3.78 mmol/g, 0.38 g, 1.47 mmole) and 2-({[(S)-phenyl[(2S)-4-(phenylmethyl)morpholin-2-yl]methyl}thio]pyridine-3-carboxamide (0.123 g, 0.29 mmole) in dry DCM (10 ml) is added 1-chloroethyl chloroformate (0.16 ml, 1.47 mmole) at room temperature and under nitrogen. The mixture is heated at 40°C for 2 hours. The reaction mixture is then filtered and concentrated *in vacuo* to give a pale yellow residue. This is taken up in methanol (10 ml)
- 10
- 15 and the solution left to stir at room temperature for 3 hours. After this time, the reaction mixture is loaded directly onto an SC10-2 column. The SC10-2 column is washed with methanol then more basic compounds are eluted with 2 N NH₃/methanol. The ammonia solution is concentrated *in vacuo* to give 2-{{[(S)-(2S)-morpholin-2-yl(phenyl)methyl]thio}pyridine-3-carboxamide as a pale yellow oil (0.097 g, 100 %). The
- 20
- 25 pale yellow oil is taken up in methanol. To this is added a solution of fumaric acid (1 equiv, 0.0153 g) in methanol. This is stirred for a couple of minutes, then EtOAc added. The resulting precipitate is collected by filtration to give the fumarate salt of 2-{{[(S)-(2S)-morpholin-2-yl(phenyl)methyl]thio}pyridine-3-carboxamide (1:1 fumarate salt) as a white solid (0.095 g). LCMS 12 min gradient method, Rt = 2.4 min, (M+H⁺) = 330

Example 13G: 2-{{[(S)-(2S)-morpholin-2-yl(phenyl)methyl]thio}pyridine-3-carbonitrile fumarate

270




i) To a degassed solution of *S*-{[(*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methyl} ethanethioate (5) (0.050 g, 0.147 mmole) and 2-chloro-3-cyanopyridine (0.020 g, 0.146 mmol) in ethanol (1.5 ml) is added a solution of sodium hydroxide in water (2 M, 0.146 ml, 0.293 mmole). The resulting solution is stirred at room temperature for ~17 hours. The reaction mixture is diluted with methanol and loaded onto an SC10-2 column preconditioned with MeOH. The column is washed with MeOH then basic material is eluted with 2 N NH₃/methanol. This ammonia solution is concentrated *in vacuo* to give 2-({[(*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methyl}thio)pyridine-3-carbonitrile as an off white solid (0.055 g, 93%). LCMS 6 min gradient method, Rt = 2.8 min, (M+H⁺) = 402

ii) To a suspension of polymer supported diisopropylamine (3.78 mmol/g, 0.181 g, 0.685 mmole) and 2-({[(*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methyl}thio)pyridine-3-carbonitrile (0.055 g, 0.137 mmole) in dry DCM (5 ml) is added 1-chloroethyl chloroformate (0.075 ml, 0.685 mmole) at room temperature and under nitrogen. The mixture is heated at 40°C for 2 hours. The reaction mixture is then filtered and concentrated *in vacuo* to give a pale orange liquid. This is taken up in methanol (5 ml) and the solution left to stir at room temperature overnight. After this time, the reaction mixture is loaded directly onto an SC10-2 column. The SC10-2 column is washed with methanol then more basic material is eluted with 2 N NH₃/methanol. The ammonia solution is concentrated *in vacuo* to give 2-{[(*S*)-(2*S*)-morpholin-2-yl(phenyl)methyl]thio}pyridine-3-carbonitrile as a pale yellow oil (0.041 g, 95 %). The pale yellow oil is taken up in methanol. To this is added a solution of fumaric acid (1 equiv, 0.0153 g) in methanol. This is stirred for a couple of minutes, then EtOAc followed by cyclohexane added. The resulting precipitate is collected by filtration to give

the fumarate salt of 2-[(*S*)-(2*S*)-morpholin-2-yl(phenyl)methyl]thio}pyridine-3-carbonitrile (1:1 fumarate salt) as a white solid (0.042 g). LCMS 12 min gradient method, $R_t = 4.6$ min, $(M+H^+) = 312$

5 **Example 14G: (2*S*)-2-[phenyl(pyridin-2-ylthio)methyl]morpholine hydrochloride**




i) To a stirred solution of (*R*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methyl methanesulfonate (0.70 g, 1.94 mmole) and 2-mercaptopyridine (0.54 g, 4.84 mmole) in anhydrous DMF, at room temperature and under nitrogen, is added potassium carbonate (0.80 g, 5.81 mmole). The reaction is left to stir at room temperature for 6 days. The reaction mixture is diluted with methanol and loaded onto an SC10-2 column preconditioned with MeOH. The column is washed with MeOH then basic material is eluted with 2 N NH_3 /methanol. This ammonia solution is concentrated *in vacuo* to give an orange residue (0.881 g). Purification by automated flash chromatography (ISCO System, 0 – 30 % ethyl acetate in *isohexane* gradient elution over 30 minutes) gave (2*S*)-2-[phenyl(pyridin-2-ylthio)methyl]-4-(phenylmethyl)morpholine as a colourless oil (0.245 g, 34 %). LCMS 6 min gradient method, $R_t = 2.7$ min, $(M+H^+) = 377$.

ii) Deprotection of the morpholine nitrogen is carried out using the method and work up as described in Example 1G, using polymer supported *diisopropylamine* (3.78 mmole/g, 0.43 g, 1.64 mmole), (2*S*)-2-[phenyl(pyridin-2-ylthio)methyl]-4-(phenylmethyl)morpholine (0.103g, 0.274 mmole), dry DCM (10 ml), 1-chloroethyl chloroformate (0.15 ml, 1.37 mmole) and methanol (10 ml). This gave a pale yellow oil (0.058 g, 74 %). Purification of this residue by automated flash chromatography (ISCO System, SiO_2 Redisep column, 10 % MeOH in DCM) gave a colourless oil (0.044 g, 54 %). This oil is taken up in ethyl acetate. To this is added a solution of hydrochloric acid in dioxane (4 M, 0.1 ml). Concentration *in vacuo* gave the hydrochloride salt of (2*S*)-2-

[phenyl(pyridin-2-ylthio)methyl] as a white solid (0.045 g). LCMS 6 min gradient method, $R_t = 1.8$ min, $(M+H^+) = 287$

Example 15G: (2*S*)-2-[(*S*)-[(3-iodopyridin-2-yl)thio](phenyl)methyl]morpholine

5 fumarate



- i) To (*S*)-phenyl[(2*S*)-4-(phenylmethyl)morpholin-2-yl]methanethiol (6) (0.50 g, 1.67 mmole) and 2-chloro-3-iodopyridine (0.48 g, 2.00 mmole) in degassed DMF (3 ml) is added cesium fluoride (0.38 g, 2.50 mmole) at room temperature and under nitrogen. The mixture is heated at between 55-75°C for 3 days. The organic layer is then loaded directly onto a 35 g ISCO column (SiO₂) and columned using automated flash chromatography (0 – 30% EtOAc in cyclohexane over 30 minutes) to give a pale yellow crystalline solid (0.55 g). The solid is taken up in DCM:MeOH (1:1) and loaded onto an SC10-2 column (10 g) preconditioned with MeOH. The column is washed with MeOH to remove 2-chloro-3-iodopyridine, then more basic material is eluted with 2 N NH₃/methanol. The ammonia solution is concentrated *in vacuo* to give (2*S*)-2-[(*S*)-[(3-iodopyridin-2-yl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine as a pale yellow solid (0.19 g, 23%). LCMS 6 min gradient method, $R_t = 3.8$ min, $(M+H^+) = 503$
- ii) To a suspension of polymer supported diisopropylamine (3.72 mmol/g, 0.285 g, 1.06 mmole) and (2*S*)-2-[(*S*)-[(3-iodopyridin-2-yl)thio](phenyl)methyl]-4-(phenylmethyl)morpholine (0.107 g, 0.21 mmole) in dry DCM (1.5 ml) is added 1-chloroethyl chloroformate (0.116 ml, 1.06 mmole) at room temperature and under nitrogen. The mixture is heated at 40°C for 2 hours. The reaction mixture is then filtered and concentrated *in vacuo* to give a pale orange liquid. This is taken up in methanol (1.5 ml) and the solution left to stir at room temperature overnight. After stirring overnight at room temperature, the reaction mixture is loaded directly onto an SC10-2 column. The SC10-2 column is washed with methanol, then more basic material is eluted with 2 N

NH₃/methanol. The ammonia solution is concentrated *in vacuo* to give (2*S*)-2-[(*S*)-[(3-iodopyridin-2-yl)thio](phenyl)methyl]morpholine as a pale yellow oil (0.047 g, 53%). This oil is taken up in methanol and to this is added a solution of fumaric acid (1 equiv, 0.013 g) in methanol. This is stirred for a couple of minutes, then EtOAc followed by
5 Et₂O added. The resulting precipitate is collected by filtration to give the fumarate salt of (2*S*)-2-[(*S*)-[(3-iodopyridin-2-yl)thio](phenyl)methyl]morpholine (1:1 fumarate salt) as a white solid (0.036 g). LCMS 12 min gradient method, Rt = 4.9 min, (M+H⁺) = 413

10 The pharmacological profile of the compounds of Formulae (IA), (IB), (IC), (ID), (IE), (IF) and (IG) can be demonstrated as follows. The preferred exemplified compounds above exhibit a K_i value less than 500nM at the norepinephrine transporter as determined using the scintillation proximity assay described below. Furthermore, the preferred exemplified compounds above selectively inhibit the norepinephrine transporter
15 relative to the serotonin and dopamine transporters by a factor of at least five using the scintillation proximity assays as described below.

Generation of stable cell-lines expressing the human dopamine, norepinephrine and serotonin transporters

20 Standard molecular cloning techniques are used to generate stable cell-lines expressing the human dopamine, norepinephrine, and serotonin transporters. The polymerase chain reaction (PCR) was used in order to isolate and amplify each of the three full-length cDNAs from an appropriate cDNA library. Primers for PCR were designed using the following published sequence data:

25

Human dopamine transporter: GenBank M95167. Reference: Vandenberg DJ, Persico AM and Uhl GR. A human dopamine transporter cDNA predicts reduced glycosylation, displays a novel repetitive element and provides racially-dimorphic TaqI RFLPs. *Molecular Brain Research* (1992) Volume 15, pages 161-166.

30

Human norepinephrine transporter: GenBank M65105. Reference: Pacholczyk T, Blakely, RD and Amara SG. Expression cloning of a cocaine- and antidepressant-sensitive human noradrenaline transporter. *Nature* (1991) Volume 350, pages 350-354.

5 Human serotonin transporter: GenBank L05568. Reference: Ramamoorthy S, Bauman AL, Moore KR, Han H, Yang-Feng T, Chang AS, Ganapathy V and Blakely RD. Antidepressant- and cocaine-sensitive human serotonin transporter: Molecular cloning, expression, and chromosomal localization. *Proceedings of the National Academy of Sciences of the USA* (1993) Volume 90, pages 2542-2546.

10

The PCR products are cloned into a mammalian expression vector (e.g., pcDNA3.1 (Invitrogen)) using standard ligation techniques. The constructs are then used to stably transfect HEK293 cells using a commercially available lipofection reagent (LipofectamineTM – Invitrogen) following the manufacturer's protocol.

15

Scintillation proximity assays for determining the affinity of test ligands at the norepinephrine transporter

The compounds of Formulae (II) and (III) of the present invention are norepinephrine reuptake inhibitors, and possess excellent activity in, for example, a
20 scintillation proximity assay (e.g., J. Gobel, D.L. Saussy and A. Goetz, *J. Pharmacol. Toxicol.* (1999) 42:237-244). Thus, ³H-nisoxetine binding to norepinephrine re-uptake sites in a cell line transfected with DNA encoding human norepinephrine transporter binding protein has been used to determine the affinity of ligands at the norepinephrine transporter.

25

Membrane Preparation:

Cell pastes from large scale production of HEK-293 cells expressing cloned human norepinephrine transporters were homogenized in 4 volumes 50mM Tris-HCl containing 300mM NaCl and 5mM KCl, pH 7.4. The homogenate was centrifuged twice
30 (40,000g, 10min, 4°C) with pellet re-suspension in 4 volumes of Tris-HCl buffer containing the above reagents after the first spin and 8 volumes after the second spin.

The suspended homogenate was centrifuged (100g, 10min, 4°C) and the supernatant kept and re-centrifuged (40,000g, 20min, 4°C). The pellet was resuspended in Tris-HCl buffer containing the above reagents along with 10%w/v sucrose and 0.1mM phenylmethylsulfonyl fluoride (PMSF). The membrane preparation was stored in aliquots (1ml) at -80°C until required. The protein concentration of the membrane preparation was determined using a bicinchoninic acid (BCA) protein assay reagent kit (available from Pierce).

[³H]-Nisoxetine Binding Assay:

- Each well of a 96 well microtitre plate was set up to contain the following:
- 50µl 2nM [N-methyl-³H]-Nisoxetine hydrochloride (70-87Ci/mmol, from NEN Life Science Products)
 - 75µl Assay buffer (50mM Tris-HCl pH 7.4 containing 300mM NaCl and 5mM KCl)
 - 25µl Test compound, assay buffer (total binding) or 10µM Desipramine HCl (non-specific binding)
 - 50µl Wheatgerm agglutinin coated poly (vinyltoluene) (WGA PVT) SPA Beads (Amersham Biosciences RPNQ0001) (10mg/ml)
 - 50µl Membrane (0.2mg protein per ml)

The microtitre plates were incubated at room temperature for 10 hours prior to reading in a Trilux scintillation counter. The results were analysed using an automatic spline fitting programme (Multicalc, Packard, Milton Keynes, UK) to provide Ki values for each of the test compounds.

Serotonin Binding Assay

- The ability of a test compound to compete with [³H]-citalopram for its binding sites on cloned human serotonin transporter containing membranes has been used as a measure of test compound ability to block serotonin uptake via its specific transporter (Ramamoorthy, S., Giovanetti, E., Qian, Y., Blakely, R., (1998) *J. Biol. Chem.* 273: 2458).

Membrane Preparation:

Membrane preparation is essentially similar to that for the norepinephrine transporter containing membranes as described above. The membrane preparation was stored in aliquots (1 ml) at -70°C until required. The protein concentration of the membrane preparation was determined using a BCA protein assay reagent kit.

5

[^3H]-Citalopram Binding Assay:

Each well of a 96 well microtitre plate was set up to contain the following:

- 50 μl 2 nM [^3H]-Citalopram (60-86 Ci/mmol, Amersham Biosciences)
- 75 μl Assay buffer (50 mM Tris-HCl pH 7.4 containing 150 mM NaCl and 5 mM KCl)
- 10 25 μl Diluted compound, assay buffer (total binding) or 100 μM Fluoxetine (non-specific binding)
- 50 μl WGA PVT SPA Beads (40 mg/ml)
- 50 μl Membrane preparation (0.4 mg protein per ml)

15 The microtitre plates were incubated at room temperature for 10 hours prior to reading in a Trilux scintillation counter. The results were analysed using an automatic spline fitting programme (Multicalc, Packard, Milton Keynes, UK) to provide K_i (nM) values for each of the test compounds.

Dopamine Binding Assay

20 The ability of a test compound to compete with [^3H]-WIN35,428 for its binding sites on human cell membranes containing cloned human dopamine transporter has been used as a measure of the ability of such test compounds to block dopamine uptake via its specific transporter (Ramamoorthy et al 1998 *supra*).

Membrane Preparation:

25 Is essentially the same as for membranes containing cloned human serotonin transporter as described above.

[^3H]-WIN35,428 Binding Assay:

- 30 Each well of a 96 well microtitre plate was set up to contain the following:
- 50 μl 4 nM [^3H]-WIN35,428 (84-87 Ci/mmol, from NEN Life Science Products)

- 75µl Assay buffer (50mM Tris-HCl pH 7.4 containing 150mM NaCl and 5mM KCl)
25µl Diluted compound, assay buffer (total binding) or 100µM Nomifensine (non-specific binding)
50µl WGA PVT SPA Beads (10mg/ml)
5 50µl Membrane preparation (0.2mg protein per ml.)

The microtitre plates were incubated at room temperature for 120 minutes prior to reading in a Trilux scintillation counter. The results were analysed using an automatic spline fitting programme (Multicalc, Packard, Milton Keynes, UK) to provide K_i values for each of the test compounds.

10

Acid Stability

- The acid stability of a compound according to the present invention was determined as a solution in buffer at 6 different pH values (HCl 0.1N, pH 2, pH 4, pH 6, pH 7, and pH 8) at 40°C over a time course of 72 hours. Samples were taken at the
15 beginning of the study and after 3, 6 and 24 hours and analysed by capillary electrophoresis. The original sample used in this study contained 0.8% of the undesired epimer as internal standard. The samples taken at the different time points during the study did not show any significant change in the percentage of the undesired epimer. This assay confirms that compounds of the present invention are chemically and
20 configurationally stable under acidic conditions.

In Vitro Determination of the Interaction of compounds with CYP2D6 in Human Hepatic Microsomes

- Cytochrome P450 2D6 (CYP2D6) is a mammalian enzyme which is commonly
25 associated with the metabolism of around 30% of pharmaceutical compounds. Moreover, this enzyme exhibits genetic polymorphism, resulting in the presence of both normal and poor metabolizers in the population. A low involvement of CYP2D6 in the metabolism of compounds (i.e. the compound being a poor substrate of CYP2D6) is desirable in order to reduce any variability from subject to subject in the pharmacokinetics of the compound.
30 Also, compounds with a low inhibitor potential for CYP2D6 are desirable in order to avoid drug-drug interactions with co-administered drugs that are substrates of CYP2D6.

Compounds can be tested both as substrates and as inhibitors of this enzyme by means of the following assays.

CYP2D6 substrate assay

Principle:

This assay determines the extent of the CYP2D6 enzyme involvement in the total oxidative metabolism of a compound in microsomes. Preferred compounds of the present invention exhibit less than 75% total metabolism via the CYP2D6 pathway.

For this in vitro assay, the extent of oxidative metabolism in human liver microsomes (HLM) is determined after a 30 minute incubation in the absence and presence of Quinidine, a specific chemical inhibitor of CYP2D6. The difference in the extent of metabolism in absence and presence of the inhibitor indicates the involvement of CYP2D6 in the metabolism of the compound.

Materials and Methods:

Human liver microsomes (mixture of 20 different donors, mixed gender) were acquired from Human Biologics (Scottsdale, AZ, USA). Quinidine and β -NADPH (β -Nicotinamide Adenine Dinucleotide Phosphate, reduced form, tetrasodium salt) were purchased from Sigma (St Louis, MO, USA). All the other reagents and solvents were of analytical grade. A stock solution of the new chemical entity (NCE) was prepared in a mixture of Acetonitrile/Water to reach a final concentration of acetonitrile in the incubation below 0.5%.

The microsomal incubation mixture (total volume 0.1 mL) contained the NCE (4 μ M), β -NADPH (1 mM), microsomal proteins (0.5 mg/mL), and Quinidine (0 or 2 μ M) in 100 mM sodium phosphate buffer pH 7.4. The mixture was incubated for 30 minutes at 37 °C in a shaking waterbath. The reaction was terminated by the addition of acetonitrile (75 μ L). The samples were vortexed and the denaturated proteins were removed by centrifugation. The amount of NCE in the supernatant was analyzed by liquid chromatography /mass spectrometry (LC/MS) after addition of an internal standard. A sample was also taken at the start of the incubation (t=0), and analysed similarly.

Analysis of the NCE was performed by liquid chromatography /mass spectrometry. Ten μL of diluted samples (20 fold dilution in the mobile phase) were injected onto a Spherisorb CN Column, 5 μM and 2.1 mm x 100 mm (Waters corp. Milford, MA, USA). The mobile phase consisting of a mixture of Solvent A/Solvent B, 30/70 (v/v) was pumped (Alliance 2795, Waters corp. Milford, MA, USA) through the column at a flow rate of 0.2 ml/minute. Solvent A and Solvent B were a mixture of ammonium formate 5.10^{-3} M pH 4.5/ methanol in the proportions 95/5 (v/v) and 10/90 (v/v), for solvent A and solvent B, respectively. The NCE and the internal standard were quantified by monitoring their molecular ion using a mass spectrometer ZMD or ZQ (Waters-Micromass corp, Machester, UK) operated in a positive electrospray ionisation.

The extent of CYP2D6 involvement (% of CYP2D6 involvement) was calculated comparing the extent of metabolism in absence and in presence of quinidine in the incubation.

The extent of metabolism without inhibitor (%) was calculated as follows:

$$\frac{(\text{NCE response in samples without inhibitor})_{\text{time 0}} - (\text{NCE response in samples without inhibitor})_{\text{time 30}}}{(\text{NCE response in samples without inhibitor})_{\text{time 0}}} \times 100$$

The extent of metabolism with inhibitor (%) was calculated as follows:

$$\frac{(\text{NCE response in samples without inhibitor})_{\text{time 0}} - (\text{NCE response in samples with inhibitor})_{\text{time 30}}}{(\text{NCE response in samples without inhibitor})_{\text{time 0}}} \times 100$$

where the NCE response is the area of the NCE divided by the area of the internal standard in the LC/MS analysis chromatogram, time0 and time30 correspond to the 0 and 30 minutes incubation time.

The % of CYP2D6 involvement was calculated as follows :

$$\frac{(\% \text{ extent of metabolism without inhibitor}) - (\% \text{ extent of metabolism with inhibitor})}{\% \text{ extent of metabolism without inhibitor}} \times 100$$

CYP2D6 inhibitor assay

Principle:

The CYP2D6 inhibitor assay evaluates the potential for a compound to inhibit CYP2D6. This is performed by the measurement of the inhibition of the bufuralol 1'-

hydroxylase activity by the compound compared to a control. The 1'-hydroxylation of bufuralol is a metabolic reaction specific to CYP2D6. Preferred compounds of the present invention exhibit an IC_{50} higher than 6 μ M for CYP2D6 activity, the IC_{50} being the concentration of the compound that gives 50 % of inhibition of the CYP2D6 activity.

5

Materials and Methods:

Human liver microsomes (mixture of 20 different donors, mixed gender) were acquired from Human Biologics (Scottsdale, AZ). β -NADPH was purchased from Sigma (St Louis, MO). Bufuralol was purchased from Ultrafine (Manchester, UK). All the other reagents and solvents were of analytical grade.

Microsomal incubation mixture (total volume 0.1 mL) contained bufuralol 10 μ M, β -NADPH (2 mM), microsomal proteins (0.5 mg/mL), and the new chemical entity (NCE) (0, 5, and 25 μ M) in 100 mM sodium phosphate buffer pH 7.4. The mixture was incubated in a shaking waterbath at 37 °C for 5 minutes. The reaction was terminated by the addition of methanol (75 μ L). The samples were vortexed and the denaturated proteins were removed by centrifugation. The supernatant was analyzed by liquid chromatography connected to a fluorescence detector. The formation of the 1'-hydroxybufuralol was monitored in control samples (0 μ M NCE) and in the samples incubated in presence of the NCE. The stock solution of NCE was prepared in a mixture of Acetonitrile/Water to reach a final concentration of acetonitrile in the incubation below 1.0%.

The determination of 1'-hydroxybufuralol in the samples was performed by liquid chromatography with fluorimetric detection as described below. Twenty five μ L samples were injected onto a Chromolith Performance RP-18e column (100 mm x 4.6 mm) (Merck KGaA, Darmstadt, Germany). The mobile phase, consisting of a mixture of solvent A and solvent B whose the proportions changed according the following linear gradient, was pumped through the column at a flow rate of 1 ml/min:

Time (minutes)	Solvent A (%)	Solvent B (%)
0	65	35
2.0	65	35

2.5	0	100
5.5	0	100
6.0	65	35

Solvent A and Solvent B consisted of a mixture of 0.02 M potassium dihydrogenophosphate buffer pH3/ methanol in the proportion 90/10 (v/v) for solvent A and 10/90 (v/v) for solvent B. The run time was 7.5 minutes. Formation of 1'-

5 hydroxybufuralol was monitored by fluorimetric detection with extinction at λ 252 nm and emission at λ 302 nm.

The IC_{50} of the NCE for CYP2D6 was calculated by the measurement of the percent of inhibition of the formation of the 1'-hydroxybufuralol in presence of the NCE compared to control samples (no NCE) at a known concentration of the NCE.

The percent of inhibition of the formation of the 1'-hydroxybufuralol is calculated as follows:

$$\frac{(\text{1'-hydroxybufuralol formed without inhibitor}) - (\text{1'-hydroxybufuralol formed with inhibitor})}{(\text{1'-hydroxybufuralol area formed without inhibitor})} \times 100$$

15 The IC_{50} is calculated from the percent inhibition of the formation of the 1'-hydroxybufuralol as follows (assuming competitive inhibition):

$$\frac{\text{NCE Concentration} \times (100 - \text{Percent of inhibition})}{\text{Percent of inhibition}}$$

20 The IC_{50} estimation is assumed valid if inhibition is between 20% and 80% (Moody GC, Griffin SJ, Mather AN, McGinnity DF, Riley RJ. (1999) Fully automated analysis of activities catalyzed by the major human liver cytochrome P450 (CYP) enzymes: assessment of human CYP inhibition potential. *Xenobiotica* 29(1): 53-75).

25 The invention being thus described, it is obvious that the same can be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the present invention, and all such modifications as would be obvious to one skilled in the art are intended to be included within the scope of the following claims.